

Generalization bounds for regularized portfolio selection with market side information

T. Bazier-Matte,
E. Delage

G-2018-77

October 2018

La collection *Les Cahiers du GERAD* est constituée des travaux de recherche menés par nos membres. La plupart de ces documents de travail a été soumis à des revues avec comité de révision. Lorsqu'un document est accepté et publié, le pdf original est retiré si c'est nécessaire et un lien vers l'article publié est ajouté.

Citation suggérée: T. Bazier-Matte, E. Delage (Octobre 2018). Generalization bounds for regularized portfolio selection with market side information, Rapport technique, Les Cahiers du GERAD G-2018-77, GERAD, HEC Montréal, Canada.

Avant de citer ce rapport technique, veuillez visiter notre site Web (<https://www.gerad.ca/fr/papers/G-2018-77>) afin de mettre à jour vos données de référence, s'il a été publié dans une revue scientifique.

The series *Les Cahiers du GERAD* consists of working papers carried out by our members. Most of these pre-prints have been submitted to peer-reviewed journals. When accepted and published, if necessary, the original pdf is removed and a link to the published article is added.

Suggested citation: T. Bazier-Matte, E. Delage (October 2018). Generalization bounds for regularized portfolio selection with market side information, Technical report, Les Cahiers du GERAD G-2018-77, GERAD, HEC Montréal, Canada.

Before citing this technical report, please visit our website (<https://www.gerad.ca/en/papers/G-2018-77>) to update your reference data, if it has been published in a scientific journal.

La publication de ces rapports de recherche est rendue possible grâce au soutien de HEC Montréal, Polytechnique Montréal, Université McGill, Université du Québec à Montréal, ainsi que du Fonds de recherche du Québec – Nature et technologies.

Dépôt légal – Bibliothèque et Archives nationales du Québec, 2018
– Bibliothèque et Archives Canada, 2018

The publication of these research reports is made possible thanks to the support of HEC Montréal, Polytechnique Montréal, McGill University, Université du Québec à Montréal, as well as the Fonds de recherche du Québec – Nature et technologies.

Legal deposit – Bibliothèque et Archives nationales du Québec, 2018
– Library and Archives Canada, 2018

GERAD HEC Montréal
3000, chemin de la Côte-Sainte-Catherine
Montréal (Québec) Canada H3T 2A7

Tél. : 514 340-6053
Télec. : 514 340-5665
info@gerad.ca
www.gerad.ca

Generalization bounds for regularized portfolio selection with market side information

Thierry Bazier-Matte ^a

Erick Delage ^{b,c}

^a Caisse de dépôt et placement du Québec (Québec)
Canada, H2Z 2B3

^b GERAD, Montréal (Québec), Canada, H3T 2A7

^c HEC Montréal, Montréal (Québec), Canada,
H3T 2A7

erick.delage@hec.ca

October 2018
Les Cahiers du GERAD
G–2018–77

Copyright © 2018 GERAD, Bazier-Matte, Delage

Les textes publiés dans la série des rapports de recherche *Les Cahiers du GERAD* n'engagent que la responsabilité de leurs auteurs. Les auteurs conservent leur droit d'auteur et leurs droits moraux sur leurs publications et les utilisateurs s'engagent à reconnaître et respecter les exigences légales associées à ces droits. Ainsi, les utilisateurs:

- Peuvent télécharger et imprimer une copie de toute publication du portail public aux fins d'étude ou de recherche privée;
- Ne peuvent pas distribuer le matériel ou l'utiliser pour une activité à but lucratif ou pour un gain commercial;
- Peuvent distribuer gratuitement l'URL identifiant la publication.

Si vous pensez que ce document enfreint le droit d'auteur, contactez-nous en fournissant des détails. Nous supprimerons immédiatement l'accès au travail et enquêterons sur votre demande.

The authors are exclusively responsible for the content of their research papers published in the series *Les Cahiers du GERAD*. Copyright and moral rights for the publications are retained by the authors and the users must commit themselves to recognize and abide the legal requirements associated with these rights. Thus, users:

- May download and print one copy of any publication from the public portal for the purpose of private study or research;
- May not further distribute the material or use it for any profit-making activity or commercial gain;
- May freely distribute the URL identifying the publication.

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract: Drawing on statistical learning theory, we derive out-of-sample and optimality guarantees about the investment strategy obtained from a regularized portfolio optimization model which attempts to exploit side information about the financial market in order to reach an optimal risk-return tradeoff. This side information might include for instance recent stock returns, volatility indexes, financial news indicators, etc. In particular, we demonstrate that an investment policy that linearly combines this side information in a way that is optimal from the perspective of a random sample set is guaranteed to perform also relatively well (*i.e.*, within a perturbing factor of $O(1/\sqrt{n})$) with respect to the unknown distribution that generated this sample set. We also demonstrate that these performance guarantee are lost in a high dimensional regime where the size of the side information vector is of an order that is comparable to the sample size. We finally extend these results to the case where non-linear investment policies are considered using a kernel operator and show that with radial basis function kernels the performance guarantees become insensitive to how much side information is used.

Keywords: Portfolio optimization, Generalization bound, Utility maximization, learning theory

Acknowledgments: E. Delage gratefully acknowledges the support of the Canadian Natural Sciences and Engineering Research Council [Grant 386416-2010].

1 Introduction

There is no doubt that modern portfolio management theory has been dramatically affected by two important historical events. First, Markowitz in 1952 highlighted in his seminal paper Markowitz (1952) how investment decisions needed to inherently trade-off between risk (typically measured using variance) and returns (in the form of expected returns). This was later reinterpreted as a special case of characterizing risk aversion using expected utility theory von Neumann and Morgenstern (1944). The flexibility of such a theory has since then been demonstrated in many occasions regarding the wide diversity of investors' risk aversion that it can represent (see Ingersoll (1987) and reference therein for an overview of the type of attitudes that can be modeled).

The second turning point of this theory can be considered to have occurred with the financial crisis of 2008 which provided strong evidence that the use of statistics such as variance and value-at-risk, and of distribution models that are calibrated using historical data could provide a false sense of security Salmon (2009). In an attempt to address some of these new challenges, researchers have proposed using more robust statistical estimators Madan et al. (1998); Goldfarb and Iyengar (2003); Olivares-Nadal and DeMiguel (2018) while others encouraged the use of robust portfolio management models that are designed to produce out-of-samples guarantees by exploiting the use of a confidence region for the distribution of future returns Delage and Ye (2010); Huang et al. (2010); Esfahani and Kuhn (2017); Bertsimas and Van Parys (2017).

In this work, we draw on statistical learning theory to establish what are the out-of-sample guarantees that can be obtained when using regularization in an expected utility model that allows to exploit side information about the financial markets (see Brandt et al. (2009) where non-regularized version was introduced). This side information could consist of fundamental analysis (as was famously done in Fama and French (1993)), but also of technical analysis, financial news, etc. Overall, we consider our contribution to be four-fold.

1. We derive a lower bound on the out-of-sample performance of the investment strategy returned by this regularized model. In this respect, our results differ from the usual statistical learning and stability theory results in the sense that our guarantees will not be in terms of quality of fit of a model (*e.g.*, expected squared loss, hinge loss, etc.), but rather in terms of the actual performance perceived by the investor (through the notion of a certainty equivalent).
2. We derive an upper bound on the suboptimality of the investment strategy when compared to the optimal strategy that would be derived using the full knowledge of the sample distribution. Note that such finite sample guarantees have not been established for data-driven or distributionally robust optimization.
3. Considering that nowadays a growing amount of side-information can be exploited by individuals to make their investments, we establish precisely how these bounds are affected at a high-dimensional (or "big data") regime.
4. Finally, we present how the out-of-sample and sub-optimality bounds can be extended to a multi-asset portfolio selection problem and a "kernelized" single-asset portfolio selection problem which can produce investment strategies that are linear functions of a lifting of the side-information to a possibly infinite dimensional space.

It is worth mentioning that contributions 1–3 are similar in spirit to those of Rudin and Vahn (2015) who applied stability theory to provide generalization bounds for a newsvendor problem. There are however a number of distinctions regarding how stability theory needs to be articulated for the two applications. For example, our paper deals with a more general performance function which is non-linear and possibly unbounded on both sides, and we focus on generalization and sub-optimality bounds which, in our opinion, are more informative than the measure that is bounded in Rudin and Vahn (2015). Moreover, we attempt to be more precise in our analysis on characterizing the effect of data dimensionality on the out-of-sample performance. To the best of our knowledge, Contribution 4 is entirely original given that the work of Rudin and Vahn (2015) did not consider a multi-dimensional decision vector and that it used kernels in the context of kernel density estimator unlike in this work where a kernel operator will be used to define the space of possible investment strategies. In the field of finance, one can also identify some interesting applications of kernels (*e.g.* in Györfi et al. (2006) and Takano and Gotoh (2014)) but again none of these applications provide out-of-sample guarantees for the use of a kernel operator in a single-stage investment problem.

The rest of the paper is divided as follows. First, we formally introduce our model and assumptions in Section 2. Section 3 then presents what kind of out-of-sample guarantees can be provided on the certainty equivalent (CE) of the investor using a sample of market returns and side information when assuming a stationary market distribution. We then proceed in Section 4 to show that the same kind of guarantees can also be derived for the CE suboptimality, before showing in Section 5 what kind of behaviour can be expected in “big-data” situation. We then present extensions of our results to the case of multiple risky assets in Section 6 and to the case where investment strategies are defined using kernels in Section 7. Finally, we conclude in Section 8. All proofs have been pushed to the appendix section.

2 Model and assumptions

We consider a classical financial portfolio selection problem involving a risky asset with random return rate R and a risk-free asset with return rate of 0% for simplicity of exposure. We also suppose that the investor’s risk aversion can be characterized using expected utility theory using a strictly increasing concave utility function u , and that the investor has access to side information regarding the returns. This information might be the result of processing the most recent financial or economic news, etc. We let this information be described as a vector of p normalized random features $X \in \mathbb{R}^p$. In this context, if the the distribution F of the pair (X, R) of side information and return is known, a linear investment policy that exploits the side information optimally for this investor can be obtained by solving the following optimization problem:

$$\underset{q \in \mathbb{R}^p}{\text{maximize}} \quad E_F[u(R \cdot q^T X)], \quad (1)$$

where an investment policy consists of investing a $q^T X$ proportion of the wealth in the risky asset and a $1 - q^T X$ proportion in the risk-free one, and where it is assumed that short-selling is permitted.¹

In practice however, the exact distribution describing the relation between X and R is not available at the time of designing the investment policy and one might instead need to exploit a sample set $\mathcal{S}_n := \{(x_i, r_i)\}_{i=1}^n$ where each (x_i, r_i) was drawn independently and identically from F . Unfortunately, when the sample size n is relatively small compared to p , it is well known that the version of problem (1) that uses the empirical distribution \hat{F} obtained from sample \mathcal{S}_n can suffer from overfitting the sample and produce investment policies that perform badly out of sample. This is for instance illustrated in the following example.

Example 1 Consider for instance a case where $n = p$ and each term in X is independently and identically drawn from a Gaussian distribution. Given that it is well known that the probability that the random matrix $\Xi := [X_1 \ X_2 \ \dots \ X_n]^T$ be singular is null, then one can easily establish that problem (1) with \hat{F} is unbounded. Indeed, one can verify that $r_i \bar{q}^T x_i = 1$ for all $i = 1, \dots, n$ when \bar{q} is set to $\Xi^{-1}[1/r_1 \ 1/r_2 \ \dots \ 1/r_n]^T$. Hence, one can achieve an arbitrarily large empirical expected utility by investing according to $\alpha \bar{q}$ for $\alpha > 0$.

To prevent issues associated to overfitting, one might instead seek the optimal solution of the following regularized empirical expected utility maximization problem:

$$\underset{q \in \mathbb{R}^p}{\text{maximize}} \quad E_{\hat{F}}[u(R \cdot q^T X)] + \lambda \|q\|_2^2. \quad (2)$$

We will refer to the optimal solution of this problem as \hat{q} . Note that such an optimal solution always exists given that the objective function is strongly concave.

The question remains of understanding what guarantees one has regarding out-of-sample performance of the portfolio investment policy obtained from such a regularized problem. In what follows, we establish some high confidence bounds on the out-of-sample performance and suboptimality of \hat{q} .

¹Note that in the case the risk free return rate is non-null, one should interpret R as the return in excess of the risk-free rate. Hence, the model can still take the form presented in (1) using the reduction $E_F[u((R + r_f) \cdot q^T X + r_f(1 - q^T X))] = E_F[u(R \cdot q^T X + r_f)] = E_F[v(R \cdot q^T x)]$ where $v(y) := u(y + r_f)$.

3 Out-of-sample performance bounds

In this section, we identify a high confidence bound on the out-of-sample performance of \hat{q} . In particular, since utility functions are expressed in units without any physical meaning for the investor, any guarantees derived using learning theory should be reinterpreted in terms of a guarantee on the certainty equivalent² (in percent of return) of the risky investment produced by $\hat{q}^T X$. In other words, we will be interested in bounding how different the in-sample certainty equivalent performance of \hat{q} might be compared to the out-of-sample certainty equivalent performance.

In order to shed some light on this question, we first make the following assumptions.

Assumption 1 *The random return R is supported on a bounded interval $\mathcal{S}_R \subseteq [-\bar{r}, \bar{r}]$, for some $\bar{r} \in \mathbb{R}$, such that $P(|R| \leq \bar{r}) = 1$.*

Assumption 2 *The random vector of side-information X is supported on bounded set \mathcal{S}_X such that $P(\|X\|_2 \leq \xi) = 1$ for some $\xi \in \mathbb{R}$.*

Assumption 3 *The utility function is normalized such that $u(0) = 0$ and $\lim_{r \rightarrow 0^+} u'(r) = 1$. Furthermore, it is Lipschitz continuous with a Lipschitz constant of γ , i.e., for any $r_1 \in \mathbb{R}$ and $r_2 \in \mathbb{R}$, we have that $|u(r_1) - u(r_2)| \leq \gamma|r_1 - r_2|$.*

The first assumption is relatively realistic given that one can usually assess from historical data a large enough interval of returns which could be assumed to contain R with probability one. For instance, when looking at the last 35 years of daily returns for an index such as S&P 500, this interval can legitimately be set to $[-25\%, 25\%]$ daily returns. If some side information are not known to be bounded, the second assumption might require one to pre-process the vector of side information in order to rely on the results that will be presented. This could typically be done by projecting this vector on the surface of a ball of radius ξ when $\|X\|_2 > \xi$, which is as simple as replacing X with $(\xi/\|X\|_2) \cdot X$. This assumption will be further studied in Section 5. Finally, while the last assumption is fairly common for establishing generalization bounds and can certainly accommodate any piecewise linear utility function (often used by numerical optimization methods), it is important to mention that it is not one that is commonly made in modern portfolio theory. If, for instance, an investor expresses an absolute risk aversion uniformly equal to α , this suggests the use of $u(r) := (1/\alpha)(1 - \exp(-\alpha r))$ which is not Lipschitz continuous. Fortunately, the theory that will be developed only exploits the fact that the function is Lipschitz continuous on the interval $[-\bar{r}^2 \xi^2 / (2\lambda), \bar{r}^2 \xi^2 / (2\lambda)]$.

We are now in a position to exploit a well-known learning theory result to establish a bound on the out-of-sample portfolio performance of \hat{q} based its in-sample estimation.

Theorem 1 *Given that Assumptions 1, 2 and 3 are satisfied, the certainty equivalent of the out-of-sample performance is at most $O(1/\sqrt{n})$ worse than the in-sample one. Specifically,*

$$\text{CE}(\hat{q}; F) \geq \text{CE}(\hat{q}; \hat{F}) - \Omega_1 / \lim_{\epsilon \rightarrow 0^-} u'(\text{CE}(\hat{q}; \hat{F}) + \epsilon),$$

where

$$\text{CE}(\hat{q}; F) := u^{-1}(E_F[u(R \cdot \hat{q}^T X)]),$$

and where

$$\Omega_1 := \frac{\bar{r}^2 \xi^2}{2\lambda} \left(\frac{\gamma^2}{n} + \frac{(2\gamma^2 + \gamma + 1)\sqrt{\log(1/\delta)}}{\sqrt{2n}} \right)$$

with probability $1 - \delta$.

Our proof of Theorem 1 proceeds as follow. First, borrowing from the terminology introduced by Bousquet and Elisseeff (2002), we show that this so-called “investment algorithm” is β -stable. We then show that for

²The fact that c is the certainty equivalent of a random return R implies that the investors are indifferent between being exposed to the risk of R or getting involved in a risk free investment that has a return rate of c .

this investment algorithm the amount of utility generated from exploiting different sample sets is within a range Δ . Given that these two conditions are satisfied, we can then rely on an adapted version of Bousquet-Ellissef's out-sample error bound theorem in order to establish out-of-sample guarantees in terms of expected utility. By exploiting the concavity of $u(\cdot)$, we are finally able to describe the implications in terms of certainty equivalent that are expressed in our theorem.

It is worth noting that there exists strong connections between the use of regularization and the principles of robust optimization (e.g. in Caramanis et al. (2012)). It is therefore not surprising that through regularization we are able obtain out-of-sample guarantees that are similar in nature as those obtained when using robust optimization (see for instance the work in Esfahani and Kuhn (2017)). It would furthermore be interesting to establish what are the bounds that could be obtained for problem 1 using the results presented in Shafieezadeh-Abadeh et al. (2017) which was made public during the later stages of the writing of this paper. On the other hand, to the best of our knowledge there has still been no results in the field of robust optimization regarding finite sample bounds on the sub-optimality of solutions obtained through the robustification process. This is what we do next for problem (1) when robustification is obtained using regularization.

4 Suboptimality performance bounds

We now turn our attention to the suboptimality of the problem, *i.e.*, we would like to understand the behaviour of the performance of the empirical investment policy \hat{q} compared to the optimal policy $q^* := \arg \max_q E_F[u(R \cdot q^T X)]$. It is important to realize that in general, there are situations in which the optimal performance according to (1) could be unbounded. Thus, if one wishes to establish a bound on the sub-optimality of an investment policy, it is necessary to impose additional assumptions on the class of problem that he is facing. The two following examples motivate these assumptions.

Example 2 Consider a risk neutral investor, *i.e.*, such that $u(r) = r$ and suppose $EX_i = 0$. The expected utility simply becomes

$$E_F[u(R \cdot q^T X)] = \sum_{i=1}^n q_i \text{Cov}_F(R, X_i).$$

If we simply let $\bar{q}_i = \text{Cov}_F(R, X_i)$, it follows immediately that the expected utility of $\alpha \bar{q}$ can become arbitrarily large as α goes to infinity.

Example 3 Consider another example in which there exists a j for which feature X_j induces arbitrage over F , namely that $P\{RX_j > 0\} = 1$. In such a case, if we let $\bar{q}_i = 1$ only when $i = j$ and otherwise zero, then the expected utility of $\alpha \bar{q}$ can once again take an arbitrarily large value as α goes to infinity.

Given those two examples, we now introduce two new assumptions that will ensure that problem (1) is bounded, *i.e.*, it has a finite optimal solution.

Assumption 4 The utility function is sublinear, *i.e.*, $u(r) = o(r)$.

Assumption 5 The side information X induces no linear arbitrage opportunities, that is, there exists no $q \in \mathbb{R}^p$, such that both $P\{Rq^T X < 0\} = 0$ and $P\{Rq^T x > 0\} > 0$.

In a financial context, Assumption 4 is certainly realistic since a financial investor behaviour is usually taken to be strictly risk averse for winnings, *i.e.* that there exists a possible random gain $X \geq 0$ for which $E[u(X)] \leq u(E[X])$, thus implying Assumption 4. As for Assumption 5, this notion of arbitrage relates directly to the notion of market efficiency when the side-information contained in X is considered to be publicly available. In particular the semi-strong version of market efficiency states that it should be impossible for an investor to constantly beat the market using publicly available information. See Malkiel and Fama (1970) and Fama (1991) for more details.

Theorem 2 *Given that Assumptions 1, 2, 3, 4, and 5 are satisfied, the suboptimality of the policy \hat{q} can be expressed with confidence $1 - \delta$ by*

$$CE(\hat{q}; F) \geq CE(q^*; F) - \Omega_2 / \lim_{\epsilon \rightarrow 0^-} u'(CE(\hat{q}; F) + \epsilon),$$

where

$$\Omega_2 = \lambda \|q^*\|_2^2 + \frac{8\gamma^2 \bar{r}^2 \xi^2 (32 + \log(1/\delta))}{\lambda n} + \frac{2\gamma \bar{r}^2 \xi^2 \sqrt{32 + \log(1/\delta)}}{\lambda \sqrt{n}}.$$

The first term in Ω_2 shows that, unless the regularization constant λ is brought to zero as n increases, the empirical maximization problem (2) will asymptotically converge toward a constant suboptimality bound based on the particular market distribution F and on λ . The two other terms in Ω_2 show that this bound will be reached at a $O(1/\sqrt{n})$ rate in the same fashion as with Theorem 1. Therefore, the best suboptimality performance that can be hoped to be reached is at most $-\lambda \|q^*\|_2^2 / \lim_{\epsilon \rightarrow 0^-} u'(CE(\hat{q}; F) + \epsilon)$ when λ is maintained constant. Alternatively, one could (and typically would) bring λ to zero as the size of sample set increases in order to bring the suboptimality bound to zero. In particular, this can be done by letting $\lambda = o(1/\sqrt{n})$.

5 Big data phenomenon

In this section, we question how realistic Assumption 2 is in a big data context. In particular, we expose two sets of natural conditions for the generation of the side information vector X that leads to motivating the use of a support set which diameter grows proportionally to the square root of p .

Example 4 *Consider a case where every terms of X are independent from each other, while each X_i has a mean $E[X_i] = 0$, a variance $\text{Var}[X_i] = 1$, and are supported on their respective intervals $P(X_i \in [-\nu, \nu]) = 1$ for all i . By Hoeffding's inequality, one can establish that*

$$P\left(\left|\|X\|_2^2 - \sum_{i=1}^p E[X_i^2]\right| \leq \sqrt{2p \log(\delta/2)} \nu^2\right) \geq 1 - \delta$$

so that $\|X\|_2^2 \in [p - \sqrt{2p \log(\delta/2)} \nu^2, p + \sqrt{2p \log(\delta/2)} \nu^2]$ with probability $1 - \delta$. Hence, any ball of fixed radius ξ will contain X with a probability that asymptotically converges to zero as p increases, more specifically $P(\|X\|_2^2 \leq \xi^2) \leq 2 \exp(-2p(1 - \xi^2/\sqrt{p})^2/\nu^2)$. On the other hand, this inequality somehow also prescribes that the diameter of the support \mathcal{S}_X should increase proportionally to \sqrt{p} in order to still contain X with high probability as p increases.

Example 5 *Consider a similar case as above but where the independence assumption is dropped. In this context, although we might not have as much of a strong argument to discredit the use of a constant diameter for \mathcal{S}_X , there is still a good motivation for employing a radius that grows proportionally to \sqrt{p} . Namely, if each X_i has a mean $E[X_i] = 0$ and a variance $\text{Var}[X_i] = 1$ then the random variable $Z := \|X\|_2^2$ is necessarily positive with an expected value of p . Based on Markov inequality, this implies that with probability $1 - \delta$, we have that $\|X\|_2 \leq \sqrt{p/\delta}$.*

Since we believe these two examples provide strong arguments for replacing Assumption 2 with the assumption that it is within a ball of radius $\xi\sqrt{p}$, we reformulate our previous two results as follows.

Corollary 1 *Given that Assumptions 1 and 3 are satisfied, and that $P(\|X\|_2 \leq \xi\sqrt{p}) = 1$, the certainty equivalent of the out-of-sample performance is at most $O(p/\sqrt{n})$ worse than the in-sample one. Specifically, with probability $1 - \delta$,*

$$CE(\hat{q}; F) \geq CE(\hat{q}; \hat{F}) - \Omega_3 / \lim_{\epsilon \rightarrow 0^-} u'(CE(\hat{q}; \hat{F}) + \epsilon),$$

where

$$\Omega_3 := \frac{\bar{r}^2 \xi^2 p}{2\lambda} \left(\frac{\gamma^2}{n} + \frac{(2\gamma^2 + \gamma + 1)\sqrt{\log(1/\delta)}}{\sqrt{2n}} \right).$$

Likewise, the suboptimality of the decision \hat{q} will reach a constant bound due to regularization at a rate of at most $O(p/\sqrt{n})$:

$$\text{CE}(\hat{q}; F) \geq \text{CE}(q^*; F) - \Omega_4 / \lim_{\epsilon \rightarrow 0^-} u'(CE(\hat{q}; F) + \epsilon),$$

where

$$\Omega_4 = \lambda \|q^*\|_2^2 + \frac{8\gamma^2 \bar{r}^2 p \xi^2 (32 + \log(1/\delta))}{n\lambda} + \frac{2\gamma \bar{r}^2 p \xi^2}{\lambda} \sqrt{\frac{32 + \log(1/\delta)}{n}},$$

with probability $1 - \delta$.

Note that Assumption 2 was inspired by an early version of Rudin and Vahn (2015) who also studied asymptotic properties of a regularized decision problem in its Big data regime, *i.e.*, when n and p go to infinity simultaneously. Our analysis indicate that the convergence in accuracy that is reported with such an assumption can be misleading for many problems, *e.g.*, when the features can be considered independent from each other. In particular, Corollary 1 states that asymptotic convergence in accuracy is only guaranteed to occur when $p/\lambda = o(\sqrt{n})$ and $\lambda \rightarrow 0$. For example, both the estimation error and sub-optimality converge to zero if $p = O(n^{1/4})$ and $\lambda = O(n^{1/8})$ since $p/\lambda = O(n^{3/8})$.

However, it is important to understand that Corollary 1 serves as a worst-case scenario and that we don't necessarily expect to observe downgrading performances as soon as $n = o(p^2/\lambda^2)$. Still, no matter what, there is a cost to pay in pouring more and more features into such a portfolio selection problem, and this cost is directly exhibited through ξ^2 and the loosening of the guarantees bound. One might therefore wish to be prudent when facing such high-dimension regimes.

6 Extension to multiple risky assets

In Brandt et al. (2009), the authors propose a generalization of problem (1) to a context where the portfolio can be diversified among multiple assets. Their model takes the form:

$$\underset{w \in \mathbb{R}^m, q \in \mathbb{R}^p}{\text{maximize}} \quad E_F \left[u \left(\sum_{i=1}^m R_i \cdot (w_i + q^T X_i) \right) \right], \quad (3)$$

where R_i is the return obtained from risky asset i , with $i = 1, \dots, m$, and for each risky asset $i = 1, \dots, m$, w_i captures a reference investment while $X_i \in \mathbb{R}^p$ is a random vector of side information about asset i which is used to adapt the investment to the market conditions. More generally speaking, one can consider the regularized multi-asset portfolio selection problem :

$$\underset{q \in \mathbb{R}^p}{\text{maximize}} \quad E_F [u(\mathbf{R}^T \mathbf{X} \mathbf{q})], \quad (4)$$

where $\mathbf{R} \in \mathbb{R}^m$ is the random vector of returns while $\mathbf{X} \in \mathbb{R}^m \times \mathbb{R}^p$ is a matrix containing in each of its rows the side-information that is used to make the decision about the proportion of wealth that is invested in a particular risky asset. In particular, one recovers problem (3) by composing \mathbf{X} and \mathbf{q} as follows:

$$\mathbf{X} := \begin{bmatrix} X_1^T & e_1^T \\ X_2^T & e_2^T \\ \dots & \dots \\ X_m^T & e_m^T \end{bmatrix} \in \mathbb{R}^m \times \mathbb{R}^{p+m} \quad \mathbf{q} := \begin{bmatrix} q \\ w \end{bmatrix}, \quad (5)$$

where each e_i captures the i -th column of the $m \times m$ identity matrix.

Building on the results presented in Sections 3 and 4, it is actually possible to establish generalization bounds for the solution of the regularized multi-asset portfolio selection problem. In order to do so, one first needs to adapt the two assumptions that were made about the support set of R and X to the multi-asset framework.

Assumption 6 *The random return \mathbf{R} is supported on \mathcal{S}_r , which lies inside a ball of radius \bar{r} , *i.e.* $P(\|\mathbf{R}\|_2 \leq \bar{r}) = 1$.*

Assumption 7 *The random matrix of side-information \mathbf{X} is supported on \mathcal{S}_x which lies inside a ball of radius ξ , i.e. $P(\|\mathbf{X}\|_2 \leq \xi) = 1$ where $\|\mathbf{X}\|_2$ stands for the largest singular value of \mathbf{X} .*

We now can proceed with an extension of Theorem 1 to the case of multi-asset portfolios. We refer the reader to Appendix A.3 for more details about the proof.

Theorem 3 *Given that Assumptions 6, 7 and 3 are satisfied, the certainty equivalent of the out-of-sample performance is at most $O(1/\sqrt{n})$ worse than the in-sample one. Specifically, with probability larger than $1 - \delta$ we have that*

$$\text{CE}(\hat{\mathbf{q}}; F) \geq \text{CE}(\hat{\mathbf{q}}; \hat{F}) - \Omega_1 / \lim_{\epsilon \rightarrow 0^-} u'(\text{CE}(\hat{\mathbf{q}}; \hat{F}) + \epsilon),$$

where $\text{CE}(\hat{\mathbf{q}}; F) := u^{-1}(E_F[u(\mathbf{R}^T \mathbf{X} \hat{\mathbf{q}})])$ and where F and \hat{F} refer respectively to the true and empirical joint distribution of (\mathbf{R}, \mathbf{X}) .

Similarly, in order to obtain some guarantees about the sub-optimality of $\hat{\mathbf{q}}$, it is necessary to extend our assumption that no linear arbitrage opportunities are present in the market defined through (\mathbf{X}, \mathbf{R}) . This is needed in order to ensure that problem (4) is bounded and has a bounded optimal solution.

Assumption 8 *The side information \mathbf{X} induces no linear arbitrage opportunities, i.e. that there exists no $\mathbf{q} \in \mathbb{R}^p$ such that both $P\{\mathbf{R}^T \mathbf{X} \mathbf{q} < 0\} = 0$ and $P\{\mathbf{R}^T \mathbf{X} \mathbf{q} > 0\} > 0$.*

We follow with the extension of Theorem 2 to the case of multi-asset portfolio selection.

Theorem 4 *Given that Assumptions 3, 4, 6, 7, and 8 are satisfied, the suboptimality of the policy $\hat{\mathbf{q}}$ can be expressed with confidence $1 - \delta$ by*

$$\text{CE}(\hat{\mathbf{q}}; F) \geq \text{CE}(\mathbf{q}^*; F) - \Omega_2 / \lim_{\epsilon \rightarrow 0^-} u'(CE(\hat{\mathbf{q}}; F) + \epsilon).$$

It is worth noting that while the bounds presented in Theorems 3 and 4 have exactly the same definition as in the single-asset case, i.e. they reuse the original definitions of Ω_1 and Ω_2 , they in fact rely on the new definitions of ξ and \bar{r} . In particular, if one considers the multi-asset model presented in Equation (5) (inspired from Brandt et al. (2009)) and that each $R_i \in [-\bar{r}_0, \bar{r}_0]$ and each $P(\|X_i\|_2 \leq \xi_0 \sqrt{p}) = 1$ for each $i \in \{1, 2, \dots, m\}$ as proposed in Section 5, then one could conclude that $\xi = \sqrt{m(p\xi_0^2 + 1)}$ and $\bar{r} = \bar{r}_0 \sqrt{m}$ using the argument that follows. First, with probability one we have that

$$\|\mathbf{R}\|_2^2 = \sum_{i=1}^m R_i^2 \leq m\bar{r}_0^2.$$

Also, for any \mathbf{q} such that $\|\mathbf{q}\|_2 \leq 1$ we have that

$$\|\mathbf{X} \mathbf{q}\|_2^2 = \left\| \sum_{i=1}^m X_i^T \mathbf{q} + w_i \right\|_2^2 \leq \sum_{i=1}^m \|X_i\|_2^2 \|\mathbf{q}\|_2^2 + \|e_i\|_2^2 \|w\|_2^2 = m(p\xi_0^2 + 1),$$

since $\|q\|_2^2 \leq \|\mathbf{q}\|_2^2 \leq 1$ and similarly for w .

Overall, in the multi-asset portfolio selection problem discussed in Brandt et al. (2009), we can expect that the out-of-sample performance will be at most $O(pm^2/\sqrt{n})$ worse than the in-sample one. On the other hand, the sub-optimality of $\hat{\mathbf{q}}$ will reach a constant bound due to regularization at a rate of at most $O(pm^2/\sqrt{n})$ when λ is considered constant and can be brought to zero by sizing λ appropriately. Finally, it is left for future work to confirm however whether these bounds are actually tight or not.

7 Extension to Kernel approach

As described in Hofmann et al. (2008), we let $k : \mathbb{R}^p \times \mathbb{R}^p \rightarrow \mathbb{R}$ be a positive definite kernel on the space of pairs of market side-information vectors and let this kernel be associated to a unique reproducing kernel

Hilbert space \mathcal{W} with a mapping $\Phi : \mathbb{R}^p \rightarrow \mathcal{W}$ that projects vectors of market side-information \mathbb{R}^p to \mathcal{W} . It is well known that any function w in \mathcal{W} can be characterized using $\{(\alpha_i, x_i)\}_{i=1}^m$, with each $\alpha_i \in \mathbb{R}$ and $x_i \in \mathbb{R}^p$, and some $m \in \mathbb{N}$ such that $w := \sum_{i=1}^m \alpha_i k(\cdot, x_i)$. Also, given two functions $f_1 \in \mathcal{W}$ and $f_2 \in \mathcal{W}$, we have that the inner product is defined as $\langle f_1, f_2 \rangle := \sum_{i=1}^{m_1} \sum_{j=1}^{m_2} \alpha_i^1 \alpha_j^2 k(x_i^1, x_j^2)$.

With this in hand, it is possible to consider the following kernel investment problem:

$$\underset{w \in \mathcal{W}}{\text{maximize}} \quad E_F[u(R \cdot \langle w, \Phi(X) \rangle)] . \quad (6)$$

Again, given that in practice we don't have access to the full distribution information for F but rather to a set of identically and independently generated samples $\{(x_i, r_i)\}_{i=1}^n$, in order to control over-fitting we can consider solving the regularized empirical kernel investment problem:

$$\underset{w \in \mathcal{W}}{\text{maximize}} \quad \frac{1}{n} \sum_{i=1}^n u(r_i \cdot \langle w, \Phi(x_i) \rangle) - \lambda \|w\|_2^2 . \quad (7)$$

From a practical perspective it is important to explain how problem (7) can be numerically solved, especially given that \mathcal{W} would usually be an infinite dimensional space. In particular, the following theorem shows how this problem can be reduced to a finite dimensional one.

Theorem 5 *Problem (7) is equivalent to the following finite dimensional convex optimization problem:*

$$\underset{\alpha \in \mathbb{R}^n}{\text{maximize}} \quad \frac{1}{n} \sum_{i=1}^n u(r_i \sum_{j=1}^n \alpha_j k(x_j, x_i)) - \lambda \alpha^T \mathbb{K} \alpha , \quad (8)$$

where $\mathbb{K} \in \mathbb{R}^{n \times n}$ is such that $\mathbb{K}_{ij} := k(x_i, x_j)$ and known as the Gram matrix. In particular, given an optimal solution $\hat{\alpha}$ for problem (8), one can construct an optimal portfolio policy \hat{w} for problem (7) through $\hat{w} := \sum_{j=1}^n \hat{\alpha}_j \Phi(x_j)$ for which the investment proposed under condition x takes the form $\langle \hat{w}, \Phi(x) \rangle = \sum_{j=1}^n \hat{\alpha}_j k(x_j, x)$.

Proof. This follows from the representer theorem (see Theorem 9 in Hofmann et al. (2008)) which states that the optimal solution to problem (7) always takes the form of a linear combination of the kernel expansion of the sample points. Specifically, there exists a linear combination $\alpha \in \mathbb{R}^n$ such that $\hat{w} := \sum_{j=1}^n \alpha_j \Phi(x_j)$. This implies that the optimization can be reduced to optimizing over the space of $\alpha \in \mathbb{R}^n$. When doing so, the objective function becomes:

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n u(r_i \langle \sum_{j=1}^n \alpha_j \Phi(x_j), \Phi(x_i) \rangle) - \lambda \langle \sum_{i=1}^n \alpha_i \Phi(x_i), \sum_{j=1}^n \alpha_j \Phi(x_j) \rangle \\ &= \frac{1}{n} \sum_{i=1}^n u(r_i \sum_{j=1}^n \alpha_j \langle \Phi(x_j), \Phi(x_i) \rangle) - \lambda \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j \langle \Phi(x_i), \Phi(x_j) \rangle \\ &= \frac{1}{n} \sum_{i=1}^n u(r_i \sum_{j=1}^n \alpha_j k(x_j, x_i)) - \lambda \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j k(x_i, x_j) . \end{aligned}$$

This directly leads to the objective function used in problem (8). \square

Next, in order to obtain generalization bounds on the quality of \hat{w} , we will exploit a bound on the magnitude of the norm of $\Phi(X)$ which is defined through the following assumption.

Assumption 9 *The random vector of side-information X is supported on a set \mathcal{S}_X such that $P(\|\Phi(X)\|_2^2 \leq \xi^2) P(k(X, X) \leq \xi^2) = 1$.*

It is worth providing some details about how the above assumption is affected by the choice of kernel. For instance, a polynomial kernel will consider $k(x, y) := (x^T y + c)^d$ with $c \geq 0$ and $d \in \mathbb{N}$, and therefore Assumption 9 would imply that $P(\|X\|_2 \leq \sqrt{\xi^{2/d} - c}) = 1$. Alternatively, one might instead use the popular radial basis function kernel $k(x, y) := \exp(-\|x - y\|^2 / (2\sigma^2))$, where $\|\cdot\|$ is an arbitrary norm and $\sigma > 0$ is a free parameter, for which Assumption 9 is always satisfied with $\xi = 1$ since $k(x, x) = \exp(\|x - x\|_2^2 / (2\sigma^2)) = 1$ for all values of x .

In what follows, we explain how Theorems 1 and 2 can be extended to the kernel investment problem. A summary of the key steps involved in proving these results is presented in Appendices A.5 and A.6 respectively.

Theorem 6 *Given that Assumptions 1, 3, and 9 are satisfied, the certainty equivalent of the out-of-sample performance is at most $O(1/\sqrt{n})$ worse than the in-sample one. Specifically, with probability larger than $1 - \delta$ we have that*

$$\text{CE}(\hat{w}; F) \geq \text{CE}(\hat{w}; \hat{F}) - \Omega_1 / \lim_{\epsilon \rightarrow 0^-} u'(\text{CE}(\hat{w}; \hat{F}) + \epsilon),$$

where $\text{CE}(\hat{w}; F) := u^{-1}(E_F[u(R \cdot \langle \hat{w}, \Phi(X) \rangle)])$ and where F and \hat{F} refer respectively to the true and empirical joint distribution of (R, X) .

In order to extend Theorem 2, we actually need a stronger version of the no linear arbitrage opportunity assumption. This is due by the fact that the kernel investment problem now permits the use of non-linear investment strategies. A stronger condition must therefore be imposed to ensure that problem (6) is bounded and hence sub-optimality controlled.

Assumption 10 *The side information X induces no general arbitrage opportunities, i.e. that there exists no $\mathcal{S} \subseteq \mathcal{S}_X$ such that*

$$P\{R > 0 \mid X \in \mathcal{S}\} > 0 \text{ and } P\{R < 0 \mid X \in \mathcal{S}\} = 0$$

or such that

$$P\{R < 0 \mid X \in \mathcal{S}\} > 0 \text{ and } P\{R > 0 \mid X \in \mathcal{S}\} = 0.$$

Theorem 7 *Given that Assumptions 1, 3, 4, 9, and 10 are satisfied, the suboptimality of the policy \hat{w} can be expressed with confidence $1 - \delta$ by*

$$\text{CE}(\hat{w}; F) \geq \text{CE}(w^*; F) - \Omega_2 / \lim_{\epsilon \rightarrow 0^-} u'(\text{CE}(\hat{w}; F) + \epsilon).$$

Overall, in the kernel investment problem (6), if a polynomial kernel is used and $\|X\|_2 \leq \xi_0 \sqrt{p}$ with probability one, we can expect that the out-of-sample performance will be at most $O(p^d/\sqrt{n})$ worse than the in-sample one. On the other hand, the sub-optimality of \hat{q} will reach a constant bound due to regularization at a rate of at most $O(p^d/\sqrt{n})$ when λ is considered constant and can be brought to zero by sizing λ appropriately. The guarantees become more interesting if a radial basis function kernel is used given that both convergence rates become $O(1/\sqrt{n})$ in that case and are completely unaffected by the size of p . This seems to be a strong theoretical argument to support the use of radial basis function kernels in the kernel investment problem.

8 Discussion

As a conclusion, we would like to review the main messages we hope to deliver from this paper. First off, it is possible to use side information from the market, such as market news, financial indicators, economic variables and so on in order to build portfolios of single or multiple risky assets while having guarantees on the out-of-sample performance. Second, it is also possible to obtain performance bounds on the suboptimality of the empirical decision in comparison to what might have been the best decision, given full knowledge of the market. Third, there might be a cost to pay for increasing the number of side information treated by the model when the sample size on which the decision is based on is not large enough. Finally, the guarantees have natural extensions for the case where the space of investment strategies is defined using kernel operators. In particular, it appears that radial basis function kernels become especially effective in a big data regime given that the performance guarantees for this family of kernel operators are unaffected by p .

A Appendix

A.1 Proof of Theorem 1

In this proof, we will make use of an adapted version of a theorem made famous by Bousquet and Elisseeff (2002) in the context of learning theory to analyse relevant statistical properties of the investment policy \hat{q} presented in Section 2. While this theorem discusses the use of a learning algorithm, we will rather refer to an investment algorithm which is defined next.

Definition 1 Let a investment algorithm $\pi : \mathbb{R}^{(p+1) \times n} \times \mathbb{R}^p \rightarrow \mathbb{R}$ be a procedure that generates a portfolio recommendation based on a historical sample set $\mathcal{S}_n := \{(x_i, r_i)\}_{i=1}^n$, where each $x_i \in \mathcal{S}_X$ and each $r_i \in \mathcal{S}_R$, and the current market conditions x , in other words, it produces the recommendation of investing $\pi(\mathcal{S}_n, x)$ in the risky asset and $1 - \pi(\mathcal{S}_n, x)$ in the risk-free asset.

We start by adapting Theorem 11.1 from Mohri et al. (2012) (originally found in Bousquet and Elisseeff (2002)) to our context. To do so, we need to adapt the concept of β -stability to the case of investment algorithms.

Definition 2 A investment algorithm $\pi(\cdot)$ is uniformly β -stable if for any two sample sets $\mathcal{S}_n^1 := \{(x_i^1, r_i^1)\}_{i=1}^n$ and $\mathcal{S}_n^2 := \{(x_i^2, r_i^2)\}_{i=1}^n$ that are exactly identical except for the j -th sample, i.e., $(x_i^1, r_i^1) = (x_i^2, r_i^2)$ for all $i \neq j$, the following holds:

$$|u(r \pi(\mathcal{S}_n^1, x)) - u(r \pi(\mathcal{S}_n^2, x))| \leq \beta, \forall x \in \mathcal{S}_X, \forall r \in \mathcal{S}_R.$$

We also introduce an additional property of the investment algorithm which will be used instead of the notion of bounded loss for learning algorithms since, unlike loss functions, utility functions are typically neither bounded above nor below.

Definition 3 An investment algorithm π achieves a Δ -bounded utility range if for all sample sets \mathcal{S}_n^1 and \mathcal{S}_n^2 , we have that

$$|u(r \pi(\mathcal{S}_n^1, x)) - u(r' \pi(\mathcal{S}_n^2, x'))| \leq \Delta, \forall (r, x) \in \mathcal{S}_R \times \mathcal{S}_X, \forall (r', x') \in \mathcal{S}_R \times \mathcal{S}_X.$$

Theorem 8 (First adapted version of Theorem 11.1 in Mohri et al. (2012)) Given that an investment algorithm $\hat{q}(\cdot)$ is uniformly β -stable and achieves a Δ -bounded utility range, then one is guaranteed with a confidence of $1 - \delta$ that

$$E_F[u(R \cdot \pi(\mathcal{S}_n, X))] \geq E_{\hat{F}}[u(R \cdot \pi(\mathcal{S}_n, X))] - \beta - (2n\beta + \Delta) \sqrt{\frac{\log(1/\delta)}{2n}}.$$

The proof of this theorem follows exactly the same steps as the proof proposed in Mohri et al. (2012) except that the Δ -bound is used to bound the expression $|u(r \pi(\mathcal{S}_n^1, x)) - u(r' \pi(\mathcal{S}_n^2, x'))|$ instead of using upper and lower bounds for $u(\cdot)$ over its entire domain.

We now have in hand the necessary tools to obtain the result presented in Theorem 1 by considering the investment algorithm defined as $\pi(\mathcal{S}_n, x) := \hat{q}(\mathcal{S}_n)^T x$ where $\hat{q}(\mathcal{S}_n) := \arg \max_q E_{\hat{F}}[u(R \cdot q^T X)] + \lambda \|q\|_2^2$. In particular, we will be interested in identifying the β -stability and the Δ -bound for this estimator

Lemma 1 When Assumptions 1, 2 and 3 are satisfied, the investment algorithm $\pi(\cdot)$ is uniformly β -stable with $\beta = \frac{(\gamma \bar{r} \xi)^2}{2\lambda n}$.

Proof. We first establish that for any pair $(q_1, q_2) \in \mathbb{R}^p \times \mathbb{R}^p$, one has that

$$|u(r q_1^T x) - u(r q_2^T x)| \leq \gamma |r q_1^T x - r q_2^T x| \leq \gamma |r| \|x\|_2 \|q_1 - q_2\|_2 \leq \gamma \bar{r} \xi \|q_1 - q_2\|_2, \forall r \in \mathcal{S}_R, \forall x \in \mathcal{S}_X.$$

This follows naturally from Assumption 3 which states that $u(\cdot)$ is Lipschitz continuous, and Assumptions 1 and 2. Next, we can employ similar steps as in the proof of Proposition 11.1 from Mohri et al. (2012) to

establish that $\|\hat{q}(\mathcal{S}_n^1) - \hat{q}(\mathcal{S}_n^2)\|_2 \leq \gamma\bar{r}\xi/(\lambda n)$ which would complete our proof since for all $x \in \mathcal{S}_X$ and all $r \in \mathcal{S}_R$, we would have that

$$|u(r\hat{q}(\mathcal{S}_n^1)^T x) - u(r\hat{q}(\mathcal{S}_n^2)^T x)| \leq \gamma\bar{r}\xi\|\hat{q}(\mathcal{S}_n^1) - \hat{q}(\mathcal{S}_n^2)\|_2 \leq \gamma^2\bar{r}^2\xi^2/(\lambda n)$$

To summarize how $\|\hat{q}(\mathcal{S}_n^1) - \hat{q}(\mathcal{S}_n^2)\|_2 \leq \gamma\bar{r}\xi/(\lambda n)$ is obtained, we first observe that since $\hat{q}(\mathcal{S}_n^1)$ is the maximizer of the concave function $\widehat{EU}_{\mathcal{S}_n^1}(q) - \lambda\|q\|_2^2$ where $\widehat{EU}_{\mathcal{S}_n}(q) := (1/n)\sum_{i=1}^n u(r_i q^T x_i)$, there must exist a super-gradient equal to zero at $\hat{q}(\mathcal{S}_n^1)$. In particular, there must be a super-gradient $\nabla\widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^1))$ of $\widehat{EU}_{\mathcal{S}_n^1}(q)$ at $\hat{q}(\mathcal{S}_n^1)$ such that

$$\nabla\widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^1)) - 2\lambda\hat{q}(\mathcal{S}_n^1) = 0 \Rightarrow \nabla\widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^1)) = 2\lambda\hat{q}(\mathcal{S}_n^1).$$

Together with the concavity of $\widehat{EU}_{\mathcal{S}_n^1}(q)$, this implies that

$$\begin{aligned} \widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^2)) &\leq \widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^1)) + \nabla\widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^1))(\hat{q}(\mathcal{S}_n^2) - \hat{q}(\mathcal{S}_n^1)) \\ &= \widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^1)) + 2\lambda\hat{q}(\mathcal{S}_n^1)^T(\hat{q}(\mathcal{S}_n^2) - \hat{q}(\mathcal{S}_n^1)) \end{aligned}$$

and similarly that

$$\widehat{EU}_{\mathcal{S}_n^2}(\hat{q}(\mathcal{S}_n^1)) \leq \widehat{EU}_{\mathcal{S}_n^2}(\hat{q}(\mathcal{S}_n^2)) + 2\lambda\hat{q}(\mathcal{S}_n^2)^T(\hat{q}(\mathcal{S}_n^1) - \hat{q}(\mathcal{S}_n^2)).$$

Together, these two inequalities can be used to conclude that

$$\begin{aligned} 2\lambda\|\hat{q}(\mathcal{S}_n^2) - \hat{q}(\mathcal{S}_n^1)\|_2^2 &= 2\lambda\hat{q}(\mathcal{S}_n^2)^T(\hat{q}(\mathcal{S}_n^2) - \hat{q}(\mathcal{S}_n^1)) - 2\lambda\hat{q}(\mathcal{S}_n^1)^T(\hat{q}(\mathcal{S}_n^2) - \hat{q}(\mathcal{S}_n^1)) \\ &\leq \widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^1)) - \widehat{EU}_{\mathcal{S}_n^1}(\hat{q}(\mathcal{S}_n^2)) + \widehat{EU}_{\mathcal{S}_n^2}(\hat{q}(\mathcal{S}_n^2)) - \widehat{EU}_{\mathcal{S}_n^2}(\hat{q}(\mathcal{S}_n^1)) \\ &= (1/n)(u(r_m^1\hat{q}(\mathcal{S}_n^1)^T x_m^1) - u(r_m^1\hat{q}(\mathcal{S}_n^2)^T x_m^1) + u(r_m^2\hat{q}(\mathcal{S}_n^2)^T x_m^2) - u(r_m^2\hat{q}(\mathcal{S}_n^1)^T x_m^2)) \\ &\leq (2\gamma\bar{r}\xi/n)\|\hat{q}(\mathcal{S}_n^2) - \hat{q}(\mathcal{S}_n^1)\|_2, \end{aligned}$$

where the second equality comes from the definition of $\widehat{EU}_{\mathcal{S}_n}$ and the second inequality follows from Assumptions 1, 2, and 3. By dividing both sides of the inequality by $2\lambda\|\hat{q}(\mathcal{S}_n^2) - \hat{q}(\mathcal{S}_n^1)\|_2$ we get our stated property. \square

Lemma 2 *When Assumptions 1, 2 and 3 are satisfied, the Δ -bound on the utility range for $\pi(\cdot)$ is $\Delta := \frac{(\gamma+1)\xi^2\bar{r}^2}{2\lambda}$.*

Proof. This proof relies mostly on demonstrating that $\|\hat{q}(\mathcal{S}_n)\|_2 \leq \bar{r}\xi/(2\lambda)$ with probability one with respect to the randomness of \mathcal{S}_n . Indeed, when this is the case, then we have that

$$\begin{aligned} |u(r_1\hat{q}(\mathcal{S}_n^1)^T x_1) - u(r_2\hat{q}(\mathcal{S}_n^2)^T x_2)| &\leq u(\bar{r}^2\xi^2/(2\lambda)) - u(-\bar{r}^2\xi^2/(2\lambda)) \\ &\leq |u(\bar{r}^2\xi^2/(2\lambda)) - u(0)| + |u(0) - u(-\bar{r}^2\xi^2/(2\lambda))| \\ &\leq (1+\gamma)\bar{r}^2\xi^2/(2\lambda). \end{aligned}$$

In order to identify a bound on the norm of $\hat{q}(\mathcal{S}_n)$, we reformulate problem (2) as follows

$$\begin{aligned} \underset{s \in \mathbb{R}, v \in \mathbb{R}^p}{\text{maximize}} \quad & \frac{1}{n} \sum_{i=1}^n u(sR_i X_i^T v) - \lambda s^2 \\ \text{s. t.} \quad & s \geq 0, \|v\|_2 = 1, \end{aligned}$$

such that $\hat{q}(\mathcal{S}_n) = s^* \cdot v^*$ when (s^*, v^*) is the pair of optimal assignments for this optimization problem. It is therefore clear that $s^* = \|\hat{q}(\mathcal{S}_n)\|_2$ and our proof reduces to establishing an upper bound for s^* . By recognizing that $s^* = \arg \max_{s \geq 0} g(s) := \frac{1}{n} \sum_{i=1}^n u(sR_i X_i^T v^*) - \lambda s^2$ and that $g(s)$ is a concave function, then it is necessarily the case that if there exists a $\bar{s} \geq 0$ such that $g(\cdot)$ is non-increasing at \bar{s} then $s^* \leq \bar{s}$.

We can actually show that this is the case for $\bar{s} := \bar{r}\xi/(2\lambda)$ by upper bounding the impact of taking a step of $\delta > 0$:

$$\begin{aligned}
g(\bar{s} + \delta) - g(\bar{s}) &= \frac{1}{n} \sum_{i=1}^n (u((\bar{s} + \delta)R_i X_i^T v^*) - u(\bar{s}R_i X_i^T v^*)) - \lambda((\bar{s} + \delta)^2 - \bar{s}^2) \\
&\leq \frac{1}{n} \sum_{i=1}^n (u((\bar{s} + \delta)|R_i X_i^T v^*|) - u(\bar{s}|R_i X_i^T v^*|)) - \lambda((\bar{s} + \delta)^2 - \bar{s}^2) \\
&\leq \frac{1}{n} \sum_{i=1}^n \delta |R_i X_i^T v^*| - \lambda(2\bar{s}\delta + \delta^2) \\
&\leq \delta \bar{r}\xi - 2\lambda\bar{s}\delta - \delta^2 = -\delta^2 \leq 0,
\end{aligned}$$

where we first used the fact that $u(\cdot)$ is increasing, next that $u(y + \delta) \leq u(y) + \delta$ when $\delta \geq 0$ since it is a concave function with a subgradient of one at zero. Finally, we exploited Assumptions 1 and 2 and used the definition of \bar{s} . This completes our proof. \square

We have therefore established that under Assumptions 1, 2, and 3, we have that

$$E_F[u(R \cdot \pi(\mathcal{S}_n, X))] \geq E_{\hat{F}}[u(R \cdot \pi(\mathcal{S}_n, X))] - \Omega_1. \quad (9)$$

We can now conclude this section by demonstrating how Theorem 1 follows from this fact. In particular, by concavity of the utility function, we have that

$$u(\text{CE}(\hat{q}; F)) \leq u(\text{CE}(\hat{q}; \hat{F})) + (\text{CE}(\hat{q}; F) - \text{CE}(\hat{q}; \hat{F})) \nabla u(\text{CE}(\hat{q}; \hat{F})),$$

where $\nabla u(r)$ denotes any supergradient of $u(\cdot)$ at r . In particular, since $u(\cdot)$ is an increasing concave function, it follows that $\lim_{\epsilon \rightarrow 0^-} u'(\text{CE}(\hat{q}; \hat{F}) + \epsilon) \geq 0$ is one of the supergradient at $\text{CE}(\hat{q}; \hat{F})$. Combining this inequality with the inequality presented in Equation (9), we get

$$\begin{aligned}
u(\text{CE}(\hat{q}; \hat{F})) - \Omega_1 &= E_{\hat{F}}[u(R \cdot \pi(\mathcal{S}_n, X))] - \Omega_1 \leq E_F[u(R \cdot \pi(\mathcal{S}_n, X))] \\
&= u(\text{CE}(\hat{q}; F)) \leq u(\text{CE}(\hat{q}; \hat{F})) + (\text{CE}(\hat{q}; F) - \text{CE}(\hat{q}; \hat{F})) \nabla u(\text{CE}(\hat{q}; \hat{F}))
\end{aligned}$$

so that

$$\text{CE}(\hat{q}; F) \geq \text{CE}(\hat{q}; \hat{F}) - \Omega_1 / \nabla u(\text{CE}(\hat{q}; \hat{F}))$$

follows since it was assumed that $u(\cdot)$ is strictly increasing. This completes the proof of Theorem 1.

A.2 Proof of Theorem 2

We first show that there exist an optimal solution q^* for problem (1).

Lemma 3 *Given Assumptions 1, 2, and 5, we have that problem (1) is bounded.*

Proof. Similarly as was done in the proof of Lemma 2, we can reformulate problem (1) in terms of both an orientation vector and a scale decision variable. This gives us

$$\begin{aligned}
&\underset{s \in \mathbb{R}, v \in \mathbb{R}^p}{\text{maximize}} && E[u(sR X^T v)] \\
&\text{s. t.} && s \geq 0, \|v\|_2 = 1.
\end{aligned}$$

Note that the optimal value of the above problem is necessarily greater or equal to $u(0) = 0$. Moreover, in the case where it is exactly zero, then $s^* = 0$ is an optimal solution and we can conclude that there exists a bounded q^* . We therefore focus in what follows on the case where the optimal value is strictly positive or even unbounded.

Based on Assumption 5, since no feature induces an arbitrage opportunity, it follows that for any v of norm equal to one, either $P(RX^T v = 0) = 1$ or there exists a $\delta > 0$ and a $\varrho > 0$ such that $P\{RX^T v < -\delta\} = \varrho$.

In the former case, v cannot be an optimal assignment since $E[u(sRX^T v)] = u(0)$ for all $s \geq 0$ which we assumed was a sub-optimal objective value. Now, in the latter case, we let B be a discrete random variable with two states such that $P\{B = -\delta\} = 1 - P\{B = \bar{r}\xi\} = \varrho$. Since $|RX^T v| \leq \bar{r}\xi$, we have that $P\{B \geq r\} \geq P\{RX^T v \geq r\}$ for all $r \in \mathbb{R}$, i.e. that B stochastically dominates $RX^T v$, so that it must necessarily follow that $E[u(sB)] \geq E[u(sRX^T v)]$ for all $s \geq 0$. But, by the sublinearity assumption on u ,

$$\begin{aligned} \lim_{s \rightarrow \infty} E[u(sRX^T v)] &\leq \lim_{s \rightarrow \infty} E[u(sB)] = \lim_{s \rightarrow \infty} (\varrho u(-s\delta) + (1 - \varrho)u(s\bar{r}\xi)) \\ &\leq \lim_{s \rightarrow \infty} -\varrho s\delta + (1 - \varrho)o(s) = -\infty \end{aligned}$$

which shows that s^* , and therefore $\|q^*\|_2$, is bounded. \square

We next invoke a theorem from Sridharan et al. (2009) which will be of use.

Theorem 9 (See Theorem 1 in Sridharan et al. (2009)) *Let \mathcal{W} be a closed convex subset of a Banach space with norm $\|\cdot\|$ and dual norm $\|\cdot\|_*$ and consider $f(w, x, r) := \ell(\langle w, \Phi(x, r) \rangle; x, r) + \lambda\|w\|^2$, where $\ell : \mathbb{R} \times \mathbb{R}^{p+1} \rightarrow \mathbb{R}$ is L -Lipschitz and convex in its first argument and where $\Phi : \mathbb{R}^p \times \mathbb{R} \rightarrow \mathcal{W}$ is bounded such that $P(\|\Phi(X, R)\| \leq B) = 1$. Then, for any $\delta > 0$, with probability at least $1 - \delta$ over a sample of size n , we have that :*

$$F(\hat{w}) - F(w^*) \leq \frac{4L^2 B^2 (32 + \log(1/\delta))}{\lambda n},$$

where $F(w) := E[f(w, X, R)]$, $\hat{F}(w) := (1/n) \sum_{i=1}^n f(w, x_i, r_i)$, $w^* := \arg \min_{w \in \mathcal{W}} F(w)$, and $\hat{w} := \arg \min_{w \in \mathcal{W}} \hat{F}(w)$.

By considering that $\Phi(x, r) := x$, which is bounded by ξ when using the 2-norm, and that $\ell(z; x, r) := -u(zr)$ which is convex and $\gamma\bar{r}$ -Lipschitz for all $x \in \mathcal{S}_X$ and $r \in \mathcal{S}_R$, we immediately get the following corollary. Note that to get this result we also exploit the fact that $f(q, x, r)$ is 2λ -strongly convex with respect to the 2-norm which implies, based on for example Lemma 13 in Shalev-Shwartz (2007) that : $F(\hat{w}) \geq F(w^*) + \lambda\|\hat{w} - w^*\|_2^2$.

Corollary 2 *Given that Assumptions 1, 2, and 3 are satisfied, then one has with confidence of $1 - \delta$ that*

$$-\lambda\|\hat{q} - q_\lambda^*\|_2^2 \geq EU_\lambda(\hat{q}) - EU_\lambda(q_\lambda^*) \geq -\omega,$$

where $\omega := 4\gamma^2\bar{r}^2\xi^2(32 + \log(1/\delta))/(\lambda n)$ and $EU_\lambda(q) := E_F(u(R \cdot q^T X)) - \lambda\|q\|_2^2$, with $q^* := \arg \min_q EU(q)$ and $q_\lambda^* := \arg \min_q EU_\lambda(q)$.

Notice that Corollary 2 implies with confidence $1 - \delta$ that

$$EU(\hat{q}) - EU(q_\lambda^*) \geq \lambda(\|\hat{q}\|_2^2 - \|q_\lambda^*\|_2^2) - \omega \geq -\lambda(\|\hat{q} - q_\lambda^*\|_2^2 + 2\|\hat{q}\|_2\|\hat{q} - q_\lambda^*\|_2) - \omega.$$

where $EU(\hat{q}) := E_F(u(R \cdot \hat{q}^T X))$. As shown in Lemma 2, $\|\hat{q}\|_2 \leq \bar{r}\xi/(2\lambda)$. Hence, Theorem 2 further implies concerning the same $1 - \delta$ probability outcomes that $\|\hat{q} - q_\lambda^*\|_2^2 \leq \omega/\lambda$, and therefore $\|\hat{q} - q_\lambda^*\|_2 \leq \sqrt{\omega/\lambda}$, so that we end up with

$$EU(\hat{q}) - EU(q_\lambda^*) \geq -2\omega - \bar{r}\xi\sqrt{\frac{\omega}{\lambda}}.$$

with probability $1 - \delta$. Finally, note that since by the definition of q_λ^* , we have that

$$EU(q^*) - \lambda\|q^*\|_2^2 \leq EU(q_\lambda^*) - \lambda\|q_\lambda^*\|_2^2,$$

it follows that

$$EU(q^*) - EU(q_\lambda^*) \leq \lambda(\|q^*\|_2^2 - \|q_\lambda^*\|_2^2) \leq \lambda\|q^*\|_2^2,$$

so that we can bound the suboptimality of the policy \hat{q} with probability $1 - \delta$ in the following fashion:

$$\begin{aligned} EU(\hat{q}) &= EU(q^*) + EU(\hat{q}) - EU(q_\lambda^*) + EU(q_\lambda^*) - EU(q^*) \\ &\geq EU(q^*) - 2\omega - \bar{r}\xi\sqrt{\frac{\omega}{\lambda}} - \lambda\|q^*\|_2^2 = EU(q^*) - \Omega_2. \end{aligned}$$

This relation can be exploited in a similar way as in the proof of Theorem 1 (see Section A.1) to derive the relation between certainty equivalents that is presented in our theorem.

A.3 Proof of Theorem 3

This proof follows exactly the same steps as the proof of Theorem 1. Namely, we can exploit another adapted version of Theorem 11.1 in Mohri et al. (2012) for a multi-asset investment algorithm $\pi : \mathbb{R}^{p+1 \times m \times n} \times \mathbb{R}^p \rightarrow \mathbb{R}^m$ which constructs an investment portfolio based on a training set $\mathcal{S}_n := \{(\mathbf{x}_1, \mathbf{r}_1), \dots, (\mathbf{x}_m, \mathbf{r}_m)\}$, where each $\mathbf{x}_i \in \mathbb{R}^{m \times p}$ and each $\mathbf{r}_i \in \mathbb{R}^m$, and the current market side-information matrix \mathbf{x} . The natural extension of β -stability and Δ -bound to this context is described in what follows.

Definition 4 A multi-asset investment algorithm $\pi(\cdot)$ is uniformly β -stable if for any two sample sets $\mathcal{S}_n^1 := \{(\mathbf{x}_i^1, \mathbf{r}_i^1)\}_{i=1}^n$ and $\mathcal{S}_n^2 := \{(\mathbf{x}_i^2, \mathbf{r}_i^2)\}_{i=1}^n$ that are exactly identical except for the j -th sample, i.e., $(\mathbf{x}_i^1, \mathbf{r}_i^1) = (\mathbf{x}_i^2, \mathbf{r}_i^2)$ for all $i \neq j$, the following holds:

$$|u(\mathbf{r}^T \pi(\mathcal{S}_n^1, \mathbf{x})) - u(\mathbf{r}^T \pi(\mathcal{S}_n^2, \mathbf{x}))| \leq \beta, \forall \mathbf{x} \in \mathcal{S}_x, \forall \mathbf{r} \in \mathcal{S}_r.$$

Definition 5 A multi-asset investment algorithm $\pi(\cdot)$ achieves a Δ -bounded utility range if for all sample sets \mathcal{S}_n^1 and \mathcal{S}_n^2 :

$$|u(\mathbf{r} \pi(\mathcal{S}_n^1, \mathbf{x})) - u(\mathbf{r}' \pi(\mathcal{S}_n^2, \mathbf{x}'))| \leq \Delta, \forall (\mathbf{r}, \mathbf{x}) \in \mathcal{S}_r \times \mathcal{S}_x, \forall (\mathbf{r}', \mathbf{x}') \in \mathcal{S}_r \times \mathcal{S}_x.$$

This leads us to adapting Theorem 8 to the context of multi-asset investment algorithms.

Theorem 10 (Second adapted version of Theorem 11.1 in Mohri et al. (2012)) Given that a multi-asset investment algorithm $\hat{\mathbf{q}}(\cdot)$ is uniformly β -stable and achieves a Δ -bounded utility range, then one is guaranteed with a confidence of $1 - \delta$ that

$$E_F[u(\mathbf{R}^T \pi(\mathcal{S}_n, \mathbf{X}))] \geq E_{\hat{F}}[u(\mathbf{R}^T \pi(\mathcal{S}_n, \mathbf{X}))] - \beta - (2n\beta + \Delta) \sqrt{\frac{\log(1/\delta)}{2n}}.$$

This time, we consider a multi-asset investment algorithm to be defined as $\pi(\mathcal{S}_n, \mathbf{x}) := \mathbf{x} \hat{\mathbf{q}}(\mathcal{S}_n)$ where $\hat{\mathbf{q}}(\mathcal{S}_n) := \arg \max_{\mathbf{q}} E_{\hat{F}}[u(\mathbf{R}^T \mathbf{X} \mathbf{q})] + \lambda \|\mathbf{q}\|_2^2$. In particular, the same steps can be used to verify that this algorithm is $\frac{(\gamma \bar{r} \xi)^2}{2\lambda n}$ -stable and that $\Delta := \frac{(\gamma+1)\xi^2 \bar{r}^2}{2\lambda}$ is a valid Δ -bound for the utility range achieved using this estimator. For completeness, we repeat the argument that is used to establish the bound on $\hat{\mathbf{q}}(\mathcal{S}_n)$.

Lemma 4 Given Assumptions 6 and 7, we have that $\|\hat{\mathbf{q}}(\mathcal{S}_n)\|_2 \leq \bar{r}\xi/(2\lambda)$ with probability one.

Proof. We first reformulate problem (2) as follows

$$\begin{aligned} & \underset{s \in \mathbb{R}, \mathbf{v} \in \mathbb{R}^p}{\text{maximize}} && \frac{1}{n} \sum_{i=1}^n u(s \mathbf{R}_i^T \mathbf{X}_i \mathbf{v}) - \lambda s^2 \\ & \text{s. t.} && s \geq 0, \|\mathbf{v}\|_2 = 1, \end{aligned}$$

such that $\hat{\mathbf{q}}(\mathcal{S}_n) = s^* \cdot \mathbf{v}^*$ when (s^*, \mathbf{v}^*) is the pair of optimal assignments for this optimization problem. It is therefore clear that $s^* = \|\hat{\mathbf{q}}(\mathcal{S}_n)\|_2$ and our proof reduces to establishing an upper bound for s^* . By recognizing that $s^* = \arg \max_{s \geq 0} g(s) := \frac{1}{n} \sum_{i=1}^n u(s \mathbf{R}_i^T \mathbf{X}_i \mathbf{v}^*) - \lambda s^2$ and that $g(s)$ is a concave function, then it is necessarily the case that if there exists a $\bar{s} \geq 0$ such that $g(\cdot)$ is non-increasing at \bar{s} then $s^* \leq \bar{s}$. We can actually show that this is the case for $\bar{s} := \bar{r}\xi/(2\lambda)$ by upper bounding the impact of taking a step of $\delta > 0$:

$$\begin{aligned} g(\bar{s} + \delta) - g(\bar{s}) &= \frac{1}{n} \sum_{i=1}^n (u((\bar{s} + \delta) \mathbf{R}_i^T \mathbf{X}_i \mathbf{v}^*) - u(\bar{s} \mathbf{R}_i^T \mathbf{X}_i \mathbf{v}^*)) - \lambda((\bar{s} + \delta)^2 - \bar{s}^2) \\ &\leq \frac{1}{n} \sum_{i=1}^n (u((\bar{s} + \delta) |\mathbf{R}_i^T \mathbf{X}_i \mathbf{v}^*|) - u(\bar{s} |\mathbf{R}_i^T \mathbf{X}_i \mathbf{v}^*|)) - \lambda((\bar{s} + \delta)^2 - \bar{s}^2) \\ &\leq \frac{1}{n} \sum_{i=1}^n \delta |\mathbf{R}_i^T \mathbf{X}_i \mathbf{v}^*| - \lambda(2\bar{s}\delta + \delta^2) \\ &\leq \delta \bar{r}\xi - 2\lambda \bar{s}\delta - \lambda\delta^2 = -\lambda\delta^2 \leq 0, \end{aligned}$$

where we first used the fact that $u(\cdot)$ is increasing, next that $u(y + \delta) \leq u(y) + \delta$ when $\delta \geq 0$ since it is a concave function with a subgradient of one at zero. Finally, we exploited Assumptions 6 and 7. \square

A.4 Proof of Theorem 4

The steps of this proof follows almost exactly as the steps presented for the proof of Theorem 2 in Appendix A.2. We can first demonstrate that problem (1) remains bounded in the multi-asset setting using the same steps as in the proof of Lemma 3. We then apply Theorem 9 using this time $\phi(\mathbf{x}, \mathbf{r}) := \mathbf{x}^T \mathbf{r}$ (instead of $\phi(x, r) := x$), which is bounded by $\bar{r}\xi$ when using the 2-norm, and using $\ell(z; \mathbf{x}, \mathbf{r}) := -u(z)$ (instead of $\ell(z; x, r) := -u(zr)$) which is convex and γ -Lipschitz. Together with the fact that $f(q, \mathbf{x}, \mathbf{r})$ is 2λ -strongly convex with respect to the 2-norm, we immediately get the following corollary.

Corollary 3 *Given that Assumptions 3, 6, and 7 are satisfied, then one has with confidence of $1 - \delta$ that*

$$-\lambda \|\hat{\mathbf{q}} - \mathbf{q}_\lambda^*\|_2^2 \geq EU_\lambda(\hat{\mathbf{q}}) - EU_\lambda(\mathbf{q}_\lambda^*) \geq -\omega,$$

where $\omega := 4\gamma^2 \bar{r}^2 \xi^2 (32 + \log(1/\delta)) / (\lambda n)$ and where $EU(\mathbf{q}) := E_F(u(\mathbf{R}^T \mathbf{X} \mathbf{q}))$ and $EU_\lambda(\mathbf{q}) := E_F(u(\mathbf{R}^T \mathbf{X} \mathbf{q})) - \lambda \|\mathbf{q}\|_2^2$, with $\mathbf{q}^* := \arg \min_{\mathbf{q}} EU(\mathbf{q})$ and $\mathbf{q}_\lambda^* := \arg \min_{\mathbf{q}} EU_\lambda(\mathbf{q})$.

The rest of the proof is straightforward and exploits the bound on $\hat{\mathbf{q}}$ established in Lemma 4.

A.5 Proof of Theorem 6

This proof follows exactly the same steps as the proof of Theorem 1 and 3. Namely, we exploit exactly the version of Theorem 11.1 in Mohri et al. (2012) presented in Theorem 8 with the investment algorithm $\pi : \mathbb{R}^{p+1 \times m \times n} \times \mathbb{R}^p \rightarrow \mathbb{R}^m$ defined as $\pi(\mathcal{S}_n, x) := \langle \hat{w}(\mathcal{S}_n), \Phi(x) \rangle$. Once again, the same steps can be used, in the reproducing Hilbert kernel space \mathcal{W} replacing q with w and x with $\Phi(x)$ in the analysis, to verify that this algorithm is $\frac{(\gamma \bar{r} \xi)^2}{2\lambda n}$ -stable and that $\Delta := \frac{(\gamma+1)\xi^2 \bar{r}^2}{2\lambda}$ is a valid Δ -bound for the utility range achieved using this estimator. For completeness, we repeat the argument that is used to establish the bound on $\hat{w}(\mathcal{S}_n)$.

Lemma 5 *Given Assumptions 1 and 9, we have that $\|\hat{w}(\mathcal{S}_n)\|_2 \leq \bar{r}\xi / (2\lambda)$ with probability one.*

Proof. We first reformulate problem (7) as follows

$$\begin{aligned} & \underset{s \in \mathbb{R}, w \in \mathcal{W}}{\text{maximize}} && \frac{1}{n} \sum_{i=1}^n u(r_i \cdot \langle sw, \Phi(x_i) \rangle) - \lambda \|sw\|_2^2 \\ & \text{s. t.} && s \geq 0, \|w\|_2 = 1, \end{aligned}$$

such that $\hat{w} = s^* \cdot w^*$ when (s^*, w^*) is the pair of optimal assignments for this optimization problem. It is therefore clear that $s^* = \|\hat{w}\|_2$ and our proof reduces to establishing an upper bound for s^* . By recognizing that $s^* = \arg \max_{s \geq 0} g(s) := \frac{1}{n} \sum_{i=1}^n u(r_i \cdot \langle sw, \Phi(x_i) \rangle) - \lambda \|sw\|_2^2$ and that $g(s)$ is a concave function, then it is necessarily the case that if there exists a $\bar{s} \geq 0$ such that $g(\cdot)$ is non-increasing at \bar{s} then $s^* \leq \bar{s}$. We can actually show that this is the case for $\bar{s} := \bar{r}\xi / (2\lambda)$ by upper bounding the impact of taking a step of $\delta > 0$:

$$\begin{aligned} g(\bar{s} + \delta) - g(\bar{s}) &= \frac{1}{n} \sum_{i=1}^n (u(r_i \cdot \langle (\bar{s} + \delta)w, \Phi(x_i) \rangle) - u(r_i \cdot \langle \bar{s}w, \Phi(x_i) \rangle)) - \lambda((\bar{s} + \delta)^2 - \bar{s}^2) \\ &\leq \frac{1}{n} \sum_{i=1}^n (u((\bar{s} + \delta)|r_i \cdot \langle w, \Phi(x_i) \rangle) - u(\bar{s}|r_i \cdot \langle w, \Phi(x_i) \rangle)) - \lambda((\bar{s} + \delta)^2 - \bar{s}^2) \\ &\leq \frac{1}{n} \sum_{i=1}^n \delta |r_i \langle w, \Phi(x_i) \rangle| - \lambda(2\bar{s}\delta + \delta^2) \\ &\leq \delta \bar{r}\xi - 2\lambda \bar{s}\delta - \lambda\delta^2 = -\lambda\delta^2 \leq 0, \end{aligned}$$

following exactly the same arguments as in the proof of Theorem 2. □

A.6 Proof of Theorem 7

The steps of this proof follows exactly as the steps presented for the proof of Theorem 2 but in the reproducing Hilbert kernel space \mathcal{W} . We can first demonstrate that problem (6) remains bounded yet this time when Assumption 10 is satisfied.

Lemma 6 *Given Assumptions 1, 2, and 10, we have that problem (6) is bounded.*

Proof. Similarly as was done in the proof of Lemma 3, we can reformulate problem (6) in terms of both an orientation vector and a scale decision variable. This gives us

$$\underset{s \in \mathbb{R}, w \in \mathcal{W}}{\text{maximize}} \quad E[u(R \cdot \langle sw, \Phi(X) \rangle)] \quad (10a)$$

$$\text{s. t.} \quad s \geq 0, \|w\|_2 = 1. \quad (10b)$$

We can once again focus on the case where the optimal value is strictly positive or even infinite. Let us consider any fixed w of norm equal to one and verify that we can once again identify some $\delta > 0$ and $\varrho > 0$ such that $P(R \cdot \langle w, \Phi(X) \rangle < -\delta) = \varrho$ in order for the rest of the proof to follow as before. In particular, we can first make the case that $P(|R \cdot \langle w, \Phi(X) \rangle| > 0) = 0$ necessarily leads to a sub-optimal assignment for w . Indeed, for such assignments $R \cdot \langle w, \Phi(X) \rangle = 0$ with probability one hence the objective value of problem (10) is zero which we considered sub-optimal.

Next, since we have that $P(|R \cdot \langle w, \Phi(X) \rangle| > 0) > 0$, it must be that at least one of the following statements is true:

$$\begin{aligned} P(R > 0 | X \in \mathcal{S}_X^+) P(X \in \mathcal{S}_X^+) &> 0 \\ P(R < 0 | X \in \mathcal{S}_X^+) P(X \in \mathcal{S}_X^+) &> 0 \\ P(R > 0 | X \in \mathcal{S}_X^-) P(X \in \mathcal{S}_X^-) &> 0 \\ P(R > 0 | X \in \mathcal{S}_X^-) P(X \in \mathcal{S}_X^-) &> 0, \end{aligned}$$

where $\mathcal{S}_X^+ := \{x \in \mathbb{R}^p \mid \langle w, \Phi(X) \rangle > 0\}$ while $\mathcal{S}_X^- := \{x \in \mathbb{R}^p \mid \langle w, \Phi(X) \rangle < 0\}$. For each of these cases, we can confirm that this necessarily implies that $P(R \cdot \langle w, \Phi(X) \rangle < 0) > 0$. In the first case, one can argue that

$$\begin{aligned} P(R > 0 | X \in \mathcal{S}_X^+) P(X \in \mathcal{S}_X^+) > 0 &\Rightarrow P(R > 0 | X \in \mathcal{S}_X^+) > 0 \ \& \ P(X \in \mathcal{S}_X^+) > 0 \\ &\Rightarrow P(R < 0 | X \in \mathcal{S}_X^+) > 0 \ \& \ P(X \in \mathcal{S}_X^+) > 0 \\ &\Rightarrow P(R \cdot \langle w, \Phi(X) \rangle < 0) > 0, \end{aligned}$$

where we employed Assumption 10 to get the second implication. On the other hand, the second case leads to:

$$\begin{aligned} P(R < 0 | X \in \mathcal{S}_X^+) P(X \in \mathcal{S}_X^+) > 0 &\Rightarrow P(R < 0 | X \in \mathcal{S}_X^+) > 0 \ \& \ P(X \in \mathcal{S}_X^+) > 0 \\ &\Rightarrow P(R \cdot \langle w, \Phi(X) \rangle < 0) > 0. \end{aligned}$$

The next two cases are similar. We can therefore conclude that there must exist some $\delta > 0$ and $\varrho > 0$ such that $P(R \cdot \langle w, \Phi(X) \rangle < -\delta) = \varrho$. \square

We then apply Theorem 9 using this time $\phi(x, r) := \Phi(x)$ (instead of $\phi(x, r) := x$), which is again bounded by ξ when using the 2-norm, and using again $\ell(z; x, r) := -u(z)$ which is convex and $\gamma\bar{r}$ -Lipschitz. Together with the fact that $f(w, x, r)$ is 2λ -strongly convex with respect to the 2-norm, we immediately get the following corollary.

Corollary 4 *Given that Assumptions 3, 1, and 9 are satisfied, then one has with confidence of $1 - \delta$ that*

$$-\lambda \|\hat{w} - w_\lambda^*\|_2^2 \geq EU_\lambda(\hat{w}) - EU_\lambda(w_\lambda^*) \geq -\omega,$$

where $\omega := 4\gamma^2\bar{r}^2\xi^2(32 + \log(1/\delta))/(\lambda n)$ and where $EU(w) := E_F(u(R \cdot \langle w, \Phi(X) \rangle))$ and $EU_\lambda(w) := E_F(u(R \cdot \langle w, \Phi(X) \rangle)) - \lambda \|w\|_2^2$, with $w^* := \arg \min_w EU(w)$ and $w_\lambda^* := \arg \min_w EU_\lambda(w)$.

The rest of the proof is straightforward and exploits the bound on \hat{w} established in Lemma 5.

References

- Bertsimas D, Van Parys B. 2017. Bootstrap robust prescriptive analytics. Working draft.
- Bousquet O, Elisseeff A. 2002. Stability and generalization. *The Journal of Machine Learning Research*. 2:499–526.
- Brandt MW, Santa-Clara P, Valkanov R. 2009. Parametric portfolio policies: Exploiting characteristics in the cross-section of equity returns. *Review of Financial Studies*. 22(9):3411–3447.
- Caramanis C, Mannor S, Xu H. 2012. Robust optimization in machine learning. In: Sra S, Nowozin S, Wright SJ, editors. *Optimization for machine learning*. Cambridge: MIT Press; chap. 14; p. 369–402.
- Delage E, Ye Y. 2010. Distributionally robust optimization under moment uncertainty with application to data-driven problems. *Operations Research*. 58(3):595–612.
- Esfahani P, Kuhn D. 2017. Data-driven distributionally robust optimization using the Wasserstein metric: Performance guarantees and tractable reformulations. *Mathematical Programming*.
- Fama EF. 1991. Efficient capital markets: II. *The Journal of Finance*. 46(5):1575–1617.
- Fama EF, French KR. 1993. Common risk factors in the returns on stocks and bonds. *Journal of Financial Economics*. 33(1):3–56.
- Goldfarb D, Iyengar G. 2003. Robust portfolio selection problems. *Mathematics of Operations Research*. 28(1):1–38.
- Györfi L, Lugosi G, Udina F. 2006. Nonparametric kernel-based sequential investment strategies. *Mathematical Finance*. 16(2):337.
- Hofmann T, Schölkopf B, Smola A. 2008. Kernel methods in machine learning. *Annals of Statistics*. 36(3):1171–1220.
- Huang D, Zhu S, Fabozzi F, Fukushima M. 2010. Portfolio selection under distributional uncertainty: A relative robust CVaR approach. *European Journal of Operations Research*. 203(1):185–194.
- Ingersoll J. 1987. *Theory of financial decision making*. Rowman & Littlefield Publishers.
- Madan DB, Carr PP, Chang EC. 1998. The variance gamma process and option pricing. *European Finance Review*. 2(1):79–105.
- Malkiel BG, Fama EF. 1970. Efficient capital markets: A review of theory and empirical work. *The Journal of Finance*. 25(2):383–417.
- Markowitz H. 1952. Portfolio selection. *The Journal of Finance*. 7(1):77–91.
- Mohri M, Rostamizadeh A, Talwalkar A. 2012. *Foundations of machine learning*. MIT press.
- Oliveras-Nadal AV, DeMiguel V. 2018. Technical note—a robust perspective on transaction costs in portfolio optimization. Working draft.
- Rudin C, Vahn GY. 2015. The big data newsvendor: Practical insights from machine learning. Working draft.
- Salmon F. 2009. Recipe for disaster: the formula that killed Wall Street. *WIRED Magazine*.
- Shafieezadeh-Abadeh S, Kuhn D, Esfahani P. 2017. Regularization via mass transportation.
- Shalev-Shwartz S. 2007. *Online learning: Theory, algorithms, and applications [dissertation]*. The Hebrew University of Jerusalem.
- Sridharan K, Shalev-Shwartz S, Srebro N. 2009. Fast rates for regularized objectives. In: *Advances in Neural Information Processing Systems*. p. 1545–1552.
- Takano Y, Gotoh J. 2014. Multi-period portfolio selection using kernel-based control policy with dimensionality reduction. *Expert Systems with Applications*. 41(8):3901–3914.
- von Neumann J, Morgenstern O. 1944. *Theory of games and economic behavior*. Princeton University Press.