

2022

**Centre intégré de
santé et de services
sociaux de Laval**

Preparation guide for a surgery

Ureteroscopy + Cystolitholopaxy



This guide will help you understand and get ready for your surgery.

Read it over with your family and bring this guide with you the day of your surgery.

A publication of

Direction des services professionnels
Centre intégré de santé et de services sociaux (CISSS) de Laval
Preadmission Clinic
1755, boulevard René-Laënnec
Laval (Québec) H7M 3L9
Téléphone: 450 975-5566
Website: www.lavalensante.com

Researcher and writer

Judith Dubois, nurse clinician, AIC
Preadmission Clinic

In collaboration with

Clinical nurses from the Preadmission Clinic and urologists from Hôpital Cité-de-la-Santé

Linguistic revision

Communications Department

Diffusion

Any reproduction of excerpts from this guide by any method whatsoever is prohibited without written authorization from the publisher.

Traduction and adaptation of

Guide de préparation à une chirurgie : urétéroscopie + cystolitholapaxie, 2020

© Centre intégré de santé et de services sociaux (CISSS) de Laval, 2022

Legal deposit - Bibliothèque et Archives nationales du Québec, 2022

ISBN 978-2-550-91259-0 (print)

ISBN 978-2-550-91261-3 (PDF)


TABLE OF CONTENTS

ANATOMY	1
YOUR PROCEDURE	2
BEFORE YOUR SURGERY	3
HOW TO GET READY FOR YOUR SURGERY	4
BEFORE YOUR VISITE TO PREADMISSION CLINIC	5
CONSENT TO SURGERY AND ANESTHESIA	5
TOBACCO	6
ALCOHOL	6
DISCHARGE PLANNING	6
PREOPERATIVE DIET	7
THE NIGHT BEFORE THE SURGERY	8
THE DAY OF YOUR SURGERY	9
HYGIENE BEFORE YOUR SURGERY	10
WHAT TO BRING TO THE HOSPITAL	11
WHEN YOU ARRIVE AT THE SURGERY UNIT	12
THE DAY SURGERY UNIT	12
OPERATING ROOM	13
IN THE RECOVERY ROOM	14
RETURN TO THE DAY SURGERY UNIT OR HOSPITAL UNIT	14
CONTROLLING YOUR PAIN	15
BREATHING EXERCISES	16
CIRCULATION EXERCISES	18
GETTING UP	19
YOUR DISCHARGE FROM THE UNIT	20
ONCE YOU GET BACK HOME – INSTRUCTIONS	21
NUTRITION AND HYDRATION	22
ACTIVITIES	23
COMPLICATIONS	24
RESOURCES	25
BIBLIOGRAPHY	26

ANATOMY

The urinary tract consists of the kidneys, two ureters, the bladder, the urethra and the urinary meatus.

Located under the ribcage on either side of the spine, the kidneys eliminate waste from the body. This waste is dissolved in the urine generated by the kidneys and flows through the urinary tract (renal calyces, renal pelvis, and ureters) to the bladder.



Coup d'œil sur le monde – Emorsite
<https://goo.gl/images/pfyhjw>

Bladder, ureteral and kidney stones

Stones are small, crystalline substances that can become lodged in different areas of the urinary tract. These stones form for various reasons and can get stuck in the urinary system.

YOUR PROCEDURE

URETEROSCOPY

Ureteroscopy is an endoscopic procedure (i.e. done through the body's natural openings instead of through a surgical incision) that allows your doctor to explore the entire urinary tract. This procedure is mainly performed to remove urinary (kidney) stones that get stuck in the kidney or a ureter. This procedure can also be used to diagnose and treat any unexplained bleeding or blockage of a ureter or to perform a biopsy.

Ureteroscopy involves inserting a ureteroscope (camera) through the urethra into the bladder, and then up through the ureter to the kidney. The doctor visually controls the camera from a miniature screen. X-rays are usually taken during the procedure. If your doctor sees a stone, he or she will gently remove it with an instrument that looks like tweezers or with a "snare" made of thread. The doctor may use a laser to cut up larger stones so that they can be extracted.

* Note that a "double-J" stent may be placed some time before your procedure and after if your ureter is blocked. This will protect your kidney by allowing urine to empty normally into the bladder. The stent may stay in place for a few weeks or a few months. It is called a "double-J" stent because it is curved at both ends.



http://www.uropage.com/ART_calc5.php

CYSTOLITHOLAPAXY

This procedure consists of inserting a device through the body's natural openings to dissolve a stone or stones in the bladder. After they are ground up, the stones can come out naturally.

* Note that this procedure does not require the placement of a double-J stent.

For **kidney stone surgery**, you will need to have an X-ray taken of your abdomen some time after you arrive at the hospital and before going to the operating room. This helps your doctor verify the exact location of the stone before performing the procedure.

BEFORE YOUR SURGERY

Admission date and time

- You will receive a call from the hospital's pre-admission department.
- The secretary will inform you of the date of your surgery.
- You will be informed of the time of your arrival at the hospital by phone 24 to 48 hours before the surgery.



SYMPTOMS TO MONITOR

If you have any of these symptoms or conditions one week before the date of your surgery:

- ☐ You have a sore throat, a cold or the flu.
- ☐ You have a fever.
- ☐ You are taking antibiotics.
- ☐ You have a contagious disease (e.g., chicken pox), or you have recently been exposed to someone with a contagious disease.
- ☐ You have a possible or unconfirmed pregnancy.
- ☐ Redness, inflammation, discharge, wound or any other problem at the operating site

Call immediately to inform the administrative officer:

 Urology: 450 975-5394



HOW TO GET READY FOR YOUR SURGERY

When to stop or continue your medication



At your appointment with your surgeon or preadmission nurse, you will be informed whether you need to stop or continue your medication before your surgery.

- ☐ **Aspirin®**, ☐ **Asaphen®**, ☐ **Rivasa®**, ☐ **Entrophen®**, ☐ **Novasen®**, ☐ **Persantine®**, ☐ **MSD AAS**, ☐ **Aggrenox®** (dipyridamole/ASA), etc.
☐ Stop ____ days before your surgery.
☐ Do not stop this medication.
- ☐ **Plavix®** (clopidogrel)
☐ Stop ____ days before your surgery.
☐ Do not stop this medication.
- ☐ **Effient®** (prasugrel), ☐ **Ticlid®** (ticlopidine), ☐ **Brilinta®** (ticagrelor)
☐ Stop ____ days before your surgery.
☐ Do not stop.
- **Anti-inflammatory drugs** (e.g., ibuprofen such as Advil®, Motrin® (including for children), Celebrex®, Maxidol®, Aleve®, Naprosyn®, etc.)
 Stop 2 days before your surgery.
- **All natural products** (glucosamine, omega 3, vitamin E, etc.).
 Stop 7 days before your surgery.

You can keep taking drugs such as Tylenol®, Tylenol® Extra-Strength, acetaminophen and Tempra® until midnight the night before your surgery.

If you are taking Coumadin®, Sintrom®, Pradaxa®, Xarelto®, Eliquis®, Lixiana® :

A hospital pharmacist will call you approximately 1 to 3 weeks before your surgery and may ask you to have a blood sample taken.

When the pharmacy department has received your results, you will be called again about when to stop taking this medication.



You must follow this instruction.

BEFORE YOUR VISITE TO PREADMISSION CLINIC

Your record will be transferred to the hospital's Preadmission Clinic. Someone will call you with the date and time of your Preadmission Clinic appointment.

Date and time of your appointment : _____

During your Preadmission Clinic visit

You will:

Meet with a nurse, who will explain how to prepare for surgery and what to expect during your hospital stay.

Have an ECG (electrocardiogram) if the nurse determines that you need one.

Have blood taken, if required. You will be sent to the hospital's test centre.

The nurse will tell you if you need more tests or have to meet with other doctors or professionals.



CONSENT TO SURGERY AND ANESTHESIA



At your preadmission meeting, the nurse will ask you to sign the consent to surgery and anesthesia.

This consent means that the surgeon clearly explained why you need this operation, what the procedure entails, the potential risks, and the desired results of the operation.

If you did not get the proper information, you must contact your surgeon. The preadmission clinic nurse can help you. You will need to sign the consent form the morning of your surgery.

For further information about anesthesia, please read the guide on anesthesia and pain relief Information guide which the nurse will provide when attending your preadmission meeting.

TOBACCO

Quitting smoking or reducing the amount you smoke will decrease your risk of respiratory problems after your surgery, aid in the healing of your surgical wound, and help you better manage pain.



If you need help to quit smoking, don't hesitate to contact:

- Your CLSC at **450 978-8300, extension 3169** (for Laval residents).
- Your pharmacist or family doctor.
- The Quit Smoking Centre nearest you at **1-866-JARRETE (527-7383)**.

Website: tobaccofreequebec.ca/iquitnow/.

ALCOHOL

Avoid drinking alcohol **7 days before your surgery**. Alcohol can interact with some medications and increase the risk of bleeding and complications.

To get help to stop right now, contact the regional hotline (for Laval residents):

Alcochoix+ Laval at 450 622-5110, ext. 64005.

Website: [Alcochoix+ | Gouvernement du Québec \(quebec.ca\)](http://Alcochoix+ | Gouvernement du Québec (quebec.ca))



DISCHARGE PLANNING

Before your operation, it is important that you prepare in advance for your return home.





- Ask another adult to come pick you up at the hospital. You must organize a ride home in advance. This person must be available to pick you up once your discharge is signed.
- Prepare meals in advance for the days after your operation.
- Get help for errands, housework and appointments.
- If you live by yourself and your operation reduces your mobility, you need to think about having another adult stay with you during your recovery.

PREOPERATIVE DIET

The goal of this diet is to ensure that you have the strength and nutrients you need to recover quickly.

Suggestions to boost your protein intake.



You can also take a supplement such as Ensure or Boost.

Add this	To this
Skim milk powder or protein powder supplement (Nestlé Beneprotein®)	Cooked cereals, scrambled eggs, sauces, mashed potatoes, soups, cream sauces, milk, milkshakes, cream desserts, custards, etc.
Milk (2% or 3.25% MF)	Hot cereals, soups, casseroles, hot chocolate (instead of water)
Soy beverage	Smoothies, soups
Greek yogurt	Fresh or canned fruit, vegetables, potatoes, rice, pancakes, casseroles, stews, soups, vegetable or fruit dips
Hard-boiled eggs	Sandwiches, salads, vegetables, potatoes, sauces and soups
Peanut butter or nut butter	Cookies, milkshakes, sandwiches, crackers, muffins, fruit slices, toast, ice cream
Tofu	Milkshakes, soups, casseroles, stir-fries, salads
Canned dried peas or beans, legumes and lentils (if you can tolerate these)	Casseroles, soups, stews, salads, rice, pasta and dips
Seeds and nuts (if you can tolerate these)	Salads, cereal, ice cream, yogurt
Pieces of cooked beef, pork, poultry, seafood or fish	Salads, soups, scrambled eggs, quiches, baked potato, pasta

THE NIGHT BEFORE THE SURGERY

Preoperative diet

1. The night before your surgery:

You can eat normally up to the night before your surgery.



2. Day of your surgery

➔ **For all users** – Starting from midnight the night before your surgery :

- Do not eat solid food.
- Do not consume dairy products.
- Do not consume alcohol and do not smoke.
- For the consuming of clear liquids, refer to the tables on the following page.



THE DAY OF YOUR SURGERY

At home

Your nurse has given you specific instructions:

- **You must remain fasting** (nothing to eat or drink from midnight the night before your surgery). Do not chew gum or eat candy.



You can brush your teeth but avoid swallowing the water.

OR

- **You must drink clear fluids** before the surgery.

Allowed clear fluids include:

- Water
- Juice without pulp (no pulp is mandatory)
- Coffee or black tea (no milk)



Make sure that you ONLY drink these clear fluids and nothing else.

When should I stop drinking clear fluids?

You must stop drinking these fluids the morning of your surgery. The exact time depends on when you need to arrive at the hospital that morning. Someone will call you 24 to 48 hours before your surgery will give you at what time you must arrive at the hospital.

I need to arrive at the hospital at...	I have to stop drinking clear fluids at...
Before 10 a.m.	6 a.m.
After 10 a.m.	8 a.m.
I do not have a specific time and have to wait at home to be called for my surgery.	11 a.m.

IMPORTANT:



You must follow these instructions to ensure your surgery is safe and to prevent serious complications.

If you have not followed these instructions, you must advise the nurse once you get to the hospital.

HYGIENE BEFORE YOUR SURGERY



☐ **Dexidin disinfectant soap (4%)**: The morning of the surgery, you must shower using the antimicrobial soap you purchased at the gift shop at the main entrance of Block C or Block D or at the pharmacy. You must use the soap from your chin to your toes and then rinse.

Put on clean clothes after your shower.



No makeup, no nail polish (fingers and toes), no fake nails, no fake eyelashes, no cream, deodorant or perfume/cologne, no jewelry or body piercings.

Do not shave the surgical area.



Medication


Take these medications **ONLY**
(with some water).



If all of these instructions are not respected, your surgery may be cancelled.

WHAT TO BRING TO THE HOSPITAL

- This guide.
- A valid health insurance card.
- Your hospital card.
- Your medications, drops and pumps in their original containers.
- A complete list of your medications (ask your pharmacist for this list).
- Slippers, dressing gown, clothing and comfortable shoes.
- Tissues, toothbrush and soap.
- Notebook and pencil.
- If you wear glasses, contact lenses, a hearing aid or dentures: bring your kits or containers and label them with your name.
- If you use a cane, crutches or a walker, bring them to the hospital and label them with your name.
- If you need to, bring sanitary napkins, baby diapers or incontinence products.



Please leave all your jewelry and other valuable objects at home.


The hospital is not responsible for lost or stolen items.

(The lockers do not have locks).

Rings will have to be cut off if not removed beforehand.

WHEN YOU ARRIVE AT THE SURGERY UNIT

- ☐ **If you are going to have a day surgery:** Go directly to the day surgery unit on the first floor of Block B (Room 1.165). The time of your surgery will be given to you when you arrive at the unit. **YOU WILL NOT BE STAYING AT THE HOSPITAL OVERNIGHT.**
- ☐ **If you have to stay at the hospital after your surgery:** Go to the reception area of room RC-5.



- **Only one person** can accompany you.
- After you arrive at the unit, you should expect to wait a moment until being called for your surgery. Bring something to entertain yourself if you want (something to read, a music player with headphones, etc.).



THE DAY SURGERY UNIT

At your arrival, the nurse will help you to get ready for your surgery.

She will give you an hospital gown to put on.


She will proceed to a blood test if necessary.

She will go over all preparations that you had to do before your surgery.

OPERATING ROOM

When you leave for the operating room, you must only wear the hospital gown and no other personal clothing.

You must remove your:

- 
- Glasses, **contact** lenses.
 - Underwear, **jewelry** and body piercings.
 - Dentures, hearing, **hair** piece.



Staff will direct you to the operating room.

The anesthesiologist will meet with you when you arrive in the operating room to discuss with you the most suitable methods of anesthesia and pain relief for you.

For further information about anesthesia, please read the guide on anesthesia – information guide, the nurse will provide when attending your preadmission meeting.

IN THE RECOVERY ROOM

You will wake up in the recovery room after your surgery.

No visitors are allowed in the recovery room.

Staff will place you comfortably on a stretcher or bed.

You won't be able to eat or drink right away, but the nurse will give you the go-ahead once your condition is stable.

Once you are stable and your pain is controlled, you will be transferred to the day surgery unit.

RETURN TO THE DAY SURGERY UNIT OR HOSPITAL UNIT

Staff will set you up comfortably on your stretcher or bed and will take your vital signs a few times.

The nurse will check the amount and color of your urine.

Rarely, the urologist will install a urinary catheter (tube in the bladder) that collects urine in a bag installed at the foot of your stretcher.




It is removed a few hours after you leave the operating room.



CONTROLLING YOUR PAIN

It is normal to have pain after an operation. The amount of pain is different for everyone. However, you can control your pain with the medication prescribed by your surgeon.

You will be asked to assess your pain on a scale of 0 to 10



Pain relief is important because this will help you:

- Breathe more easily.
- Move around more easily.
- Sleep better.
- Eat better.
- Recover more quickly.
- Do things that are important to you.

PAIN RELIEF TECHNICS

Analgesics (pain medication)

- You may receive injections if your pain is too intense.
- You will get your medication in tablet (pill) form once you can tolerate it or once you can eat food.




BREATHING EXERCISES

Deep breathing

****Do this as soon as you wake up from surgery****

1. Lie on your back, with your legs slightly bent. Place one hand on your stomach and the other below your breasts.



2. **Inhale slowly and deeply through your nose or mouth.** Feel your lungs inflate. Just the hand on your belly should rise.
3. Keep your lips pursed and exhale **slowly** through your mouth. This will double the length of your breath. Move your belly back in to expel the air from your lungs.



This exercise is not easy to do. Therefore, you need to practice before your operation.

Spirometer


The preadmission nurse will give you this device if you need it.

How do I use it?

Remove the device from the package. Connect the mouthpiece to the tubing. Connect the tubing to the outlet on the other side of the flow volume selector.

1. Get into a comfortable seated position.
2. Adjust the level by turning the flow volume selector to the right that will increase the difficulty of the exercise).

Tube



Flow
volume
selector

3. Hold the device upright in front of you (if you lean it to the front or back, the exercise is too easy). Exhale normally.
4. Place your lips snugly around the mouthpiece and then **inhale**. Take in enough air to lift the ball.

5. Continue inhaling to keep the ball elevated for 3 seconds. This step lets you expand your lungs as much as possible. Hold your inhalation for 3 seconds, even if the ball drops back down.
6. Then, breathe out through your mouth through pursed lips. Take a break to breathe normally, and then try again.
7. Repeat steps 4 to 6 for about 5 minutes per hour or as per your nurse's instructions.

Keep the device near you so that you remember to do the exercises. Between uses, you can keep the mouthpiece attached to the end of the tubing.

Spirometer breathing exercises help you:

- Eliminate lung secretions to prevent respiratory complications.
- Regain and maintain good lung expansion.
- Stimulate the breathing reflex, which is slowed by anesthesia and pain medication.
- Improve your well-being and resume your usual activities more quickly.

CIRCULATION EXERCISES¹

These exercises encourages blood circulation in your legs while you are lying down. They are very important because they can prevent serious complications, such as blood clots in the veins of your legs (thrombophlebitis).

- **Toe flexion and extension**

While lying on your back or sitting with your legs stretched out, point your toes to the foot of the bed and then point them toward your chin. Repeat the exercise 30 times a minute for 1 to 2 minutes, every 2 hours.



- **Ankle rotations**

While lying on your back or sitting, make ankle circles from left to right and then from right to left. Repeat this exercise 30 times a minute for 1 to 2 minutes, every 2 hours.




¹ These circulation exercises are based on those developed by Paradis and Poissant.

GETTING UP

When you get up for the first time, a staff member will be there to assist you; however, you should only get up at your own pace. You need to walk and increase the distance you walk each time you get up. Increase your pace gradually.

To help you get in and out of bed, you need to raise slightly the head of your bed.



1. Turn toward your non-operated side.
2. Push against the mattress using your elbow on the non-operated side and your other hand to sit up on the edge of the bed. Slide your legs over the bed at the same time.
3. Stay in this position for a few minutes. Take a few deep breaths and move your feet around.
4. If you do not feel well, tell the nurse or care attendant.

Staff will help you sit in an armchair if you need to.

YOUR DISCHARGE FROM THE UNIT

- Your surgeon is the one who will discharge you.
- You must ask another adult to come pick you up, since you cannot drive after your operation. You must plan a ride home.
- If you live by yourself, it is a good idea to ask another adult to stay with you for 24 hours for your safety reasons.
- The nurse may give you a prescription for pain medication, which you must get at your pharmacy. Your nurse will also give you a pamphlet about what you need to know if you need to take a narcotic medication for pain.
- The nurse will give you a follow-up appointment with your surgeon. You must absolutely go to this appointment, even if you feel well.



Surgeon name: _____

Date & time of appointment: _____

- You will receive a proof of hospitalization or medical leave from work form if you need one. You must notify your surgeon if you need these documents.
- If you have insurance forms to complete, contact your urologist's secretary at his or her private office. (See Urologists' Reference on p. 25).




**All forms must be sent to the private office.
No forms will be filled out at the hospital on the day of the surgery.**

ONCE YOU GET BACK HOME – INSTRUCTIONS

Your incision

The procedure is performed through the natural route of the urethra, so no incision on the belly is required.



HYGIENE

1. You can take a shower or bath after your procedure.
2. You may feel a burning sensation when urinating and see either pinkish or reddish urine and some blood clots. This is normal after this operation and is from the instruments passing through the urinary tract and the double-J stent.
3. Common symptoms from this procedure include back pain, bladder spasms, frequent urination, and the feeling of your bladder not emptying.
4. Staying hydrated (one 250-ml glass per hour during the day when you are awake, if you have no medical restrictions against this amount of fluid intake) up to a maximum of 8 to 10 glasses per day and frequent urination will help relieve these symptoms. Taking a hot bath and your prescribed pain relievers will also keep you comfortable.



GOING BACK TO WORK

Your surgeon and nurse will give you more details about your recovery, which will depend on your procedure and the type of work that you do.



BREASTFEEDING

If you are breastfeeding, ask the surgeon or nurse if you can continue.



Generally, you need to wait 2 to 3 hours after having general anesthesia before breastfeeding your baby. As soon as you return home, you can breastfeed if you feel alert and comfortable

NUTRITION AND HYDRATION

Generally, you can eat normally after your surgery. Start with light meals and gradually increase as you can tolerate.

If you need to adjust your diet after your operation, the surgeon and nurse will give you instructions. If you have questions, please do not hesitate to ask them.



If you have nausea (you feel sick to your stomach), start by drinking clear fluids and gradually increase the amount and change the texture of the foods you eat as you can tolerate them.

To avoid constipation, which can be caused by pain medication:

- Eat plenty of fiber (grains, whole-grain bread, fruit, vegetables, etc.).
- Drink 7 to 8 glasses of water a day (unless you have a medical restriction).
- Walking can help with bowel function.



If, despite these tips, you are unable to have a bowel movement:

- You can use a mild laxative such as Metamucil®, Colace®, Lax A day® or Prodiem® at a pharmacy. Ask your pharmacist for advice.



If you have not had a bowel movement for at least 3 days despite these tips, consult a health care professional (family doctor, pharmacist, Info-Santé at 811).

ACTIVITIES



You can drive when:

- **You no longer feel dizzy.**
- **You no longer have pain and you have stopped taking narcotic medications for at least 24 hours.**

- Depending on your procedure, your surgeon or nurse may give you other instructions that are appropriate for your situation. Do not hesitate to ask questions.
- You need to stay active after your operation, but you also need to rest. It is normal to feel tired.
- Pain should not prevent you from your daily activities such as getting dressed, washing yourself or eating. Take your pain medication if the pain is too intense, and take it at least 30 minutes before doing exercise, if you need to.
- Walking is one of the best exercises, except if restricted by your surgeon. Increase the distance you walk everyday and alternate with periods of rest. Avoid vigorous exercise, sudden movements, or contact sports.
- You must restrict your activities for the first week. You can resume your activities gradually thereafter, depending on your tolerance. You can go up and down the stairs.
- A double-J stent often increases bladder pain, frequency of urination, and an urgent need to urinate. Patients also commonly feel pain in their side when urinating or when their bladder is full. They may also see blood in the urine. These symptoms can be aggravated by activity.
- Avoid vigorous exercise, sudden movement, contact sports, and travelling. To prevent bleeding, avoid lifting heavy weights (10 kg / 20 lbs.) for 2 weeks after your follow-up visit with your urologist and not until your urine starts running clear.
- You can resume sexual activity about 1 week after your surgery and once your pain has subsided.



COMPLICATIONS

If you have difficulty breathing:

**Immediately call
Urgences-Santé at 9-1-1**



If you have one or more of the following signs or symptoms:




Fever (38.5 °C or 101 °F or higher) for more than 24 hours

Your pain increases and is not relieved by medication.



You have cramps or constant pain in your calf.

- An inability to urinate after several hours and severe abdominal pain.
- Constant dark-red urine. Urinating multiple times without the urine clearing up (despite abundant hydration).
- Abundant blood clots in the urine
- Nausea and vomiting
- Hard and swollen abdomen



Contact a nurse at Info-Santé by calling 811 at any time (24 hours a day).

For all other questions: Contact one of the resources listed on next page



RESOURCES

For emergencies, call 911
Info-Santé – CLSC, call 811
24 hours a day, 7 days a week

Outpatient Clinic

Preadmission (preoperative only) 450 975-5566
Urology 450 975-5913, poste 4

Private offices of Urologists in Laval

Polyclinique médicale Concorde
Adress: 300, blvd de la Concorde East, Laval (Québec)
Téléphone450 667-5310

CLSC

Région de Laval

Accueil première ligne..... 450 627-2530, poste 64922
CLSC des Mille-Îles450 661-2572
CLSC du Ruisseau-Papineau450 682-5690
CLSC et CHSLD Sainte-Rose.....450 622-5110
CLSC de l'Ouest-de-l'île450 627-2530
CLSC et CHSLD Idola-Saint-Jean.....450 668-1803

Région des Laurentides

Centre intégré de santé et de services sociaux des Laurentides :
Thérèse de Blainville.....450 433-2777
Des sommets.....819 324-4000
St-Jérôme450 432-2777
Pays d'en haut450 229-6601
Jean-Olivier Chenier450 433-2777
Argenteuil.....450 562-3761
Antoine Labelle819 275-2118

Région de Lanaudière

Lanaudière Sud.....450 654-2572
Lanaudière Nord450 839-3864

BIBLIOGRAPHY

- McGill University Health Centre. Patient Education Office. (2013). *A Guide to Your Minimally Invasive Hysterectomy*.
- Centre de santé et de services sociaux de Laval.(2006). *Insertion of a double-J stent*.
- Canadian Urological Association. (2014). *Ureteroscopy*.
https://www.cua.org/sites/default/files/Flipbooks/PIB/PIB43_en/mobile/index.html
- Centre de santé et de services sociaux Jardins Roussillon, Preadmission Clinic. (2009). *Ureteroscopy, stone removal, and installation of a double-J stent*.
- Vinet, M. (2006, December). *Removal of ureteral and bladder stones* [Brochure]. Centre de santé et de services sociaux de Laval, Preadmission Clinic. [Unpublished manuscript].

[illegible]

**Centre intégré
de santé
et de services sociaux
de Laval**

Québec 

November 2022
Direction des services professionnels
62-100-098