

A note on continuous p -defense-sum problems

Nenad Mladenović

*GERAD and École des Hautes Études Commerciales
3000 chemin de la Côte-Sainte-Catherine
Montréal, Canada H3T 2A7 SMG and ISRO
Université Libre de Bruxelles
Bruxelles, Belgium*

Frank Plastria

*Dept. Management Informatics
Vrije Universiteit Brussel
Brussel, Belgium*

September, 2000

Les Cahiers du GERAD

G-2000-43

Copyright © 2000 GERAD

Abstract

The continuous p -defense-sum problem consists of locating p facilities in a convex polyhedron, such that the sum of the distances among all their pairs is maximized. We indicate that it is sufficient to search for optimal sites at the polyhedron's vertices only, and show that the optimal solution can be degenerate, i.e., more than one facility being located at the same point. An integer programming formulation is also given, taking the possible degeneracy into account.

Keywords: Optimization, Location, p -defense-sum, degeneracy.

Résumé

Le problème de p -défense-somme continu consiste à localiser p facilités dans un polyèdre convexe de sorte que la somme des distances entre paires de facilités soit maximisée. Nous montrons qu'il suffit de choisir des sites optimaux parmi des sommets de ce polyèdre, et que la solution optimale peut être dégénérée, c'est-à-dire que plus d'une facilité peut être localisée en le même point. Nous donnons aussi une formulation de programmation entière, tenant compte de la dégénérescence éventuelle.

Mots-clés: Optimisation, Localisation, p -défense-somme, dégénérescence.

Acknowledgment: The first author was supported by the *Research in Brussels action* grant.

The discrete p -defense-sum problem is one of the known noxious facility location problems. Among a given set F of n potential undesirable facility sites (nuclear waste, garbage dumpsites, missile silos, chemical plants etc.), find p sites such that the sum of distances among all pairs of facilities is maximized. It is usually formulated as

$$\max_{X \subseteq F} \sum_{i \in X} \sum_{j \in X} d_{ij} \quad (1)$$

$$\text{subject to } |X| = p,$$

where $|F| = n$ and d_{ij} are distances (shortest paths or some other appropriate metric). This strongly NP-hard problem [5] is formulated as an 0-1 integer program in [7], with $O(n^2)$ constraints using $O(n^2)$ 0-1 variables and in [3], involving $O(n)$ variables and $O(n)$ constraints. An exact solution method based on branch and bound is suggested in [1], while heuristic methods are proposed for example in [2], [6] and [4].

The continuous version we shall consider here consists in finding p facilities in a polyhedron $S \subset R^q$ and with use of any kind of norm or gauge distance evaluation (see e.g. [8], e.g. Euclidean distances. It can be formulated as follows:

$$\max \sum_{i=1}^p \sum_{j=1}^p d_{ij}(x_i, x_j) \quad (2)$$

subject to

$$x_j \in S = \{x \in R^q \mid Ax \leq b\}, \quad j = 1, \dots, p.$$

where $b \in R^m$, $A \in R^{mq}$ and each d_{ij} is some gauge on R^q .

Proposition 1 *There exists some optimal solution x_j^* , $j = 1, \dots, p$ of the continuous p -defense-sum problem (2), with each point x_j^* located at an extreme point of polyhedron S .*

Proof. Let $X = (x_1, x_2, \dots, x_p) \in R^{qp}$ and let $S^p = S \times S \times \dots \times S$. Then (2) can be expressed as

$$\max_{X \in S^p} g(X) = \sum_{i=1}^p \sum_{j=1}^p d_{ij}(x_i, x_j) \quad (3)$$

Since $d_{ij}(x_i, x_j)$ is convex function of X , (see for example [8], $g(X)$ is convex as well (as a sum of convex functions). Now (3) is a convex maximization problem and thus its optimal value is reached at some extreme point X^* of S^p . Extreme points of

the product of convex sets are products of extreme points, i.e. $\text{ext}(S^p) = (\text{ext}(S))^p$, hence $(x_1^*, x_2^*, \dots, x_p^*) = (e_1, e_2, \dots, e_p)$, where $e_j \in \text{ext}(S)$. q.e.d.

From Proposition 1 it is clear that continuous p -defense-sum problems (2) can be reformulated as discrete maximization problems. The next proposition shows that the thus obtained problem is not equivalent to the usual discrete p -defense-sum problem (1). We first define a *degenerate* solution of (2) as a solution where not all points x_j are different.

Proposition 2 *The optimal solution of the continuous p -defense-sum problem (2) can be degenerate, if $p > 3$.*

Proof. It is easy to see that solution is never degenerate if $p \leq 3$ and $|S| > 1$. We construct a counter-example with $n = 4$ and $p = 4$. Let S be defined by the following extreme points: e_1 is the center of the ball with radius 10; three other extreme points are located on the boundary of this ball with distances among them equal to 1, 1 and 2 respectively. If $(x_1, x_2, x_3, x_4) = (e_1, e_2, e_3, e_4)$, the corresponding objective function value is equal to $10 + 10 + 10 + 2 + 1 + 1 = 34$. But if $(x_1, x_2, x_3, x_4) = (e_1, e_2, e_3, e_1)$, then we have $f = 10 + 10 + 0 + 1 + 10 + 10 = 41$. Thus, a larger (better) value is obtained in the degenerate solution since e_1 is repeated twice. q.e.d.

The difference between the two formulations is of course the fact that in a discrete problem it is easy to impose the (implicit) additional constraint that all chosen facility sites should be different, while in a continuous problem this is much less evident. Indeed, constraints such as $x_i \neq x_j$ define open feasible sets, and this leads to the possibility of ill formulation without optimal solutions, as clearly shown by previous counterexample.

In order to obtain a discrete formulation of the continuous p -defense-sum problem (2), the set of extreme points of S is copied $p - 3$ times to allow duplications (Proposition 2). Therefore the cardinality of the set of possible facility locations F is $n = |F| = n' + (p - 3) \cdot n' = (p - 2) \cdot n'$, where $n' = |\text{ext}(S)|$. The new $p - 3$ levels of the extreme points set $\text{ext}(S)$ are constructed such that holds

$$d(x_{i+\ell n'}, x_{j+kn'}) = d(x_i, x_j), \forall i, j \in \text{ext}(S), \forall \ell, k = 1, \dots, p - 3. \quad (4)$$

Now combinatorial formulation (1) can be applied.

The integer programming formulation can be given as 0-1 *quadratic knapsack* problem:

$$\max \sum_{\ell=1}^{p-2} \sum_{k=1}^{p-2} \sum_{i=1}^{n'} \sum_{j=1}^{n'} d(x_i, x_j) y_{i+(\ell-1)n'} \cdot y_{j+(k-1)n'} \quad (5)$$

subject to

$$\sum_{k=1}^n y_k = p$$

$$y_k \in \{0, 1\}, \quad 1 \leq k \leq n,$$

where y_k is set to 1 if the candidate point k is selected and 0 otherwise.

There is, however, a combinatorial problem with this formulation, in the sense that a same solution may be encoded in different ways, by selecting a same site at other levels, leading to many symmetrical equivalent solutions. One way to avoid this is to break this symmetry by imposing that some site may be used at a given level only when all previous levels have been used also, which may be expressed by the additional symmetry breaking constraints

$$y_{i+\ell n'} \leq y_{i+kn'} \quad \text{for all } i \text{ and all } \ell < k.$$

References

- [1] ERKUT, E., BAPTIE, T., and HOHENBALKEN, B., *The discrete p-maxian location problem*, Computers and Operations Research, Vol.17, pp.51-61, 1990.
- [2] ERKUT, E., and NEUMAN, S., *Analytic models for locating undesirable facilities*, European Journal of Operational Research, Vol.40, pp.275-291, 1989.
- [3] ERKUT, E., and NEUMAN, S., *Comparison of four models for dispersing facilities*, INFOR, Vol.29, pp.68-85, 1991.
- [4] ERKUT E., ULKUSAL, Y., and O. YENICERIOGLU, O., *A comparison of p-dispersion heuristics*, Computers and Operations Research, Vol.21, pp.1103-1113, 1994.
- [5] HANSEN, P., and MOON, I., *Dispersing facilities on a network*, Presentation at the TIMS/ORSA Washington Joint National Meeting, 1988.
- [6] KINCAID, R., *Good solutions to discrete noxious location problems via meta-heuristics*, Annals of Operations Research, Vol.40, pp.265-281, 1992.
- [7] KUBY, M., *Programming models for facility dispersion: The p-dispersion and maxisum dispersion problems*, Geographical Analysis, Vol.19, pp.315-329, 1987.

- [8] PLASTRIA, F., *Continuous Location Problems*, Facility Location: A Survey of Applications and Methods, Edited by Z. Drezner, Springer-Verlag, 1995, pp.225–262, 1995.