

Largest small polygons: A sequential convex optimization approach

C. Bingane

G-2020-50

October 2020

La collection *Les Cahiers du GERAD* est constituée des travaux de recherche menés par nos membres. La plupart de ces documents de travail a été soumis à des revues avec comité de révision. Lorsqu'un document est accepté et publié, le pdf original est retiré si c'est nécessaire et un lien vers l'article publié est ajouté.

Citation suggérée : C. Bingane (Octobre 2020). Largest small polygons: A sequential convex optimization approach, Rapport technique, Les Cahiers du GERAD G-2020-50, GERAD, HEC Montréal, Canada.

Avant de citer ce rapport technique, veuillez visiter notre site Web (<https://www.gerad.ca/fr/papers/G-2020-50>) afin de mettre à jour vos données de référence, s'il a été publié dans une revue scientifique.

La publication de ces rapports de recherche est rendue possible grâce au soutien de HEC Montréal, Polytechnique Montréal, Université McGill, Université du Québec à Montréal, ainsi que du Fonds de recherche du Québec – Nature et technologies.

Dépôt légal – Bibliothèque et Archives nationales du Québec, 2020
– Bibliothèque et Archives Canada, 2020

The series *Les Cahiers du GERAD* consists of working papers carried out by our members. Most of these pre-prints have been submitted to peer-reviewed journals. When accepted and published, if necessary, the original pdf is removed and a link to the published article is added.

Suggested citation: C. Bingane (October 2020). Largest small polygons: A sequential convex optimization approach, Technical report, Les Cahiers du GERAD G-2020-50, GERAD, HEC Montréal, Canada.

Before citing this technical report, please visit our website (<https://www.gerad.ca/en/papers/G-2020-50>) to update your reference data, if it has been published in a scientific journal.

The publication of these research reports is made possible thanks to the support of HEC Montréal, Polytechnique Montréal, McGill University, Université du Québec à Montréal, as well as the Fonds de recherche du Québec – Nature et technologies.

Legal deposit – Bibliothèque et Archives nationales du Québec, 2020
– Library and Archives Canada, 2020

Largest small polygons: A sequential convex optimization approach

Christian Bingane

GERAD & Department of Mathematics and Industrial Engineering, Polytechnique Montréal (Québec) Canada, H3C 3A7

christian.bingane@polymtl.ca

October 2020
Les Cahiers du GERAD
G–2020–50

Copyright © 2020 GERAD, Bingane

Les textes publiés dans la série des rapports de recherche *Les Cahiers du GERAD* n'engagent que la responsabilité de leurs auteurs. Les auteurs conservent leur droit d'auteur et leurs droits moraux sur leurs publications et les utilisateurs s'engagent à reconnaître et respecter les exigences légales associées à ces droits. Ainsi, les utilisateurs:

- Peuvent télécharger et imprimer une copie de toute publication du portail public aux fins d'étude ou de recherche privée;
- Ne peuvent pas distribuer le matériel ou l'utiliser pour une activité à but lucratif ou pour un gain commercial;
- Peuvent distribuer gratuitement l'URL identifiant la publication.

Si vous pensez que ce document enfreint le droit d'auteur, contactez-nous en fournissant des détails. Nous supprimerons immédiatement l'accès au travail et enquêterons sur votre demande.

The authors are exclusively responsible for the content of their research papers published in the series *Les Cahiers du GERAD*. Copyright and moral rights for the publications are retained by the authors and the users must commit themselves to recognize and abide the legal requirements associated with these rights. Thus, users:

- June download and print one copy of any publication from the public portal for the purpose of private study or research;
- June not further distribute the material or use it for any profit-making activity or commercial gain;
- June freely distribute the URL identifying the publication.

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract: A small polygon is a polygon of unit diameter. The maximal area of a small polygon with $n = 2m$ vertices is not known when $m \geq 7$. Finding the largest small n -gon for a given number $n \geq 3$ can be formulated as a nonconvex quadratically constrained quadratic optimization problem. We propose to solve this problem with a sequential convex optimization approach, which is an ascent algorithm guaranteeing convergence to a locally optimal solution. Numerical experiments on polygons with up to $n = 128$ sides suggest that the optimal solutions obtained are near-global. Indeed, for even $6 \leq n \leq 12$, the algorithm proposed in this work converges to known global optimal solutions found in the literature.

Keywords: Planar geometry, small polygons, maximal area, quadratically constrained quadratic optimization, sequential convex optimization, concave-convex procedure.

Acknowledgments: The author thanks Charles Audet, Professor at Polytechnique Montreal, for helpful discussions on largest small polygons and helpful comments on early drafts of this paper.

1 Introduction

The *diameter* of a polygon is the largest Euclidean distance between pairs of its vertices. A polygon is said to be *small* if its diameter equals one. For a given integer $n \geq 3$, the maximal area problem consists in finding the small n -gon with the largest area. The problem was first investigated by Reinhardt [1] in 1922. He proved that

- when n is odd, the regular small n -gon is the unique optimal solution;
- when $n = 4$, there are infinitely many optimal solutions, including the small square;
- when $n \geq 6$ is even, the regular small n -gon is not optimal.

The maximal area is known for even $n \leq 12$. In 1961, Bieri [2] found the largest small 6-gon, assuming the existence of an axis of symmetry. In 1975, Graham [3] independently constructed the same 6-gon, represented in Figure 2c. In 2002, Audet, Hansen, Messine, and Xiong [4] combined Graham's strategy with global optimization methods to find the largest small 8-gon, illustrated in Figure 3c. In 2013, Henrion and Messine [5] found the largest small 10- and 12-gons by also solving globally a nonconvex quadratically constrained quadratic optimization problem. They also found the largest small axially symmetrical 14- and 16-gons. In 2017, Audet [6] showed that the regular small polygon has the maximal area among all equilateral small polygons. In 2020, Audet, Hansen, and Svrtan [7] determined analytically the largest small axially symmetrical 8-gon.

The diameter graph of a small polygon is defined as the graph with the vertices of the polygon, and an edge between two vertices if the distance between these vertices equals one. Graham [3] conjectured that, for even $n \geq 6$, the diameter graph of a small n -gon with maximal area has a cycle of length $n - 1$ and one additional edge from the remaining vertex. The case $n = 6$ was proven by Graham himself [3] and the case $n = 8$ by Audet, Hansen, Messine, and Xiong [4]. In 2007, Foster and Szabo [8] proved Graham's conjecture for all even $n \geq 6$. Figure 1, Figure 2, and Figure 3 show diameter graphs of some small polygons. The solid lines illustrate pairs of vertices which are unit distance apart.

In addition to exact results and bounds, uncertified largest small polygons have been obtained both by metaheuristics and nonlinear optimization. Assuming Graham's conjecture and the existence of an axis of symmetry, Mossinghoff [9] in 2006 constructed large small n -gons for even $6 \leq n \leq 20$. In 2018, using a formulation based on polar coordinates, Pinter [10] presented numerical solutions estimates of the maximal area for even $6 \leq n \leq 80$. However, the solutions obtained by Pinter are not optimal for even $n \geq 32$.

The maximal area problem can be formulated as a nonconvex quadratically constrained quadratic optimization problem. In this work, we propose to solve it with a sequential convex optimization approach, also known as the concave-convex procedure [11, 12]. This approach is an ascent algorithm guaranteeing convergence to a locally optimal solution. Numerical experiments on polygons up to $n = 128$ sides suggest that the optimal solutions obtained are near-global. Indeed, without assuming Graham's conjecture nor the existence of an axis of symmetry in our quadratic formulation, optimal n -gons obtained with the algorithm proposed in this work verify both conditions within the limit of numerical computations. Moreover, for even $6 \leq n \leq 12$, this algorithm converges to known global optimal solutions. The algorithm is implemented as a MATLAB-based package, OPTIGON, which is available on GitHub [13]. OPTIGON requires that CVX [14] be installed.

The remainder of this paper is organized as follows. In Section 2, we recall principal results on largest small polygons. Section 3 presents the quadratic formulation of the maximal area problem and the sequential convex optimization approach to solve it. We report in Section 4 computational results. Section 5 concludes the paper.




Figure 1: Two small 4-gons ($P_4, A(P_4)$)




Figure 2: Three small 6-gons ($P_6, A(P_6)$)




Figure 3: Three small 8-gons ($P_8, A(P_8)$)

2 Largest small polygons

Let $A(P)$ denote the area of a polygon P . Let R_n denote the regular small n -gon. We have

$$A(R_n) = \begin{cases} \frac{n}{2} \left(\sin \frac{\pi}{n} - \tan \frac{\pi}{2n} \right) & \text{if } n \text{ is odd,} \\ \frac{n}{8} \sin \frac{2\pi}{n} & \text{if } n \text{ is even.} \end{cases}$$

We remark that $A(R_n) < A(R_{n-1})$ for all even $n \geq 6$ [15]. This suggests that R_n does not have maximum area for any even $n \geq 6$. Indeed, when n is even, we can construct a small n -gon with a larger area than R_n by adding a vertex at distance 1 along the mediatix of an angle in R_{n-1} . We denote this n -gon by R_{n-1}^+ and we have

$$A(R_{n-1}^+) = \frac{n-1}{2} \left(\sin \frac{\pi}{n-1} - \tan \frac{\pi}{2n-2} \right) + \sin \frac{\pi}{2n-2} - \frac{1}{2} \sin \frac{\pi}{n-1}.$$

Theorem 1 (Reinhardt [1]) For all $n \geq 3$, let A_n^* denote the maximal area among all small n -gons and let $\bar{A}_n := \frac{n}{2} \left(\sin \frac{\pi}{n} - \tan \frac{\pi}{2n} \right)$.

- When n is odd, $A_n^* = \bar{A}_n$ is only achieved by R_n .
- $A_4^* = 0.5 < \bar{A}_4$ is achieved by infinitely many 4-gons, including R_4 and R_3^+ illustrated in Figure 1.
- When $n \geq 6$ is even, $A(R_n) < A_n^* < \bar{A}_n$.

The maximal area A_n^* is known for even $n \leq 12$. Using geometric arguments, Graham [3] determined analytically the largest small 6-gon, represented in Figure 2c. Its area $A_6^* \approx 0.674981$ is about 3.92% larger than $A(R_6) \approx 0.649519$. The approach of Graham, combined with methods of global optimization, has been followed by [4] to determine the largest small 8-gon, represented in Figure 3c. Its area $A_8^* \approx 0.726868$ is about 2.79% larger than $A(R_8) \approx 0.707107$. Henrion and Messine [5] found that $A_{10}^* \approx 0.749137$ and $A_{12}^* \approx 0.760730$.

For all even $n \geq 6$, let U_n denote the largest small n -gon.

Theorem 2 (Graham [3], Foster and Szabo [8]) For even $n \geq 6$, the diameter graph of U_n has a cycle of length $n - 1$ and one additional edge from the remaining vertex.

Conjecture 1 For even $n \geq 6$, U_n has an axis of symmetry corresponding to the pending edge in its diameter graph.

From Theorem 2, we note that R_{n-1}^+ has the same diameter graph as the largest small n -gon U_n . Conjecture 1 is only proven for $n = 6$ and this is due to Yuan [16]. However, the largest small polygons obtained by [4] and [5] are a further evidence that the conjecture may be true.

3 Nonconvex quadratically constrained quadratic optimization

We use cartesian coordinates to describe an n -gon P_n , assuming that a vertex v_i , $i = 0, 1, \dots, n - 1$, is positioned at abscissa x_i and ordinate y_i . Placing the vertex v_0 at the origin, we set $x_0 = y_0 = 0$. We also assume that the n -gon P_n is in the half-plane $y \geq 0$ and the vertices v_i , $i = 1, 2, \dots, n - 1$, are arranged in a counterclockwise order as illustrated in Figure 4, i.e., $y_{i+1}x_i \geq x_{i+1}y_i$ for all $i = 1, 2, \dots, n - 2$. The maximal area problem can be formulated as follows

$$\max_{\mathbf{x}, \mathbf{y}, \mathbf{u}} \sum_{i=1}^{n-2} u_i \tag{1a}$$

$$\text{s. t. } (x_j - x_i)^2 + (y_j - y_i)^2 \leq 1 \quad \forall 1 \leq i < j \leq n - 1, \tag{1b}$$

$$x_i^2 + y_i^2 \leq 1 \quad \forall 1 \leq i \leq n - 1, \tag{1c}$$

$$y_i \geq 0 \quad \forall 1 \leq i \leq n - 1, \tag{1d}$$

$$2u_i \leq y_{i+1}x_i - x_{i+1}y_i \quad \forall 1 \leq i \leq n - 2, \tag{1e}$$

$$u_i \geq 0 \quad \forall 1 \leq i \leq n - 2. \tag{1f}$$

At optimality, for all $i = 1, 2, \dots, n - 2$, $u_i = (y_{i+1}x_i - x_{i+1}y_i)/2$, which corresponds to the area of the triangle $v_0v_iv_{i+1}$. It is important to note that, unlike what was done in [4, 5], this formulation does not make the assumption of Graham's conjecture, nor of the existence of an axis of symmetry.

Problem (1) is a nonconvex quadratically constrained quadratic optimization problem and can be reformulated as a difference-of-convex optimization (DCO) problem of the form

$$\max_{\mathbf{z}} g_0(\mathbf{z}) - h_0(\mathbf{z}) \tag{2a}$$

$$\text{s. t. } g_i(\mathbf{z}) - h_i(\mathbf{z}) \geq 0 \quad \forall 1 \leq i \leq m, \tag{2b}$$

where g_0, \dots, g_m and h_0, \dots, h_m are convex quadratic functions. We note that the feasible set

$$\Omega := \{ \mathbf{z} : g_i(\mathbf{z}) - h_i(\mathbf{z}) \geq 0, i = 1, 2, \dots, m \}$$




Figure 4: Definition of variables: Case of $n = 6$ vertices

is compact with a nonempty interior, which implies that $g_0(\mathbf{z}) - h_0(\mathbf{z}) < \infty$ for all $\mathbf{z} \in \Omega$.

For a fixed \mathbf{c} , we have $\underline{g}_i(\mathbf{z}; \mathbf{c}) := g_i(\mathbf{c}) + \nabla g_i(\mathbf{c})^T (\mathbf{z} - \mathbf{c}) \leq g_i(\mathbf{z})$ for all $i = 0, 1, \dots, m$. Then the following problem

$$\max_{\mathbf{z}} \quad \underline{g}_0(\mathbf{z}; \mathbf{c}) - h_0(\mathbf{z}) \quad (3a)$$

$$\text{s. t.} \quad \underline{g}_i(\mathbf{z}; \mathbf{c}) - h_i(\mathbf{z}) \geq 0 \quad \forall 1 \leq i \leq m \quad (3b)$$

is a convex restriction of the DCO problem (2) as stated by Proposition 1. Constraint (1e) is equivalent to

$$(y_{i+1} - x_i)^2 + (x_{i+1} + y_i)^2 + 8u_i \leq (y_{i+1} + x_i)^2 + (x_{i+1} - y_i)^2$$

for all $i = 1, 2, \dots, n - 2$. For a fixed $(\mathbf{a}, \mathbf{b}) \in \mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$, if we replace (1e) in (1) by

$$(y_{i+1} - x_i)^2 + (x_{i+1} + y_i)^2 + 8u_i \leq 2(b_{i+1} + a_i)(y_{i+1} + x_i) - (b_{i+1} + a_i)^2 + 2(a_{i+1} - b_i)(x_{i+1} - y_i) - (a_{i+1} - b_i)^2$$

for all $i = 1, 2, \dots, n - 2$, we obtain a convex restriction of the maximal area problem.

Proposition 1 *If \mathbf{z} is a feasible solution of (3) then \mathbf{z} is a feasible solution of (2).*

Proof. Let \mathbf{z} be a feasible solution of (3), i.e., $\underline{g}_i(\mathbf{z}; \mathbf{c}) - h_i(\mathbf{z}) \geq 0$ for all $i = 1, 2, \dots, m$. Then $g_i(\mathbf{z}) - h_i(\mathbf{z}) \geq \underline{g}_i(\mathbf{z}; \mathbf{c}) - h_i(\mathbf{z}) \geq 0$ for all $i = 1, 2, \dots, m$. Thus, \mathbf{z} is a feasible solution of (2). \square

Proposition 2 *If \mathbf{c} is a feasible solution of (2) then (3) is a feasible problem. Moreover, if \mathbf{z}^* is an optimal solution of (3) then $g_0(\mathbf{c}) - h_0(\mathbf{c}) \leq g_0(\mathbf{z}^*) - h_0(\mathbf{z}^*)$.*

Proof. Let \mathbf{c} be a feasible solution of (2), i.e., $g_i(\mathbf{c}) - h_i(\mathbf{c}) \geq 0$ for all $i = 1, 2, \dots, m$. Then there exists $\mathbf{z} = \mathbf{c}$ such that $\underline{g}_i(\mathbf{c}; \mathbf{c}) - h_i(\mathbf{c}) = g_i(\mathbf{c}) - h_i(\mathbf{c}) \geq 0$ for all $i = 1, 2, \dots, m$. Thus, (3) is a feasible problem. Moreover, if \mathbf{z}^* is an optimal solution of (3), we have $g_0(\mathbf{c}) - h_0(\mathbf{c}) = \underline{g}_0(\mathbf{c}; \mathbf{c}) - h_0(\mathbf{c}) \leq \underline{g}_0(\mathbf{z}^*; \mathbf{c}) - h_0(\mathbf{z}^*) \leq g_0(\mathbf{z}^*) - h_0(\mathbf{z}^*)$. \square

From Proposition 2, the optimal small n -gon (\mathbf{x}, \mathbf{y}) obtained by solving a convex restriction of Problem (1) constructed around a small n -gon (\mathbf{a}, \mathbf{b}) has a larger area than this one. Proposition 3 states that if (\mathbf{a}, \mathbf{b}) is the optimal n -gon of the convex restriction constructed around itself, then it is a local optimal n -gon for the maximal area problem.

Proposition 3 *Let \mathbf{c} be a feasible solution of (2). We suppose that $\underline{\Omega}(\mathbf{c}) := \{\mathbf{z} : \underline{g}_i(\mathbf{z}; \mathbf{c}) - h_i(\mathbf{z}) \geq 0, i = 1, 2, \dots, m\}$ satisfies Slater condition. If \mathbf{c} is an optimal solution of (3) then \mathbf{c} is a critical point of (2).*

Proof. If \mathbf{c} is an optimal solution of (3) then there exist m scalars $\mu_1, \mu_2, \dots, \mu_m$ such that

$$\begin{aligned} \nabla g_0(\mathbf{c}; \mathbf{c}) + \sum_{i=1}^m \mu_i \nabla g_i(\mathbf{c}; \mathbf{c}) &= \nabla h_0(\mathbf{c}) + \sum_{i=1}^m \mu_i \nabla h_i(\mathbf{c}), \\ g_i(\mathbf{c}; \mathbf{c}) &\geq h_i(\mathbf{c}) && \forall i = 1, 2, \dots, m, \\ \mu_i &\geq 0 && \forall i = 1, 2, \dots, m, \\ \mu_i g_i(\mathbf{c}; \mathbf{c}) &= \mu_i h_i(\mathbf{c}) && \forall i = 1, 2, \dots, m. \end{aligned}$$

Since $\underline{g}_i(\mathbf{c}; \mathbf{c}) = g_i(\mathbf{c})$ and $\nabla \underline{g}_i(\mathbf{c}; \mathbf{c}) = \nabla g_i(\mathbf{c})$ for all $i = 0, 1, \dots, m$, we conclude that \mathbf{c} is a critical point of (2). \square

We propose to solve the DCO problem (2) with a sequential convex optimization approach given in Algorithm 1, also known as concave-convex procedure. A proof of showing that a sequence $\{\mathbf{z}_k\}_{k=0}^{\infty}$ generated by Algorithm 1 converges to a critical point \mathbf{z}^* of the original DCO problem (2) can be found in [11, 12].

Model 1 Sequential convex optimization

- 1: Initialization: choose a feasible solution \mathbf{z}_0 and a stopping criteria $\varepsilon > 0$.
 - 2: $\mathbf{z}_1 \in \arg \max\{g_0(\mathbf{z}; \mathbf{z}_0) - h_0(\mathbf{z}); \underline{g}_i(\mathbf{z}; \mathbf{z}_0) - h_i(\mathbf{z}) \geq 0, i = 1, 2, \dots, m\}$
 - 3: $k := 1$
 - 4: **while** $\frac{\|\mathbf{z}_k - \mathbf{z}_{k-1}\|}{\|\mathbf{z}_k\|} > \varepsilon$ **do**
 - 5: $\mathbf{z}_{k+1} \in \arg \max\{g_0(\mathbf{z}; \mathbf{z}_k) - h_0(\mathbf{z}); \underline{g}_i(\mathbf{z}; \mathbf{z}_k) - h_i(\mathbf{z}) \geq 0, i = 1, 2, \dots, m\}$
 - 6: $k := k + 1$
 - 7: **end while**
-

4 Computational results

Problem (1) was solved in MATLAB using CVX 2.2 with MOSEK 9.1.9 and `default precision` (tolerance $\varepsilon = 1.49 \times 10^{-8}$). All the computations were carried out on an Intel(R) Core(TM) i7-3540M CPU @ 3.00 GHz computing platform. Algorithm 1 was implemented as a MATLAB package: OPTIGON, which is freely available at <https://github.com/cbingane/optigon>. OPTIGON requires that CVX be installed. CVX is a MATLAB-based modeling system for convex optimization, which turns MATLAB into a modeling language, allowing constraints and objectives to be specified using standard MATLAB expression syntax [14].

We chose the following values as initial solution:

$$\begin{aligned} a_0 &= 0, & b_0 &= 0, \\ a_i &= \frac{\sin \frac{2i\pi}{n-1}}{2 \cos \frac{\pi}{2n-2}} = -a_{n-i}, & b_i &= \frac{1 - \cos \frac{2i\pi}{n-1}}{2 \cos \frac{\pi}{2n-2}} = b_{n-i} \quad \forall i = 1, \dots, n/2 - 1, \\ a_{n/2} &= 0, & b_{n/2} &= 1, \end{aligned}$$

which define the n -gon \mathbb{R}_{n-1}^+ , and the stopping criteria $\varepsilon = 10^{-5}$. Table 1 shows the optimal values A_n^* of the maximal area problem for even numbers $n = 6, 8, \dots, 84, 90, 100, 110, 120, 128$, along with the areas of the initial n -gons \mathbb{R}_{n-1}^+ , the best lower bounds \underline{A}_n found in the literature, and the upper bounds \overline{A}_n . We also report the number k of iterations in Algorithm 1 for each n . The results support the following keypoints:

1. For $6 \leq n \leq 12$, $\underline{A}_n - A_n^* \leq 10^{-8}$, i.e., Algorithm 1 converges to the best known optimal solutions found in the literature.
2. For $32 \leq n \leq 80$, $\underline{A}_n < A(\mathbb{R}_{n-1}^+) < A_n^*$, i.e., the solutions obtained by Pinter [10] are suboptimal.

Table 1: Maximal area problem

n	$A(\mathbb{R}_{n-1}^+)$	\underline{A}_n		\bar{A}_n	A_n^*	# ite. k
6	0.6722882584	0.6749814429	[2, 3, 9]	0.6961524227	0.6749814387	5
8	0.7253199909	0.7268684828	[4, 9]	0.7350842599	0.7268684802	10
10	0.7482573378	0.7491373459	[5, 9]	0.7531627703	0.7491373454	16
12	0.7601970055	0.7607298734	[5, 9]	0.7629992851	0.7607298710	24
14	0.7671877750	0.7675310111	[9]	0.7689359584	0.7675310093	33
16	0.7716285345	0.7718613220	[9]	0.7727913493	0.7718613187	43
18	0.7746235089	0.7747881651	[9]	0.7754356273	0.7747881619	55
20	0.7767382147	0.7768587560	[9]	0.7773275822	0.7768587517	68
22	0.7782865351	0.7783773308	[10]	0.7787276939	0.7783773228	81
24	0.7794540033	0.7795240461	[10]	0.7797927529	0.7795240330	95
26	0.7803559816	0.7804111201	[10]	0.7806217145	0.7804111058	109
28	0.7810672517	0.7811114192	[10]	0.7812795297	0.7811114002	122
30	0.7816380102	0.7816739255	[10]	0.7818102598	0.7816739044	136
32	0.7821029651	0.7818946320	[10]	0.7822446490	0.7821325276	148
34	0.7824867354	0.7823103007	[10]	0.7826046775	0.7825113660	159
36	0.7828071755	0.7826513767	[10]	0.7829063971	0.7828279054	169
38	0.7830774889	0.7829526627	[10]	0.7831617511	0.7830950955	177
40	0.7833076096	0.7832011589	[10]	0.7833797744	0.7833226804	183
42	0.7835051276	0.7834135187	[10]	0.7835674041	0.7835181187	185
44	0.7836759223	0.7835966860	[10]	0.7837300377	0.7836871900	184
46	0.7838246055	0.7837554636	[10]	0.7838719255	0.7838344336	179
48	0.7839548353	0.7838942710	[10]	0.7839964516	0.7839634510	172
50	0.7840695435	0.7840161496	[10]	0.7841063371	0.7840771278	162
52	0.7841711020	0.7841233641	[10]	0.7842037903	0.7841778072	150
54	0.7842614465	0.7842192995	[10]	0.7842906181	0.7842674010	138
56	0.7843421691	0.7843044654	[10]	0.7843683109	0.7843474779	128
58	0.7844145892	0.7843807534	[10]	0.7844381066	0.7844193386	118
60	0.7844798073	0.7844492943	[10]	0.7845010402	0.7844840717	109
62	0.7845387477	0.7845111362	[10]	0.7845579827	0.7845425886	101
64	0.7845921910	0.7834620877	[10]	0.7846096710	0.7845956631	94
66	0.7846408000	0.7845910589	[10]	0.7846567322	0.7846439473	88
68	0.7846851407	0.7846139029	[10]	0.7846997026	0.7846880001	82
70	0.7847256986	0.7846403575	[10]	0.7847390429	0.7847283036	77
72	0.7847628920	0.7847454020	[10]	0.7847751508	0.7847652718	72
74	0.7847970830	0.7845564840	[10]	0.7848083708	0.7847992622	68
76	0.7848285863	0.7847585719	[10]	0.7848390031	0.7848305850	64
78	0.7848576763	0.7845160579	[10]	0.7848673094	0.7848595143	61
80	0.7848845934	0.7848252941	[10]	0.7848935195	0.7848862871	58
82	0.7849095487	–		0.7849178354	0.7849111119	55
84	0.7849327284	–		0.7849404352	0.7849341725	52
90	0.7849931681	–		0.7849994298	0.7849943223	46
100	0.7850706272	–		0.7850751877	0.7850714422	38
110	0.7851278167	–		0.7851312404	0.7851284086	32
120	0.7851712379	–		0.7851738734	0.7851716781	28
128	0.7851988626	–		0.7852010332	0.7851992126	25

3. For all n , the solutions obtained with Algorithm 1 verify, within the limit of the numerical computations, Theorem 2 and Conjecture 1, i.e.,

$$\begin{aligned}
x_{n/2} &= 0, & y_{n/2} &= 1, \\
\|\mathbf{v}_{n/2-1}\| &= 1, & \|\mathbf{v}_{n/2+1}\| &= 1, \\
\|\mathbf{v}_{i+n/2} - \mathbf{v}_i\| &= 1, & \|\mathbf{v}_{i+n/2+1} - \mathbf{v}_i\| &= 1 & \forall i = 1, 2, \dots, n/2 - 2, \\
\|\mathbf{v}_{n-1} - \mathbf{v}_{n/2-1}\| &= 1, & & & \\
x_{n-i} &= -x_i, & y_{n-i} &= y_i & \forall i = 1, 2, \dots, n/2 - 1.
\end{aligned}$$

We illustrate the largest small 16-, 32- and 64-gons in Figure 5. Furthermore, we remark that Theorem 2 and Conjecture 1 are verified by each polygon of the sequence generated by Algorithm 1. All 6-gons generated by the algorithm are represented in Figure ?? and the coordinates of their vertices are given in Table 2.




Figure 5: Three largest small n -gons (U_n, A_n^*)

Table 2: Vertices of 6-gons generated by Algorithm 1

6-gon	Coordinates (x_i, y_i)					Area
	(x_1, y_1)	(x_2, y_2)	(x_3, y_3)	(x_4, y_4)	(x_5, y_5)	
R_5^+	(0.500000, 0.363271)	(0.309017, 0.951057)	(0.000000, 1.000000)	(-0.309017, 0.951057)	(-0.500000, 0.363271)	0.6722882584
P_6^1	(0.500000, 0.397460)	(0.339680, 0.940541)	(0.000000, 1.000000)	(-0.339680, 0.940541)	(-0.500000, 0.397460)	0.6749414624
P_6^2	(0.500000, 0.401764)	(0.343285, 0.939231)	(0.000000, 1.000000)	(-0.343285, 0.939231)	(-0.500000, 0.401764)	0.6749808685
P_6^3	(0.500000, 0.402283)	(0.343715, 0.939074)	(0.000000, 1.000000)	(-0.343715, 0.939074)	(-0.500000, 0.402283)	0.6749814310
P_6^4	(0.500000, 0.402345)	(0.343766, 0.939055)	(0.000000, 1.000000)	(-0.343766, 0.939055)	(-0.500000, 0.402345)	0.6749814386
P_6^5	(0.500000, 0.402352)	(0.343773, 0.939053)	(0.000000, 1.000000)	(-0.343773, 0.939053)	(-0.500000, 0.402352)	0.6749814387

5 Conclusion

We proposed a sequential convex optimization approach to find the largest small n -gon for a given even number $n \geq 6$, which is formulated as a nonconvex quadratically constrained quadratic optimization problem. The algorithm, also known as the concave-convex procedure, guarantees convergence to a locally optimal solution.

Without assuming Graham’s conjecture nor the existence of an axis of symmetry in our quadratic formulation, numerical experiments on polygons with up to $n = 128$ sides showed that each optimal n -gon obtained with the algorithm proposed verifies both conditions within the limitation of the numerical computations. Furthermore, for even $6 \leq n \leq 12$, the n -gons obtained correspond to the known largest small n -gons.

References

- [1] Karl Reinhardt. Extremale polygone gegebenen durchmessers. Jahresbericht der Deutschen Mathematiker-Vereinigung, 31:251–270, 1922.
- [2] H Bieri. Ungelöste probleme: Zweiter nachtrag zu nr. 12. Elem. Math, 16:105–106, 1961.
- [3] Ronald L Graham. The largest small hexagon. Journal of Combinatorial Theory, Series A, 18(2):165–170, 1975.
- [4] Charles Audet, Pierre Hansen, Frédéric Messine, and Junjie Xiong. The largest small octagon. Journal of Combinatorial Theory, Series A, 98(1):46–59, 2002.
- [5] Didier Henrion and Frédéric Messine. Finding largest small polygons with GloptiPoly. Journal of Global Optimization, 56(3):1017–1028, 2013.
- [6] Charles Audet. Maximal area of equilateral small polygons. The American Mathematical Monthly, 124(2):175–178, 2017.
- [7] Charles Audet, Pierre Hansen, and Dragutin Svrtan. Using symbolic calculations to determine largest small polygons. Journal of Global Optimization, pages 1–8, 2020.
- [8] Jim Foster and Tamas Szabo. Diameter graphs of polygons and the proof of a conjecture of Graham. Journal of Combinatorial Theory, Series A, 114(8):1515–1525, 2007.

-
- [9] Michael J Mossinghoff. Isodiametric problems for polygons. *Discrete & Computational Geometry*, 36(2):363–379, 2006.
 - [10] János D Pintér. Largest small n -polygons: Numerical results and conjectured optima. http://www.optimization-online.org/DB_FILE/2018/11/6950.pdf, 2018.
 - [11] Barry R Marks and Gordon P Wright. A general inner approximation algorithm for nonconvex mathematical programs. *Operations research*, 26(4):681–683, 1978.
 - [12] Gert R Lanckriet and Bharath K Sriperumbudur. On the convergence of the concave-convex procedure. In *Advances in neural information processing systems*, pages 1759–1767, 2009.
 - [13] Christian Bingane. OPTIGON: Extremal small polygons. <https://github.com/cbingane/optigon>, September 2020.
 - [14] CVX Research, Inc. CVX: MATLAB software for disciplined convex programming, version 2.2. <http://cvxr.com/cvx>, August 2012.
 - [15] Charles Audet, Pierre Hansen, and Frédéric Messine. Ranking small regular polygons by area and by perimeter. *Journal of Applied and Industrial Mathematics*, 3(1):21–27, 2009.
 - [16] Bao Yuan. The Largest Small Hexagon. Master’s thesis, National University of Singapore, 2004.