



Second Generation HIV Surveillance Within Montreal Communities from Sub-Saharan Africa and the English-Speaking Caribbean



Produced by the Surveillance and Protection sector
Montreal Public Health Branch
1301 Sherbrooke St. East
Montreal, QC H2L 1M3
514-528-2400
www.dsp.santemontreal.qc.ca

Authors

Aïssatou Fall
Alix Adrien
Pascale Leclerc
Joseph Cox

In this document, the masculine form is meant to be inclusive of both genders and is used merely to simplify the text.

Information contained in the report may be cited provided the source is acknowledged.

This paper is available online in the Publications section of the Public Health Branch website:
www.dsp.santemontreal.qc.ca

© Public Health Branch
Montreal Health and Social Services Agency (2014)
All rights reserved.

ISBN 978-2-550-77128-9 (Print Version)
ISBN 978-2-550-77128-9 (PDF)
Legal deposit – Bibliothèque et Archives nationales du Québec, 2014
Legal deposit – Library and Archives Canada, 2014

ACKNOWLEDGMENTS

The Second Generation HIV Surveillance Survey of Montreal Communities from Sub-Saharan Africa and the English-Speaking Caribbean (SGS-AFCAR Survey) was completed with the cooperation of many people and organizations. It was successfully executed through collaboration with research scientists, a community advisory committee, survey coordinators, interviewers, professionals from the Public Health Agency of Canada, and community organizations.

We would like to thank all stakeholders involved in this project:

Research Team

Alix Adrien, Principal Investigator, Public Health Branch of the Montreal Health and Social Services Agency (PHB of the Montreal Agency)
 Pascale Leclerc, Co-investigator, PHB of the Montreal Agency
 Joseph Cox, Co-investigator, PHB of the Montreal Agency
 Aïssatou Fall, Research Coordinator, PHB of the Montreal Agency
 Jérémie Butoyi, Field Coordinator, PHB of the Montreal Agency
 Susanna Ogunnaike-Cooke, Public Health Agency of Canada (PHAC)
 Jill Tarasuk, PHAC
 Erin Rutherford, PHAC
 Robert Remis, University of Toronto
 Raphael Bitéra, Institut national de santé publique du Québec (INSPQ)
 Marie Niyongere, sub-Saharan African community representative
 Janice Dayle, English-speaking Caribbean community representative
 Aurélie Hot, Coalition des organismes communautaires québécois de lutte contre le sida (COCQ-SIDA)

Community Advisory Committee

Alix Adrien, PHB of the Montreal Agency
 Aïssatou Fall, PHB of the Montreal Agency
 Jérémie Butoyi, PHB of the Montreal Agency
 Marie Niyongere, Centre for AIDS Services of Montreal–Women (CASM)
 Janice Dayle
 Joseph Jean-Gilles, GAP-VIES
 Adina Ungureanu, Alliance des communautés culturelles pour l'égalité dans la santé et services sociaux
 Geneviève Némouthé, COCQ-SIDA
 Aurélie Hot, COCQ-SIDA
 Cora-Lee Conway
 Michael Farkas
 Marcel Rachid
 Dorothy Williams, Black Community Resource Centre (BCRC)

Interviewers

Mamadou Bady Balde
Mamvula Dada Bakombo
Camille Kouessi Dagbo
Gorgui Diouf
Elhadji Mamadou Dioukhane
Samira Edwards
Irénikatché M. Hazoumé
Mawudjro Johnson-Zigbe
Samuels Keithus
Janice Dayle
Liliane Mefanche
Yannick André Tala Talom
Marion Tétégan

Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada (CCDIC, PHAC)

Chris Archibald
Susanna Ogunnaike-Cooke
Jill Tarasuk
Erin Rutherford
Stephen Cule
Tory Atwood

National Microbiology Laboratory, Public Health Agency of Canada

Laurie Malloch

Community organizations, churches, organizations of Quebecers of African and English-speaking Caribbean descent on the Island of Montreal**Funding**

Public Health Agency of Canada
Quebec Ministry of Health and Social Services

Report Writing Team

Aïssatou Fall, Data Analysis and Drafting Officer
Alix Adrien, Author
Pascale Leclerc, Author
Joseph Cox, Author

DIRECTOR'S MESSAGE

Monitoring health status and its determinants has been at the heart of public health practice in Quebec for decades. People from HIV-endemic countries are particularly vulnerable to sexually transmitted and blood-borne infections (STBBIs). However, little information is available to develop appropriate measures to combat these infections.

This surveillance project, conducted on the Island of Montreal between June 2013 and February 2014, is an anonymous and voluntary cross-sectional behavioural and biological survey. The main objectives of the Second Generation HIV Surveillance Survey of Montreal Communities from Sub-Saharan Africa and the English-Speaking Caribbean (SGS-AFCAR Survey) were to describe the prevalence of HIV and other STBBIs, and identify associated risk behaviours. This survey was the first to characterize the HIV epidemic in these communities. There was a high level of participation in the survey, which involved 1,602 participants.

The Public Health Branch of the Montreal Health and Social Services Agency is pleased to present this report, outlining the main findings of the SGS-AFCAR Survey. It provides important information for the development of policies, programs and interventions to prevent and control HIV infection and other STBBIs among Montrealers from sub-Saharan Africa and the English-speaking Caribbean.

I would like to thank all the people from sub-Saharan Africa and the English-speaking Caribbean living on the Island of Montreal who took the time to participate in this survey. I also want to thank the community organizations, churches, and organizations of Quebecers of African and English-speaking Caribbean descent for their support throughout this project. Community involvement makes it likely that these data will soon be used to implement preventive measures. Promotion, prevention and protection help us fulfill our mission to keep our world healthy.

Director, Public Health Branch



Richard Massé

ABSTRACT

Human immunodeficiency virus (HIV) infections continue to be a public health issue in Quebec. In 2013, the Montreal area alone recorded 61.5% of all new HIV diagnoses, and most cases occurred among people from HIV-endemic countries. According to the Public Health Agency of Canada (PHAC), people born in these countries continue to be overrepresented among those infected with HIV in Canada. Although there is a legal requirement to collect epidemiological information on HIV infection and AIDS in order to monitor public health, these data do not document risk behaviours in the most exposed populations. Therefore, “strengthened systems,” known as second generation surveillance systems, are needed to collect data linking behavioural data with biological data on HIV infection and other related infections (hepatitis C, syphilis, etc.).

This report outlines the findings of the Second Generation HIV Surveillance Survey of Montreal Communities from Sub-Saharan Africa and the English-Speaking Caribbean (SGS-AFCAR Survey) conducted on the Island of Montreal from June 2013 to February 2014. This was an anonymous and voluntary cross-sectional behavioural and biological survey conducted by the Public Health Branch of the Montreal Health and Social Services Agency (PHB of the Montreal Agency) in conjunction with the Public Health Agency of Canada (PHAC).

During the study period, 1,602 participants, including 1,105 Montrealers from sub-Saharan Africa and 497 English-speaking Caribbeans meeting the selection criteria, were recruited on the Island of Montreal. Recruitment sites were locations used by both communities for social interactions. These could be fixed or temporary locations at events or in specific locations. Just over half the participants (51.7%) in the two target communities were recruited using a vehicle to meet them in their neighbourhoods at grocery stores, ethnic product stores, hairdressing shops, shopping malls, subway stations, supermarkets, restaurants and parks.

Of the 1,105 Montrealers from sub-Saharan Africa who were recruited, 60.5% (668/1,105) were men, while 53.5% (266/497) of Montrealers from the English-speaking Caribbean were women.

HIV prevalence for all participants was 1.4% [95% CI: 0.9–2.1]; the sub-Saharan African women had the highest prevalence at 2.4% [95% CI: 1.2–4.2]. A total of 22 HIV-positive tests (18 among sub-Saharan Africans and 4 among English-speaking Caribbeans) were identified, and of those who tested positive for HIV, 68.2% (15/22) were unaware they were HIV positive. Age was significantly associated with HIV ($p<0.0001$), with most HIV-positive tests detected in 30–49 year-old [86.4% (19/22)].

Seroprevalence of hepatitis C virus (HCV) among all participants was 0.5% [95% CI: 0.2–1.01%], for a total of 8 positive HCV tests. In addition, one case of HIV and HCV coinfection was identified in a sub-Saharan African woman. Syphilis results will be available at a later date.

In total, 80.4% (866/1,077) of sub-Saharan Africans vs. 61.3% (302/493) of English-speaking Caribbeans had already undergone HIV testing, and just over half had undergone testing in the previous two years.

The frequency of high-risk sexual behaviour was significant in both target communities, including multiple partners (with regular or casual partners), and low condom use during high-risk sex, especially among women. The survey revealed a significant incidence of unprotected vaginal or anal intercourse among some participants during recent trips to HIV-endemic countries; the incidence was greater among women. These risk behaviours have a significant impact on the spread of the HIV epidemic, as people affected by these practices are at greater risk of acquiring or transmitting HIV.

Participants from sub-Saharan Africa were more aware of modes of HIV transmission than English-speaking Caribbeans. Certain misconceptions remain, such as the belief that a healthy-looking person cannot have HIV or that HIV is transmitted through mosquito bites. Correcting major misconceptions about HIV transmission is as important as providing accurate information on real modes of HIV transmission.

Moreover, 60.7% (963/1,587) of participants recruited in both communities did not have a family doctor. Among all participants who required medical care in the 12 previous months, 15.1% (223/1,473) reported having difficulty accessing these health care services during this period [i.e. 15.7% of Sub-Saharan Africans and 10.1% of English-speaking Caribbeans].

The surveillance survey revealed that some subgroups in both target communities differed in attitudes and sexual risk behaviours, as well as knowledge of modes of HIV transmission, and were therefore more vulnerable to HIV/AIDS: women, the 30–49 year age group, less educated people and socially vulnerable people. These subgroups should be priority targets for measures promoting HIV/AIDS prevention and protection.

In Quebec and elsewhere in Canada, the majority character of heterosexual transmission of HIV infection among people from sub-Saharan Africa and the Caribbean, as well as the preponderance of infection among women, are constants. In the SGS-AFCAR Survey, all HIV positive cases (n=22) occurred among heterosexuals, and just over half the cases [54.5% (12/22)] occurred in women.

These findings could be used as a benchmark for monitoring trends in HIV and HCV prevalence, as well as screening habits and behaviours associated with the acquisition and transmission of these infections among people from HIV-endemic countries, in general, but more specifically in the sub-Saharan African and English-speaking Caribbean communities. Moreover, the findings should help guide development of policies, programs and interventions to prevent and control HIV infection and other STBBIs by targeting the needs of ethno-cultural communities on the Island of Montreal.

TABLE OF CONTENTS

ACKNOWLEDGMENTS.....	i
DIRECTOR'S MESSAGE	iii
ABSTRACT.....	iv
TABLE OF CONTENTS.....	vii
LIST OF TABLES.....	ix
LIST OF ACRONYMS AND ABBREVIATIONS	xii
INTRODUCTION	1
1. BACKGROUND	2
1.1. Epidemiological background of HIV/AIDS.....	2
1.2. HIV Epidemiological Surveillance System	4
1.3. Objectives	6
1.4. SGS-AFCAR Survey feasibility study	6
2 METHODOLOGY.....	8
2.1 Target population and inclusion criteria.....	8
2.2 Sample size for the Island of Montreal.....	9
2.5. Recruitment strategies and sites	11
2.6. Data collection	11
2.7. Ethical considerations.....	14
2.8. Data analysis	15
3. Results	16
3.1. Summary of data collection results	16
3.2. Sociodemographic characteristics	19
3.3. Sexual behaviours	29
3.4. Travel history	40
3.5. History of testing for HIV or other STBBIs	42
3.6. HIV prevalence in the SGS-AFCAR Survey.....	48
3.7. Seroprevalence of HCV infection	51
3.8. Sources of information, knowledge and beliefs about HIV infection.....	52
3.9. Access to and use of health systems and services.....	57
3.10. HIV prevalence rate based on certain characteristics	60
4. STRENGTHS/LIMITATIONS	64
5. SUMMARY OF RESULTS, RISK PROFILE AND POTENTIAL INTERVENTIONS	67
5.1. Vulnerabilities common to both communities	67
5.2. Communities from sub-Saharan Africa.....	69
5.3. Communities from the English-speaking Caribbean.....	70
6. OTHER INTERVENTIONS AND RECOMMENDATIONS	72
6.1. Reality and needs of the communities involved.....	72
6.2. Commitment and awareness of health professionals and managers	73
6.3. Political responsibility	73

CONCLUSION	74
REFERENCES.....	75
APPENDICES.....	78

LIST OF TABLES

Table 1: Distribution of target population (15–49 years) on the Island of Montreal by geographical area of origin	9
Table 2: Breakdown of refusals and main reasons for refusal	16
Table 3: Data collection period, sample size and percentage of dried blood spot (DBS) samples provided	17
Table 4: Sample compared to 2006 Census data	17
Table 5: Distribution of different recruitment methods	18
Table 6: Breakdown of participants by sex and age group	19
Table 7: Distribution of participants by geographical area of birth	20
Table 8: Distribution of participants by immigration period and status	22
Table 9: Distribution of participants by sexual orientation and marital status	23
Table 10: Distribution of participants by the highest level of education	24
Table 11: Distribution of participants by employment status	24
Table 12: Distribution of participants by annual household income	25
Table 13: Distribution of participants by CCSS area of residence	26
Table 14: Distribution of participants by type of house	27
Table 15: Distribution of participants by number of lifetime sexual partners, men and women with vaginal or anal penetration	29
Table 16: Distribution of participants by age at first sexual intercourse with vaginal or anal penetration	30
Table 17: Number of lifetime sexual partners from sub-Saharan Africa, the Caribbean or Central/South Asia	30
Table 18: Sexual intercourse (vaginal or anal) in the past year	31
Table 19: Condom use (male or female) during the last sexual encounter regardless of the type of sexual partner	31
Table 20: Number of sexual partners with vaginal or anal penetration in the last 12 months	32
Table 21: Number of sexual partners with vaginal or anal penetration in the last 12 months	32
Table 22: Condom use with regular sexual partners in the last 12 months	33
Table 23: Number of casual sex partners with vaginal or anal penetration in the last 12 months	33
Table 24: Condom use with casual sex partners in the last 12 months	34
Table 25: Condom use during last sexual encounter with a casual sex partner	34
Table 26: HIV status of last regular or casual sex partner	35
Table 27: Sex in exchange for goods or services in the last 12 months	35
Table 28: Use of alcohol or drugs during vaginal or anal intercourse in the last 12 months	36
Table 29: Injection drug use history	37
Table 30: Vaginal drying in women during intercourse in the last 12 months	38
Table 31: Practice of excision in women	38
Table 32: Practice of circumcision in men	39
Table 33: Travel over the past five years in a country where HIV is highly endemic	40
Table 34: Vaginal or anal intercourse during a trip, with a partner living in a country where HIV is highly endemic	40
Table 35: Condom use during intercourse while travelling in a country where HIV is highly endemic	41
Table 36: HIV test history	42
Table 37: HIV test history during last pregnancy	43
Table 38: Reasons for not having had an HIV test in the last two years	43
Table 39: Self-reported HIV status	44

Table 40: Length of time since self-reported HIV positive status, circumstances and place of infection	44
Table 41: HIV monitoring and treatment	45
Table 42: Reported history of diagnosed STBBIs and tuberculosis	46
Table 43: HIV seroprevalence and known HIV status	48
Table 44: HCV seroprevalence, HIV and HCV coinfection	51
Table 45: Sources of information on HIV/AIDS	52
Table 46: Knowledge and beliefs about HIV	53
Table 47: Type of access to health care systems	57
Table 48: Lack of access to health care systems	58
Table 49: History of immunization against diseases and diagnostics	59
Table 50: HIV prevalence based on certain characteristics and behaviours	61

LIST OF FIGURES

Figure 1: Estimated number of new HIV infections by exposure category.....	2
Figure 2: HIV prevalence in sub-Saharan Africa and the Caribbean among adults (15–49 years).....	4
Figure 3: Sub-Saharan African and English-speaking Caribbean areas and countries	10
Figure 4: Cotton fibre filter paper for collecting organic fluids.....	12
Figure 5: Dried blood spot collection kit	12
Figure 6: Dried blood spot samples on drying rack.....	12
Figure 7: Distribution of recruitment sites in both target communities	18
Figure 8: Distribution of participants by area and country of origin.....	21
Figure 9: Distribution of participants by immigration period	22
Figure 10: Distribution of the number of HIV-positive cases by geographical area of birth.....	49
Figure 11: Percentage of HIV-positive test results by sex, age group and marital status.....	50
Figure 12: Percentage of knowledge and assumptions about HIV transmission in both communities.....	55
Figure 13: Percentage of knowledge and assumptions about HIV transmission in both communities by age group	55
Figure 14: Percentage of knowledge and assumptions regarding HIV transmission in both communities by highest level of education	56

LIST OF ACRONYMS AND ABBREVIATIONS

ACCÉSSS	Alliance des communautés culturelles pour l'égalité dans la santé et services sociaux
AIDS	Acquired immunodeficiency syndrome
BCRC	Black Community Resource Centre
CASM	Centre for AIDS Services of Montreal
CCDIC	Centre for Communicable Diseases and Infection Control
CLSC	Centre local de services communautaires – Local Community Service Centre
CO	Community Organization
COCQ-SIDA	Coalition des organismes communautaires québécois de lutte contre le sida
CSSS	Health and Social Services Centre
DBS	Dried blood spot
EC	Endemic country
EIA	Enzyme immunoassay
FI	Federal Initiative to Address HIV/AIDS in Canada
GAP-VIES	Groupe d'action pour la prévention de la transmission du VIH et l'éradication du SIDA
HCV	Hepatitis C virus
Het-endemic	Heterosexuals from HIV endemic areas
HIV	Human immunodeficiency virus
IDU	Injection drug user
INSPQ	Institut national de santé publique du Québec
LSPQ	Laboratoire de santé publique du Québec
MSM	Men who have sex with men
NHRL	National HIV and Retrovirology Laboratories
NML	National Microbiology Laboratory
Notifiable disease	Notifiable disease
PHA	People living with HIV/AIDS
PHAC	Public Health Agency of Canada
REB	Research Ethics Board
SGS-AFCAR	Second Generation Surveillance Within Montreal Communities from Sub-Saharan Africa and the English-Speaking Caribbean
SIH	Social inequalities in health
STBBI	Sexually transmitted and blood-borne infections
STI	Sexually transmitted infection
TP-PA	Treponema pallidum particle agglutination test
UNAIDS	Joint United Nations Programme on HIV/AIDS
WHO	World Health Organization

INTRODUCTION

According to the Public Health Agency of Canada (PHAC), people born in countries where the human immunodeficiency virus (HIV) is endemic continue to be overrepresented among those infected with HIV in Canada.^{1 2} According to the Joint United Nations Programme on HIV/AIDS (UNAIDS), these countries are located mainly in the Caribbean and sub-Saharan Africa, which is still severely affected by the epidemic, and accounted for 70% of all new HIV infections in 2013.³

In Canada, in 2011 the estimated rate of new infections among people from HIV-endemic countries was nine times higher than in other Canadians⁴, and heterosexual transmission was a major characteristic of HIV/AIDS in this population. Based on the various HIV exposure categories, men who have sex with men (MSM) was still the main risk category in Canada in 2012, followed by heterosexual contact and injection drug users (IDU).¹ There is an increasing percentage of new HIV infections included in the heterosexuals from HIV-endemic areas (Het-endemic) exposure category, i.e. 13.2% of Het-endemic persons of the 32.6% of reported cases involving heterosexual contacts.¹

In Quebec, in 2013 two-thirds of cases of HIV infections through heterosexual contact were from the Caribbean (essentially Haiti) or sub-Saharan Africa.⁵ Montreal area alone recorded 61.5% of all new HIV diagnoses, and most cases occurred among immigrants from HIV-endemic countries.⁵

Naturally, routine monitoring provides a set of information on persons who have recently been diagnosed with HIV⁵ and on AIDS cases.⁶ These data are used to analyze trends in people infected with HIV, but are not designed to document risk behaviours among populations most at risk. Therefore, additional data collection systems are necessary. The Second Generation HIV Surveillance Survey of Montreal Communities from Sub-Saharan Africa and the English-Speaking Caribbean (SGS-AFCAR Survey) is an anonymous and voluntary cross-sectional behavioural and biological survey implemented to fill this gap. It is part of the E-Track⁷ second generation surveillance system, a national public health surveillance system for monitoring HIV and related infections in populations from HIV-endemic countries, implemented in various sentinel sites, including Montreal (SGS-AFCAR Survey), Ottawa and Toronto in collaboration with PHAC.

The SGS-AFCAR Survey helps the Public Health Branch of the Montreal Health and Social Services Agency (PHB of the Montreal Agency) fulfill its mandate. The PHB's responsibility within the regional STBBI prevention program is to monitor health risk factors, and introduce effective intervention measures in order to improve strategies for preventing HIV/AIDS and protecting the most vulnerable populations.

This report discusses the findings of the SGS-AFCAR Survey conducted on the Island of Montreal from June 2013 to February 2014 within Montreal communities from sub-Saharan Africa and the English-speaking Caribbean. Having participated in two previous epidemiological studies,^{8 9 10} Montreal's Haitian community was not targeted in this study.

The report provides an overview of the epidemiological context of HIV/AIDS, describes the methodology and presents the main findings of the behavioural and biological surveillance system run by the PHB of the Montreal Agency in collaboration with PHAC. It concludes with a discussion and recommendations for the various stakeholders and partners.

1. BACKGROUND

1.1. Epidemiological background of HIV/AIDS

UNAIDS estimates that in 2013 the total number of people living with HIV/AIDS worldwide was 35.0 million (range: 33.2 to 37.2 million), an increase from 2001 (29.8 million, range: 28.1 to 31.9 million).³ There are significant geographic variations in HIV prevalence, incidence and modes of transmission. However, sub-Saharan Africa remains the most affected region. It is estimated that nearly 1.5 million people (range: 1.3 to 1.6 million) were infected in 2013. HIV prevalence among people aged 15 to 49 was 4.7% [4.4%–4.9%].³ In the Caribbean, an estimated 12,000 people (range: 9,400 to 14,000) were infected in 2013. HIV prevalence among people aged 15 to 49 was 1.1% [0.9%–1.2%].³

HIV is considered endemic in a country (see list in Appendix 1)² when HIV prevalence among adults (15 to 49 years) is greater than 1%, and when one of the following criteria are met:

- At least 50% of HIV cases are attributed to heterosexual transmission;
- The male/female ratio is 2:1 for prevalent infections; or
- HIV prevalence is greater than or equal to 2% among women receiving prenatal care.

HIV/AIDS remains a concern in **Canada**. Between 2008 and 2011, the estimated number of people with HIV/AIDS rose from 64,000 (range: 53,000 to 75,000) to 71,300 (range: 58,600 to 84,000), an 11.4% increase.⁴ The estimated prevalence rate in Canada in 2011 was 208.0 per 100,000 inhabitants (range: 171.0 to 245.1 per 100,000).¹¹ National estimates produced by PHAC suggest that an estimated 535 (range: 370 to 700) new infections in 2011 could be attributed to the heterosexuals from HIV-endemic areas exposure category. This subcategory is growing (see Figure 1) and accounted for 16.9% of new infections in Canada in 2011,^{4 11} whereas about 2.2% of Canada's population was born in an HIV-endemic country according to the 2006 Census.¹¹ When taking into account second-generation Canadians with at least one parent born in an HIV-endemic country, the proportion of the total population of people from these countries increases to 2.7%.²




Figure 1: Estimated number of incident HIV infections in Canada over time, by exposure category

Source: Public Health Agency of Canada⁴

In **Quebec**, an estimated 19,300 people (range: 15,600 to 23,000) were infected with HIV in 2011 (including those who developed AIDS).¹² Epidemiological data on HIV infection has been collected in Quebec to monitor public health since April 2002. Program data do not consider the prevalence or incidence of

infection in the province; they are based on cases confirmed by the Laboratoire de santé publique du Québec (LSPQ). From April 2002 to December 2013, 4,316 new diagnoses^{5 a} of HIV infection were recorded in Quebec, 63.8% (2,752/4,316) of which occurred in Montreal.⁵ The proportion of specimens tested in the public system whose results were confirmed positive by LSPQ was stable at less than 1%.⁵ In 2013, the 358 new diagnoses recorded in Quebec were grouped in the following major exposure categories: 63.1% were MSM; 15.9% were heterosexuals from HIV-endemic countries (Het-endemic); 12.0% were other heterosexual men and women from those countries; 4.7% were injection drug users (IDU); and 2.0% were in the “Other” category.⁵

The cases were concentrated in the **Montreal** area, which makes up 25% of Quebec’s population, but reported 61.5% (220/358) of new diagnoses made in 2013, 18.6% (41/220) of which were in the Het-endemic category.⁵ In Montreal, of the new diagnoses made in 2013, 65.7% (23/35) of women and 9.7% (18/185) of the men were from the Caribbean (mostly from Haiti) or from sub-Saharan Africa.⁵ In addition, 78.0% of the Het-endemic cases had never undergone HIV testing.⁵ A relatively high percentage of newly diagnosed cases were in late stages of the disease (less than 200 CD4 lymphocytes/ μ L).¹³ A significant percentage of cases of HIV infection were attributable to recently infected people in the primary infection period.^{13 14}

Significant progress has been made in reducing the number of new cases of HIV infection and AIDS-related deaths in **sub-Saharan Africa**. The number of new HIV infections in adults (15–49 years) decreased by 33% between 2005 and 2013.¹⁵ However, alarming trends persist in many areas.¹⁶ In 2013, sub-Saharan Africa accounted for 70% of all new HIV infections [1.5 million (range: 1.3–1.6 million)], 71% of people with HIV [24.7 million (range: 23.5 to 26.1 million); 58% of whom were women] and 72% of AIDS-related deaths [1.1 million (range: 1.0–1.3 million)].¹⁵ However, prevalence can vary considerably between sub-regions (West, Central, East and Southern) [see Figure 2],¹⁶ and countries, and sometimes within the same country.¹⁶ Southern and East Africa, which together represented 6% of the total population, accounted for 52% of all people living with HIV in 2012 [18.5 million (range: 7.5–19.5 million)] and almost half of newly infected people worldwide. In West Africa, prevalence rates were between 1.5% and 3%. Compared to the rest of Africa, West and Central African countries have always appeared less affected by the epidemic, but prevalence can reach relatively high levels in some of these countries.^{15 16}

The **Caribbean** remains the region with the highest prevalence outside Africa [1.1% (range 0.9–1.2%)].¹⁵ However, the epidemic has slowed down considerably since the mid-1990s and new HIV infections have dropped by 40% since 2005.¹⁵ In 2013, the total number of new HIV infections was 12,000 (range: 9,400–14,000), whereas 250,000 (range: 230,000–280,000) people were living with HIV (PHA) and there were 11,000 (range: 8,300–14,000) AIDS-related deaths.¹⁵ In total, 57% of new infections in the Caribbean occurred in Haiti [in 2012: 8,500 (range: 6,900–11,000) new infections; 150,000 (range: 130,000–160,000) PHA; 2.1% prevalence (range: 1.9–2.3%)] and 12% in Jamaica [in 2012: 1,400 (range: <1,000–2,000) new infections; 28,000 (range: 23,000–34,000) PHA; 1.7% prevalence (range: 1.4–2.0%)].^{15 16}

^a A “new diagnosis” is a case where no prior HIV testing was done or all previous tests were negative.




Figure 2: HIV prevalence in sub-Saharan Africa and the Caribbean among adults (15–49 years)

1.2. HIV Epidemiological Surveillance System

In Quebec, monitoring health status is a fundamental aspect of public health practice. The importance of continuously assessing public health makes this activity a core mandate of regional public health branches. Also, implementing public health activities involves regularly developing profiles of public health and related factors, monitoring temporal and spatial trends, and identifying priority problems.

AIDS and HIV infection epidemiology in Quebec can be described as:

- The number of reported AIDS cases: passive surveillance data come from the Quebec AIDS Surveillance Program;^{6 17}
- The incidence of AIDS-related mortality: data come from the Institut de la statistique du Québec;
- The number of cases of HIV infection: data are collected by the Quebec HIV infection surveillance program.⁵

In Quebec, HIV and AIDS are notifiable diseases only if the individual with HIV or AIDS has donated blood, organs or tissue or has received blood, blood products, organs or tissue.¹⁸ However, there is a legal requirement to collect epidemiological information on HIV infection and AIDS in order to monitor public health, under sections 10 to 14 of the *Minister's Regulation under the Public Health Act*.¹⁸ Furthermore, HIV and AIDS reports do not include names nor do they identify anyone. Public and private laboratories have been operating under this framework since April 2002. They perform HIV infection tests and send all specimens with positive, indeterminate or inconclusive results to the Laboratoire de santé publique du Québec (LSPQ) for final confirmation. Following positive HIV test results, a public health worker at LSPQ starts collecting the necessary epidemiological information. This involves only a telephone conversation between the public health worker and the doctor or health professional who requested the HIV test. The doctor who diagnosed a disease indicative of AIDS must collect the epidemiological information using form SP-100, "AIDS surveillance for ongoing monitoring of public health."^{18 19}

Data from the HIV infection / AIDS surveillance program is used to estimate the number of people living with HIV (PHA) who contact health services for HIV testing; identify the percentage of people screened at an advanced stage of the disease (late detection); and describe the distribution of cases among vulnerable populations.

Naturally, routine case surveillance provides a set of information on people recently diagnosed with HIV, and reflects the increased risk of HIV infection in certain subgroups. However, HIV case monitoring systems are not designed to provide information on risk behaviours among the most vulnerable subgroups, as well as certain high-risk populations with little access to health care, which means additional data collection systems are required.

According to UNAIDS and World Health Organization (WHO) (2001) recommendations,²⁰ second generation HIV surveillance systematically integrates information from multiple sources to provide a more complete picture of HIV epidemiology and trends in a given area or region. Also known as “strengthened systems,”²⁰ they supplement STBBI epidemiological data by linking behavioural data with biological data on HIV infection and other related infections (hepatitis C, syphilis, etc.). They help focus data collection on populations most at risk of being recently infected with HIV (populations with high levels of risk behaviour or young people who are becoming sexually active),²⁰ and provide a better understanding of epidemic trends and underlying behaviours in order to plan and improve HIV transmission prevention programs, early screening and case management while targeting the most vulnerable populations.

In Canada, the core principles of second generation HIV surveillance are established by developing, implementing and disseminating biological and behavioural surveillance data. To date, systems that have been designed and implemented include the I-Track system focused on injection drug users, the M-Track system, which focuses on men who have sex with men, and the P-Track system for PHA. A pilot survey for the A-Track surveillance system focusing on Aboriginal people was conducted in Regina, Saskatchewan, in the winter of 2011 and spring of 2012.⁷ In addition, PHAC is conducting second generation STBBI surveillance among various other populations at risk: Enhanced Surveillance of Canadian Street Youth (15 to 24 years) (E-SYS/SAJR) and women at risk, two key populations identified in the Federal Initiative to Address HIV/AIDS in Canada (FI).

E-Track is a second generation HIV surveillance system established by developing, implementing and disseminating data provided by biological and behavioural surveillance on populations from HIV-endemic countries. Various sentinel sites, including Montreal, Ottawa and Toronto, were selected as part of a national public health surveillance system designed to monitor HIV and related infections.⁷ The SGS-AFCAR Survey is the first E-Track survey in Canada. Each surveillance system is implemented in collaboration with a wide range of stakeholders, partners and participants, including provincial health departments, local health services, research scientists, national or provincial committees, and community associations.

Aside from two behavioural and biological studies involving Quebecers of Haitian descent in 1994–1996⁸ and then in 2007–2008^{9 10} by research scientists from the PHB of the Montreal Agency, GAP-VIES (Groupe d’action pour la prévention de la transmission du VIH et l’éradication du SIDA) and PHAC, no surveillance studies have been conducted within other ethnocultural communities in Quebec. The 1994–1996 study revealed 1.3% HIV prevalence among Quebecers of Haitian descent.⁸

In Montreal, the first phase of E-Track, the SGS-AFCAR Survey, focused on communities from sub-Saharan Africa and the English-speaking Caribbean. Methodologies used to implement the survey are described in Section 2.

1.3. Objectives

As a behavioural and biological surveillance system, the **overall objective** of the SGS-AFCAR Survey is to better understand trends in the HIV epidemic, related infections (hepatitis C and syphilis) and their determinants in Montreal communities from sub-Saharan Africa and the English-speaking Caribbean, to improve HIV transmission prevention and screening programs, and provide appropriate services to affected populations.

The **main objectives** of the SGS-AFCAR Survey in Montreal communities from sub-Saharan Africa and the English-speaking Caribbean are to describe:

- The prevalence of HIV and other sexually transmitted and blood-borne infections (STBBI);
- Sexual behaviours, drug use and other risk behaviours for HIV and other STBBIs;
- Behaviours in terms of testing for HIV and related STBBIs;
- Knowledge of HIV status and use of HIV treatment options;
- Basic knowledge of HIV transmission modes; and
- Access to and use of the health system and services.

Secondary objectives

- Describe the factors associated with HIV and other STBBIs; and
- Describe the factors associated with risk behaviours for HIV and other STBBIs.

This report provides the descriptive results of the Survey, with the results section focusing on the main objectives (see Section 3).

Prior to implementing the SGS-AFCAR Survey in Montreal, an exploratory phase was conducted to assess the feasibility and implementation of this type of second generation surveillance project. This study is described in the following section.

1.4. SGS-AFCAR Survey feasibility study

PHAC funded a feasibility study, conducted by the Coalition des organismes communautaires québécois de lutte contre le sida (COCQ-SIDA) in collaboration with the PHB of the Montreal Agency, to identify factors that could facilitate or hinder implementation of a second generation HIV surveillance survey among Montreal communities from sub-Saharan Africa and the English-speaking Caribbean.²¹

A community participatory approach helped foster community involvement in the decision-making stages of planning and implementing the SGS-AFCAR Survey in Montreal, through meetings, interviews and a community consultation workshop.

A community advisory committee with representatives from both target communities, community organizations, and (local, regional and national) public health stakeholders was created in 2010.

This committee comprised the following resource persons, representatives and organizations:

- Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada (CCDIC, PHAC);
- PHB of the Montreal Agency;
- Coalition des organismes communautaires québécois de lutte contre le sida (COCQ-SIDA);
- Institut national de santé publique du Québec (INSPQ);
- Groupe d'action pour la prévention de la transmission du VIH et l'éradication du SIDA (GAP-VIES);

- Centre for AIDS Services of Montreal (CASM);
- Alliance des communautés culturelles pour l'égalité dans la santé et services sociaux (ACCÉSSS);
- Round Table on Black History Month;
- Black Community Resource Centre (BCRC);
- Source de vie pour les nations (SVN);
- DESTA Lion Wolf School;
- Dr. Robert Remis, Professor of Epidemiology, University of Toronto Dalla Lana School of Public Health.

The committee's mandate was to advise the research team on:

- Items to be included in the base protocol;
- Key activities to be conducted (literature reviews, consultations, etc.) to explore whether the cultural communities found the proposed surveillance system acceptable;
- Preferred channels for community participation;
- Sampling strategies, questionnaire themes and obtaining preliminary community support;
- Mobilization of community organizations to promote the survey, identify recruitment sites, and the like;
- Criteria and process for selecting local survey teams (research scientists, contributors, research coordinator(s), interviewers and community advisory groups);
- Developing strategies to overcome data collection issues during the survey;
- Analysis of preliminary data results, key research questions to be answered during data analysis, data distribution strategies and messages for communities.

Based on the data collected (through individual interviews, Advisory Committee meetings and the community consultation workshop), the feasibility study found that the proposed second generation HIV surveillance within Montreal communities from sub-Saharan Africa and the English-speaking Caribbean was acceptable and feasible. It also identified meeting places and the distribution of the various communities concerned on the Island of Montreal, while noting potential challenges and obstacles to be overcome, as well the strengths supporting effective implementation of the SGS-AFCAR Survey.²¹

These strengths included the fact that the communities have a representative community and religious meeting places, as well the support of some influential community and religious leaders. However, the risk of stigmatization was a challenge to be overcome by conducting awareness and promotion activities in the target communities, while mobilizing and involving community leaders.²¹

Implementing the SGS-AFCAR Survey therefore promoted community involvement at all stages of the process (planning, development, assessment). It built capacity to reach the target ethno-cultural communities and to integrate the information acquired through consultations and collaboration with resource persons throughout the SGS-AFCAR Survey design and implementation process, as well as through the knowledge transfer and acquisition process.

2 METHODOLOGY

The SGS-AFCAR Survey is an anonymous and voluntary cross-sectional behavioural and biological survey. Participants were recruited mainly using methods based on “time-space sampling” (TSS) or “time-location sampling” (TLS) principles. Recruitment was performed based on the target population’s meeting locations, days and times. Participation was voluntary, the questionnaires were anonymous, and participants were required to provide informed consent.

The survey included a behavioural component and an optional biological component. The behavioural component documented sociodemographic characteristics; sexual behaviour; travel history; psychoactive substance use; access to and use of health systems and services; screening tests performed; care and treatment received for HIV; as well as knowledge and beliefs about HIV. Responses were collected using a questionnaire completed by the participant and/or by an interviewer.

The biological component involved a finger prick dried blood sample (DBS) collected by an interviewer to screen for the following infections: HIV, hepatitis C virus (HCV) and syphilis.

2.1 Target population and inclusion criteria

As mentioned earlier, in Quebec in 2013, two-thirds of cases of HIV infection through heterosexual contact were from the Caribbean (essentially Haiti) or sub-Saharan Africa. Having participated in two previous studies, the Haitian community was excluded from the SGS-AFCAR Survey. People from sub-Saharan Africa and the English-speaking Caribbean living on the Island of Montreal were the target population for this study.

Research funds were limited, which precluded conducting separate studies for these two populations. Funding constraints also prevented us from recruiting enough people from the English-speaking Caribbean community for a robust estimate of HIV prevalence, given the limited number of HIV cases expected in this community. We chose to conduct the study with these two groups because it was important to document sexual risk behaviours and other risk factors associated with HIV infection in this community. We also wanted to avoid the negative impact generated by the misunderstanding that populations included in the feasibility study, but excluded from the survey, were excluded because they were less exposed to HIV infection. The purpose of the study was not to compare the two populations to one another, but to describe each one.

The inclusion criteria were as follows:

- Reside on the Island of Montreal;
- Be between 15 and 49 years of age;
- Belong to the first generation, i.e. have been born in sub-Saharan Africa or the English-speaking Caribbean;
- Be able to speak and understand English or French;
- Be able to give informed consent.

2.2 Sample size for the Island of Montreal

The sample size of the SGS-AFCAR Survey was estimated taking into account the following:

- Estimated prevalence of the main behavioural factors of interest and HIV prevalence in sub-Saharan African (West, East, Central and Southern) and English-speaking Caribbean countries:
 - o Sample size needed to detect expected HIV prevalence with a “reasonable” degree of accuracy, based on known HIV prevalence in the country and geographical area of origin.
 - o Sample size needed to detect an expected 50% prevalence rate for the main behavioural factors of interest with a “reasonable” degree of accuracy.
- The target population aged 15 to 49 and identification of priority countries of origin based on 2006 Census data.
- Availability of financial resources.

Based on the above criteria, the sample size to be recruited was 1,100 people from sub-Saharan Africa and 500 people from the English-speaking Caribbean, for a total of 1,600 people to be recruited for the SGS-AFCAR Survey. (Table 1)

Based on the 2006 Census,²² this sample of 15–49 year-olds included 5% (1,100/24,300) of immigrants from sub-Saharan Africa, and 3% (500/16,040) of immigrants from the English-speaking Caribbean; these communities represented 4.3% (24,300/560,380) and 2.9% (16,040/560,380) of all first-generation immigrants residing on the Island of Montreal.

Table 1: Distribution of target population (15–49 years) on the Island of Montreal by geographical area of origin

	Target Population Based on the 2006 Census	Sample to Be Recruited for the SGS-AFCAR Survey	
	N	n	%
Central Africa	8,655	392	35.6
West Africa	7,645	346	31.5
East Africa	7,515	340	30.9
Southern Africa	485	22	2.0
Total Sub-Saharan Africa	24,300	1,100	
Total English-Speaking Caribbean	16,040	500	
Total population		1,600	

Due to the heterogeneity of the target population and practical difficulties involved in obtaining a representative sample of the two target communities, recruitment of participants focused on people from countries whose nationals formed a majority ($\geq 1,000$ people) on the Island of Montreal according to the 2006 Census. However, people from other countries in the target areas (whose nationals were a minority on the Island of Montreal) and who met the inclusion criteria were also eligible for the survey. In addition, to make it easier to generalize our results, we considered their distribution by sex, age group, health and social services centre (CSSS) and local community service centre (CLSC) area of residence on the Island of Montreal.

Demographic (2006 Census)²² and epidemiological data indicated that:

- 82.4% of people of African descent lived in the census metropolitan area of Montreal. Two-thirds (66.3%) resided in the administrative region of Montreal and the three districts of the City of Montreal where the population of African descent was most concentrated, namely Côte-des-Neiges/Notre-Dame-de-Grâce, Ahuntsic-Cartierville and Villeray-Saint-Michel-Parc-Extension. Most members of the African community aged 15 and over were first generation (82.9%).
- The vast majority of Caribbeans (93.3%) lived in the census metropolitan area of Montreal. Over two-thirds (68.9%) resided in the administrative region of Montreal and the three districts of the City of Montreal where the English-speaking Caribbean population was most concentrated, namely Côte-des-Neiges/Notre-Dame-de-Grâce, followed by the boroughs of LaSalle and Pierrefonds-Senneville. Over two-thirds (68.6%) of the members of the Caribbean community aged 15 and over were first generation.
- Finally, the majority of people from HIV-endemic countries lived in Montreal.

The map of all target sub-Saharan African and English-speaking Caribbean countries is presented in Figure 3.




Figure 3: Sub-Saharan African and English-speaking Caribbean areas and countries

2.5. Recruitment strategies and sites

Recruitment strategies were dependent on time, budget, restricted access to target communities, and prioritized suitable locations or activities for recruiting people from countries belonging to the two communities.

Different strategies were used to select recruitment sites in the target community, including:

- Reviewing newspapers, magazines, local online media or any other sources of relevant information;
- Networking, interviewing and maintaining regular contact with community leaders, representatives of community organizations, business owners and other resource persons;
- Consulting with members of both communities at every opportunity.

Recruitment sites were locations used by both communities for social interactions. These could be fixed or temporary locations at events or in specific locations.

The various recruitment sites used by the target population were:

- Fixed locations: community organizations (e.g., Black Community Resource Centre [BCRC] located in the Côte-des-Neiges Community resource centre [CRC])
- Mobile strategies (vehicle): grocery stores, ethnic product stores, hairdressing shops, shopping malls, subway stations, supermarkets, restaurants and parks
- Churches
- Associations/Groups: groups of nationals from the same countries
- Adult schools / universities
- Social events: cultural, sports, festivals

The recruitment period was from June 2013 to February 2014. SGS-AFCAR Survey promotion sessions were conducted before and during the recruitment period for the target population, leaders, community organizations and various groups of nationals from the same country. Also, information slips and business cards listing the inclusion criteria and a phone number for scheduling an appointment to participate in the study were made available in various popular locations and distributed to the target populations.

2.6. Data collection

2.6.1. Questionnaire


The E-Track survey questionnaire was created by a working group of experts from various regions of Canada (Ottawa, Montreal, Toronto), taking into account the questionnaire developed for surveys in Montreal for Quebecers of Haitian descent in 2007 and 2008.^{9 10} It contained questions regarding sociodemographic characteristics, sexual behaviour, drug use, screening practices for HIV, HCV and other STBBIs, knowledge and attitudes related to HIV infection and other STBBIs, and use of and access to health care and services.

Data for the SGS-AFCAR Survey were only collected using an electronic, English- or French-language questionnaire, which was either self-administered or completed by an interviewer or with his help. Each participant was assigned an anonymous serial identification code. This code was entered on the DBS sample collection form and the informed consent form for each participant, which was signed by the interviewer after he received the participant's verbal consent. To identify duplicates, a unique encrypted 12-digit alphanumeric identification code was generated (based on the participant's initials, date of birth and sex) for each participant using unidirectional software at the recruitment site.

2.6.2. Biological sample collection and analysis

Interviewers were required to obtain informed consent from participants for collecting finger prick dried blood spot samples (DBS) to screen for HIV, HCV and syphilis. The samples will be stored for future analysis, except where consent was explicitly refused at the time of the survey. Participants could also choose not to provide a DBS.

The biological sample was collected on a cotton fibre filter paper (Whatman 903® Protein Saver Card) designed to collect organic fluids (Figure 4). The fingertip to be punctured was cleaned with an alcohol swab and a micro-lancet was then used to collect a finger prick blood sample from the cleaned area (Figure 5). Drops of blood were collected on the sample card (Figure 6) and the puncture site was covered with a bandage.



Spec# = Serial code

Figure 4: Cotton fibre filter paper for collecting organic fluids



Figure 5: Dried blood spot collection kit



Figure 6: Dried blood spot samples on drying rack

Biological samples were labelled at the sampling point with an anonymous serial identification code matching the code on the participant's questionnaire. The coded samples were sent to the National HIV and Retrovirology Laboratories (NHRL) in Ottawa for HIV and HCV testing, and were subsequently sent to the National Microbiology Laboratory (NML) in Winnipeg for syphilis testing.

HIV testing of dried blood spot (DBS) specimens was performed using the Bio-Rad GS rLAV HIV-1 EIA assay. Confirmatory testing was subsequently performed using the Bio-Rad GS HIV-1 Western Blot assay. A positive result indicated a current HIV infection. Both the EIA and Western Blot assays used are approved by Health Canada as diagnostic assays for use with DBS specimens.

Hepatitis C testing of DBS specimens was performed using the Ortho® HCV version 3.0 EIA. Confirmatory testing was not performed for samples that tested positive. A positive result indicated past or present hepatitis C infection and did not discriminate acute from chronic or resolved infections.

Syphilis testing was performed using the Serodia® *Treponema pallidum* particle agglutination assay (TP-PA). As with HCV, confirmatory testing was not performed for samples that tested positive. A positive result indicated the presence of antibodies against a past or present syphilis infection. The syphilis testing results will be presented later in an addendum.

Because the surveillance system is anonymous, participants did not receive their test results. However, they were encouraged to undergo testing for HIV and other STBBIs at the various screening centres. Interviewers gave participants leaflets with contact information for health care providers and local screening centres.

2.6.3. Data management

Data entry and cleanup processes

CCDIC was responsible for managing all data, including lab results. Lab results were managed by NHRL, NML and CCDIC, and were sent to the PHB of the Montreal Agency as soon as they were available. Standardized methods were used to record all data.

Survey and training staff

The **principle investigator in Montreal** was responsible for overall direction and management of the study within the sentinel site, including finalizing the questionnaire, hiring staff for the local study team, supervising implementation of the survey, liaising with community stakeholders and PHAC, ensuring data security and quality, directing implementation of analysis plans, and writing a report.

A **coordinator** was appointed to oversee the SGS-AFCAR Survey's daily operations, and to plan and coordinate the **field team's (field coordinator and interviewers)** work. The coordinator provided the principal investigator, co-investigators and the PHAC team with regular progress reports on the survey.

Interviewers were responsible for recruiting all survey participants and obtaining their informed consent. They were also responsible for collecting DBS samples, answering participants' questions and providing contact information for related services. All interviewers (n=13) were from sub-Saharan Africa and the English-speaking Caribbean. Questionnaires were administered at the selected recruitment sites, and to the extent possible, a gender approach was adopted where necessary, i.e. interviewers worked in pairs (one woman / one man) and questioned participants of the same sex.

All team members received one week of training on the protocol and procedure prior to survey implementation. Training sessions were conducted at the PHB of the Montreal Agency. The following topics were covered during training: ethics, recruitment, sampling methods, questionnaire administration, universal precautions, DBS sampling techniques, and report content. Employee safety issues were also discussed with the Montreal study team.

The **E-Track National Coordinator**, a PHAC epidemiologist, helped the Montreal team implement the survey, train the local survey team, manage daily needs of the entire surveillance system, and answer any questions.

2.7. Ethical considerations

The surveillance project protocol, questionnaire, consent forms and information, as well as procedures were approved by Health Canada and PHAC's Research Ethics Board.

Quebec differs from some other provinces with respect to the need to have surveillance projects approved by a research ethics committee (we prefer the term "surveillance project" rather than "surveillance study"). Among other things, because "... it is the purpose of these projects (to monitor changes in specific public health issues and their determinants), apart from the potentially similar methods they employ, that differentiates these activities from those that require ethics review."²³ In light of these considerations, the Ethics Research Board of McGill University Health Centre (REB-MUHC) found that the SGS-AFCAR Survey did not meet all the criteria for review by a research ethics board.

This interpretation is consistent with statements made by the Quebec Ministry of Health and Social Services' Research Ethics Unit in the document entitled "Note de clarification relative aux compétences matérielles et territoriales des comités d'éthique de la recherche" [Clarification note on the material and territorial jurisdiction of research ethics committees] published in 2007.²⁴ The Ethics Unit document distinguishes between research and other activities parallel or closely related to research, and projects related to research requiring review by a research ethics committee, but not projects related to other activities.

However, some indicators measured as part of this project are included in the Plan régional de surveillance de Montreal [Montreal regional surveillance plan] (PRSM) 2011–2015. In its assessment of PRSM, the Public Health Ethics Committee (PHEC), emphasized the need to avoid stigmatizing the population targeted by the SGS-AFCAR Survey.²⁵ To reflect PHEC's concern about avoiding any negative impact on the target populations in connection with dissemination of project results, close collaboration was maintained with members of the sub-Saharan African and English-speaking Caribbean communities, as well as organizations working with them.

Participation in the survey was voluntary and anonymous. The questionnaires were administered and DBS collected only after informed consent was obtained. To protect the anonymity of participants, the interviewers signed the consent form.

Each participant had an anonymous serial identification code to link the questionnaire to their biological results. To avoid duplicates, an encrypted identification code (based on the participant's initials, date of birth and sex) was generated for each participant using unidirectional software at the recruitment site.

Participants received a fact sheet summarizing the information in the consent form and the contact details for the research ethics committees and the survey coordinator, and a small stipend (\$20) to cover time and travel expenses.

Participants did not receive their test results. Interviewers provided participants with counselling and contact information for health care providers and local screening centres, and encouraged them to undergo testing for HIV and other STBBIs at the various screening locations.

Interviewer training focused on the importance of maintaining data confidentiality, a point reinforced throughout the survey.

The surveillance project database is password-protected, and all documents will be kept under lock and key for a maximum period of five years.

2.8. Data analysis

Analyses performed in this report are mainly descriptive. They provide a simple, comprehensive overview of data collected for the SGS-AFCAR Survey. The refusal rate among eligible people contacted and the number of completed surveys (behavioural and/or biological component) were quantified.

Univariate descriptive analyses were performed to characterize participants recruited in Montreal communities from sub-Saharan Africa and the English-speaking Caribbean. Frequency tables were used for categorical variables, and central tendency and dispersion measurements were used for continuous variables. These analyses included: sociodemographic characteristics, sexual behaviour, travel history in HIV-endemic countries, history of testing for HIV and other STBIs, access to HIV-related treatment, knowledge of HIV transmission modes, and access to health care.

HIV and HCV seroprevalence was assessed for the study population, each target community and other subgroups. Syphilis seroprevalence results will be available later.

Bivariate analyses were performed to assess risk factors associated with HIV seroprevalence, but also to look for associations between sociodemographic characteristics and risk behaviours. The Chi-square test was used to compare percentages, and the Student's t-test to compare means.

Unadjusted odds ratios (OR) and 95% confidence intervals (CI) were calculated to measure the strength of associations.

Statistical significance was set at 5% for all analyses, based on alternative hypotheses. All analyses were performed using IBM® SPSS® Statistics 22 software for Windows.

2.8.1 Presentation of results

The data are presented in tables based on the two target communities (where numbers are sufficient for analysis).

The response categories are presented in order of decreasing frequency except for ordinal variables, and the "Other" category always comes last.

No statistical techniques were used to account for missing data in this report because most analyses involved only cross-tabulations. Missing data are shown in an independent row in each table (where possible). Denominators used to calculate percentages do not include questions where values are missing (unless otherwise indicated). To ensure data presented in tables are interpreted correctly, each table mentions the specific subgroup of respondents to which the analysis applies, which is referred to as the "Eligible" subgroup in the report. The "Total" row in each table shows the total size of the "eligible" sample.

Results presented in the report reflect the broad outlines of the SGS-AFCAR Survey questionnaire (Appendix 2).

3. RESULTS

3.1. Summary of data collection results

Between June 28, 2013, and February 8, 2014, 1,958 eligible persons were approached, and 1,602 actually participated in the SGS-AFCAR Survey, an 81.8% (1,602/1,958) participation rate.

A total of 341 eligible people refused to take part in the SGS-AFCAR Survey. The abandonment and duplication figure was 15.

Some recruitment sites (street, church, etc.) and the context in which the SGS-AFCAR Survey was being promoted made it impossible to establish the total number of ineligible people (nature and eligibility criteria) approached throughout the survey.

Sex, age group and main reasons for refusal were recorded for most people who refused to participate [76.7% (241/314)]. (Table 2)

Table 2: Breakdown of refusals and main reasons for refusal

(ELIGIBLE: All cases of refusal)

Refusals	n (%)
Communities	
Sub-Saharan Africa	167 (69.3)
English-Speaking Caribbean	74 (30.7)
(n)	241
Sex	
Men	124 (51.5)
Women	117 (48.5)
(n)	241
Age group	
<15 years	2 (0.8)
20–29 years	82 (34.0)
30–39 years	122 (50.6)
40–49 years	35 (14.5)
(n)	241
Reasons for refusal (not mutually exclusive)	
Did not want to provide a blood sample	120 (35.2)
Sub-Saharan Africa	73 (60.8)
Caribbean	47 (39.2)
Lack of time	79 (23.2)
Will make an appointment	51 (15.0)
Sceptical / Lack of trust	36 (10.6)
Did not want to answer questions	26 (7.6)
Afraid of being identified	13 (3.8)
Needs to ask for permission	4 (1.2)

The main reason for refusal was unwillingness to provide a dried blood spot (DBS) sample [35.2% (120/241)].

Those eligible were invited to participate in both parts of the survey, the behavioural questionnaire and dried blood spot (DBS) sample collection. (Table 3) All participants (N=1,602) completed an electronic questionnaire in French or English with an interviewer present. They either completed the questionnaire on their own [15.2% (244/1,602)] or with the help of the interviewer [32.3% (518/1,602)], or the questionnaire was administered entirely by the interviewer [52.4% (840/1,602)]. It took an average of 24.2 minutes to complete the questionnaire (n=1,577) (range: 10 to 98 min; median: 22 min).

Table 3: Data collection period, sample size and percentage of dried blood spot (DBS) samples provided

SGS-AFCAR Survey Implementation Period	Total Number of Participants n	Completed Questionnaires n (%)	DBS Provided n (%)
June 28, 2013 to February 8, 2014	1,602	1,602 (100.0)	1,546 (96.5%)

Of all the participants, 96.5% (1,546/1,602) provided a dried blood spot (DBS) sample for biological testing. (Table 3)

In total, 69.0% (1,105/1,602) of participants were from sub-Saharan Africa, and 31.0% (497/1,602) from the English-speaking Caribbean.

Table 4 provides a comparison of participants recruited using 2006 Census data, broken down by geographical area of birth, sex and age group.

Table 4: Sample compared to 2006 Census data

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan African		English-Speaking Caribbean	
	SGS-AFCAR Survey N=1105 % [95% CI]	2006 Census N=24,300 % [Difference]	SGS-AFCAR Survey N=497 % [95% CI]	2006 Census N=16,400 % [Difference]
Breakdown by region				
Central Africa	39.5 [37.2-42.4]	35.6 [+3.9]	—	—
West Africa	37.1 [34.8-40.0]	31.5 [+5.6]	—	—
East Africa	20.8 [18.9-23.2]	30.9 [-10.1]	—	—
Southern Africa	2.5 [1.8-3.5]	2.0 [+0.5]	—	—
Sex				
Percentage of men	60.5 [58.1-63.3]	52.9 [+7.6]	46.5 [42.9-50.9]	42.6 [+3.9]
Age group				
15–24 years	29.2 [27.0-31.9] →	16.8 [+12.4]	33.2 [29.8-37.3] →	15.3 [+17.9]
25–34 years	35.5 [33.2-38.3] →	28.1 [+7.4]	26.6 [23.4-30.4]	24.9 [+1.7]
35–44 years	27.2 [25.1-29.9]	30.8 [-3.6]	27.4 [24.2-31.3]	31.2 [-3.8]
45–49 years	8.1 [6.7-9.7]	—	12.9 [10.5-15.8]	—

Based on 2006 Census data, fewer people from East Africa were recruited. The distribution of participants from other African sub-regions did not differ greatly from Census data.

Because of certain recruitment strategies (driving around in a vehicle), slightly more men than women were recruited in both target communities. The same applied for the 15–24 and 25–34 age groups.

Note that according to the 2006 Census, the sub-Saharan African community (15–49 years) included slightly more men (52.9%) than women (47.1%), whereas the opposite was true in the English-speaking Caribbean community (57.4% women vs. 42.6% men).

Figure 7 shows the number and percentage of participants recruited using the different recruitment strategies.




Figure 7: Distribution of recruitment sites in both target communities

Just over half of the participants (51.7%) in the two target communities were recruited using a vehicle to meet them in their neighbourhoods at grocery stores, ethnic product stores, hairdressing shops, shopping malls, subway stations, supermarkets, restaurants and parks.

The distribution of different participant recruitment methods is described in Table 5.

Table 5: Distribution of different recruitment methods

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa N=1105 n (%)	English-Speaking Caribbean N=497 n (%)	Total N=1,602 n (%)
At recruitment site	610 (55.2)	231 (46.5)	841 (52.5)
Word of mouth	457 (41.4)	244 (49.1)	701 (43.8)
On recommendation	86 (7.8)	46 (9.3)	132 (8.2)
Community organizations	56 (5.1)	3 (0.6)	59 (3.7)
Other	11 (1.0)	2 (0.4)	13 (0.8)

Participants could specify several modes. The “Other” category included scheduling a telephone appointment and/or information obtained by reading flyers that had been distributed.

3.2. Sociodemographic characteristics

Of the 1,105 Montrealers from sub-Saharan Africa who were recruited, 60.5% (668/1,105) were men, while 53.5% (266/497) of Montrealers from the English-speaking Caribbean were women.

Age group

The average age of all participants (N=1,602) was 30.75 years (± 9.5) [range: 15 to 49 years; median = 31 years]. In people from sub-Saharan Africa, the average age was higher among men than women [31.09 vs. 29.49 years; $p=0.004$]. (Table 6)

The distribution of age groups was different when comparing the two target communities ($p<0.0001$), as was distribution by sex. One-third of participants were in the 15–24 age group [30.5% (488/1,602), i.e. 29.2% (323/1,105) of sub-Saharan Africans and 33.2% (165/497) of English-speaking Caribbeans]. The 25–29 and 30–34 age groups were larger in sub-Saharan Africans, while the 40–44 and 45–49 age groups were larger in English-speaking Caribbeans. (Table 6)

Table 6: Breakdown of participants by sex and age group

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Age						
Average [SD]	30.46 [± 9.11]	31.09 [± 9.15] **	29.49 [± 8.90]	31.39 [± 10.31]	30.91 [± 10.68]	31.80 [± 9.97]
Age group						
15–19	182 (16.5) ***	107 (16.0) *	75 (17.2)	89 (17.9)	46 (19.9) *	43 (16.2)
20–24	141 (12.8)	77 (11.5)	64 (14.6)	76 (15.3)	41 (17.7)	35 (13.2)
25–29	179 (16.2)	91 (13.6)	88 (20.1)	45 (9.1)	18 (7.8)	27 (10.2)
30–34	213 (19.3)	135 (20.2)	78 (17.8)	87 (17.5)	32 (13.9)	55 (20.7)
35–39	195 (17.6)	130 (19.5)	65 (14.9)	68 (13.7)	38 (16.5)	30 (11.3)
40–44	106 (9.6)	71 (10.6)	35 (8.0)	68 (13.7)	24 (10.4)	44 (16.5)
45–49	89 (8.1)	57 (8.5)	32 (7.3)	64 (12.9)	32 (13.9)	32 (12.0)
(n)	1105	668	437	497	231	266

In each table, the percentages of the “Total Sub-Saharan Africa” column are compared to those of the “Total English-speaking Caribbean” column. Also, in each community, the percentages of the “Men” column are compared with those of the “Women” column.

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Area and country of origin

The geographical areas of birth of sub-Saharan Africans are presented in Table 7.

Table 7: Distribution of participants by geographical area of birth

(ELIGIBLE: All respondents surveyed)

	Men n (%)	Women n (%)	Total n (%)
Sub-Saharan Africa			
Central Africa	268 (40.1)	169 (38.7)	437 (39.5)
West Africa	252 (37.7)	158 (36.2)	410 (37.1)
East Africa	130 (19.5)	100 (22.9)	230 (20.8)
Southern Africa	18 (2.7)	10 (2.3)	28 (2.5)
(n)	668	437	1105
English-Speaking Caribbean			
(n)	231	266	497

In all, 497 English-speaking Caribbean participants from 12 countries were recruited. Also, 1,105 people from 36 sub-Saharan African countries participated in the SGS-AFCAR Survey: 6 Central African countries [39.5% (437/1,105)]; 13 West African countries [37.1% (410/1,105)]; 10 East African countries [20.8% (230/1,105)] and 7 Southern African countries [2.5% (28/1,105)] (Figure 8).




Figure 8: Distribution of participants by area and country of origin

Immigration period and status

Sub-Saharan Africans had immigrated to Canada [range: 1969 to 2013; median=2009; mode=2013] more recently ($p < 0.0001$) than people from the English-speaking Caribbean [range: 1966 to 2013; median=2000; mode=2000], with 86.3% (951/1,102) vs. 50.2% respectively immigrating after 2000. (Table 8 and Figure 9)

Also, Caribbean women had immigrated more recently than men ($p = 0.034$). (Table 8)

Compared to people from sub-Saharan Africa, most English-speaking Caribbeans were Canadian citizens [91.7% vs. 85.0%; $p = 0.0001$]. (Table 8)

The percentage of people with refugee status was almost identical in both communities: [2.4% (26/1,105) of sub-Saharan Africans (19 men and 7 women) vs 2.0% (10/492) of English-speaking Caribbeans (7 men and 3 women)].


In addition, the majority of those recruited from both communities had spent most of their childhood (0 to 18 years) in their country of birth. This was truer for sub-Saharan Africans than English-speaking Caribbeans [86.5% (953/1,102) vs. 73.2% (364/497); $p < 0.0001$].

Table 8: Distribution of participants by immigration period and status

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Wome	Total	Men	Women
Immigration period						
<1990	19 (1.7) ***	13 (2.0)	6 (1.4)	114 (23.2)	65 (28.3) *	49 (18.7)
1990–1999	132 (12.0)	68 (10.2)	64 (14.7)	131 (26.6)	54 (23.5)	77 (29.4)
≥2000	951 (86.3)	585 (87.8)	366 (83.9)	247 (50.2)	111 (48.3)	136 (51.9)
(n)	1102	666	436	492	230	262
Missing	3 (0.3)	2 (0.3)	1 (0.2)	5 (1.0)	1 (0.4)	4 (1.5)
Total	1105	668	437	497	231	266
Immigration status						
Canadian citizen / Permanent resident	939 (85.0) ***	564 (84.4)	375 (85.8)	451 (91.7)	209 (90.9)	242 (92.4)
Work permit / Temporary or student visa	103 (9.3)	59 (8.8)	44 (10.1)	13 (2.6)	6 (2.6)	7 (2.7)
Refugee / Other	63 (5.7)	45 (6.7)	18 (4.1)	28 (5.7)	15 (6.4)	13 (5.0)
(n)	1105	668	437	492	230	262
Missing	0 (0.0)	0 (0.0)	0 (0.0)	5 (1.0)	1 (0.4)	4 (1.5)
Total	1105	668	437	497	231	266

* p<0.05 ** p<0.01 *** p<0.001

**Figure 9:** Distribution of participants by immigration period

Ethnicity and religion

In total, 97.9% (1,568/1,602) of participants from both communities identified themselves as black.

Most English-speaking Caribbeans were Protestant [(38.0% (189/497)), Christian [(19.9% (99/497)) or Catholic [(17.7% (88/497))] or did not belong to a religion [(11.5% (57/497))]. Sub-Saharan Africans were Catholic [(31.0% (343/1,105)), Protestant [(26.3% (291/1,105)), Muslim [(22.4% (248/1,105))] or Christian [(13.1% (145/1,105))]. Participants in both communities who called themselves Christian did not identify as Catholic.

Sexual orientation and marital status

Nearly all participants from both communities [96.8% (1,551/1,602)] reported they were heterosexual: 97.6% of sub-Saharan Africans vs. 95.0% of English-speaking Caribbeans. (Table 9) Other participants defined themselves as bisexual [1.9% (30/1,602)], homosexual [0.6% (9/1,602)] or of an “Other” orientation [0.7% (12/1,602)]. The distribution of these subgroups was different in the two target communities ($p=0.017$), and by sex among English-speaking Caribbeans ($p=0.049$). (Table 9)

With respect to marital status, 36.8% (1,551/1,101) of persons from sub-Saharan Africa and 10.9% (54/496) of English-speaking Caribbeans were married. The spouses of 6.4% (70/1,101) of sub-Saharan Africans and 0.6% (3/496) of English-speaking Caribbeans were living outside Canada.

The percentage of singles (unmarried), couples living together or couples living apart differed significantly in the two communities ($p=0.0001$). (Table 9)

Table 9: Distribution of participants by sexual orientation and marital status

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Sexual orientation						
Heterosexuals	1079 (97.6) *	651 (97.5)	428 (97.9)	472 (95.0)	225 (97.4) *	247 (92.8)
Bisexuals	13 (1.2)	8 (1.2)	5 (1.1)	17 (3.4)	3 (1.3)	14 (5.3)
Homosexuals	6 (0.5)	4 (0.6)	2 (0.5)	3 (0.6)	2 (0.9)	1 (0.4)
Other ^a	7 (0.6)	5 (0.7)	2 (0.5)	5 (1.0)	1 (0.4)	4 (1.5)
(n)	1105	668	437	497	231	266
Missing	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	1105	668	437	497	231	266
Marital status						
Single	564 (51.2) ***	351 (52.8)	213 (48.9)	301 (60.7)	136 (58.9)	165 (62.3)
Living together as a couple	370 (33.6)	211 (31.7)	159 (36.5)	105 (21.2)	50 (21.6)	55 (20.8)
Couple not living together	167 (15.2)	103 (15.5)	64 (14.7)	90 (18.1)	45 (19.5)	45 (17.0)
(n)	1101	665	436	496	231	265
Missing	4 (0.4)	3 (0.4)	1 (0.2)	1 (0.2)	0 (0.0)	1 (0.4)
Total	1105	668	437	497	231	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

^a “Other” sexual orientation: asexual, pansexual, questioning

Level of education

People from sub-Saharan Africa had a higher level of education than English-speaking Caribbeans. In fact, 70.1% (771/1,100) vs. 37.9% (188/496) had a high school education or higher ($p=0.0001$). This applied much more to African men than women [73.2% (485/663) of men vs. 65.4% (286/437) of women; $p=0.006$], and the opposite applied to English-speaking Caribbeans [29.9% (69/231) of men vs. 44.9% (119/265) of women; $p=0.001$]. (Table 10)

Table 10: Distribution of participants by the highest level of education

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Wome	Total	Men	Women
Elementary/primary education	329 (29.9) ***	178 (26.8) ***	151 (34.6)	308 (62.1)	162 (70.1) **	146 (55.1)
Secondary studies	301 (27.4)	163 (24.6)	138 (31.6)	160 (32.3)	62 (26.8)	98 (37.0)
College/university	279 (25.4)	179 (27.0)	100 (22.9)	23 (4.6)	7 (3.0)	16 (6.0)
Higher education	191 (17.4)	143 (21.6)	48 (11.0)	5 (1.0)	0 (0.0)	5 (1.9)
(n)	1100	663	437	496	231	265
Missing	5 (0.5)	5 (0.8)	0 (0.0)	1 (0.2)	0 (0.0)	1 (0.4)
Total	1105	668	437	497	231	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Employment status and household income

The distribution of employment status was significantly different ($p=0.0001$) in both communities and by sex among sub-Saharan Africans. At the time of the survey, there were more sub-Saharan African students: 43% vs. 29.3%; more unemployed English-speaking Caribbeans: 22.2% vs. 12.2%. (Table 11)

Table 11: Distribution of participants by employment status

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Wome	Total	Men	Women
Employed full time or part time	490 (44.8) ***	320 (48.3) *	170 (39.4)	239 (48.6)	122 (52.8)	117 (44.8)
Students	470 (43.0)	268 (40.4)	202 (46.9)	144 (29.3)	58 (25.1)	86 (33.0)
Unemployed	134 (12.2)	75 (11.3)	59 (13.7)	109 (22.2)	51 (22.1)	58 (22.2)
(n)	1094	663	431	492	231	261
Missing	11 (1.0)	5 (0.8)	6 (1.4)	5 (1.0)	0 (0.0)	5 (1.9)
Total	1105	668	437	497	231	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Of those who reported their family income, just over half [51.7% (710/1,374)] had an annual income $< \$15,000$, i.e. 53.1% (530/998) of those from sub-Saharan Africa and 47.9% (180/376) of those from the English-speaking Caribbean. However, the latter had higher incomes than sub-Saharan Africans ($p=0.001$). (Table 12)

In total, 41.3% (561/1,357) of participants reported that based on their household income over the past 12 months, they had found it “somewhat difficult / very difficult” to meet their housing and food needs. This was the case for 40.0% (394/987) of sub-Saharan Africans vs. 45.1% (167/370) of English-speaking Caribbeans ($p=0.084$, not significant).

Table 12: Distribution of participants by annual household income

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Less than \$5,000	279 (28.0) **	160 (26.2)	119 (30.7)	67 (17.8)	35 (20.2)	32 (15.8)
\$5,000 to \$14,999	251 (25.2)	157 (25.7)	94 (24.3)	113 (30.1)	53 (30.6)	60 (29.6)
\$15,000 to \$29,999	228 (22.8)	142 (23.2)	86 (22.2)	83 (22.1)	41 (23.7)	42 (20.7)
\$30,000 to \$49,999	143 (14.3)	98 (16.0)	45 (11.6)	79 (21.0)	34 (19.7)	45 (22.2)
\$50,000 to \$69,999	52 (5.2)	26 (4.3)	26 (6.7)	20 (5.3)	6 (3.5)	14 (6.9)
\$70,000 or more	45 (4.5)	28 (4.6)	17 (4.4)	14 (3.7)	4 (2.3)	10 (4.9)
(n)	998	611	387	376	173	203
Missing	1 (0.1)	0 (0.0)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Don't know	89 (8.1)	43 (6.4)	46 (10.5)	115 (23.1)	56 (24.2)	59 (22.2)
Refusal	17 (1.5)	13 (1.9)	4 (0.9)	6 (1.2)	2 (0.9)	4 (1.5)
Total	1105	668	437	497	231	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Housing, family environment

The main CSSS area in which the two communities resided was La Montagne [21.8% (332/1,520)], where 19.7% (211/1,072) of sub-Saharan Africans and 27.1% (121/447) English-speaking Caribbeans were recruited. The main CSSS areas where the English-speaking Caribbeans resided [e.g., 22.8% in Dorval-Lachine-Lasalle and 20.6% in Cavendish] were different ($p<0.0001$) from those where sub-Saharan Africans lived [e.g., 18.5% in Bordeaux-Cartierville-Saint-Laurent and 10.2% in Lucille-Teasdale]. (Table 13)

Table 13: Distribution of participants by CCSS area of residence

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
CSSS de la Montagne	211 (19.7) ***	124 (19.0)	87 (20.7)	121 (27.1)	48 (23.1)	73 (30.5)
CSSS de Dorval-Lachine-LaSalle	96 (8.9)	53 (8.1)	43 (10.2)	102 (22.8)	52 (25.0)	50 (20.9)
CSSS Cavendish	71 (6.6)	43 (6.6)	28 (6.7)	92 (20.6)	41 (19.7)	51 (21.3)
CSSS de Bordeaux-Cartierville-Saint-Laurent	199 (18.5)	128 (19.6)	71 (16.9)	15 (3.4)	5 (2.4)	10 (4.2)
CSSS du Sud-Ouest-Verdun	96 (8.9)	63 (9.6)	33 (7.9)	66 (14.8)	36 (17.3)	30 (12.6)
CSSS Lucille-Teasdale	109 (10.2)	67 (10.3)	42 (10.0)	4 (0.9)	2 (1.0)	2 (0.8)
CSSS de l'Ouest-de-l'Île	94 (8.8)	56 (8.6)	38 (9.0)	29 (6.5)	16 (7.7)	13 (5.4)
CSSS d'Achims et Montréal-Nord	62 (5.8)	40 (6.1)	22 (5.2)	2 (0.4)	1 (0.5)	1 (0.4)
CSSS de Saint-Léonard et Saint-Michel	54 (5.0)	31 (4.7)	23 (5.5)	4 (0.9)	3 (1.4)	1 (0.4)
CSSS Jeanne-Mance	31 (2.9)	21 (3.2)	10 (2.4)	9 (2.0)	2 (1.0)	7 (2.9)
CSSS de la Pointe-de-l'Île	30 (2.8)	15 (2.3)	15 (3.6)	3 (0.7)	2 (1.0)	1 (0.4)
CSSS du Cœur-de-l'Île	20 (1.9)	12 (1.8)	8 (1.9)	0 (0.0)	0 (0.0)	0 (0.0)
(n)	1073	653	420	447	208	239
Missing	32 (2.9)	15 (2.2)	17 (3.9)	50 (10.1)	23 (10.2)	27 (10.2)
Total	1105	668	437	497	231	266

* p<0.05 ** p<0.01 *** p<0.001

In both communities, 8 in 10 participants were renting a house, apartment or condo [80.6% (1,287/1,596)]. More sub-Saharan Africans [81.6% (898/1,101)] were tenants than English-speaking Caribbeans [78.6% (389/495)], who were more often owners (8.5% vs. 4.1%) than Sub-Saharan Africans (p=0.009). (Table 14)

A total of 87.8% (1,394/1,588) of participants from both communities had stable housing in the previous 12 months ($\leq$ 01 move). A greater percentage of English-speaking Caribbeans than sub-Saharan Africans (16.3% vs. 10.4%; p=0.001) reported that they had moved in the past year.

Table 14: Distribution of participants by type of house

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Type of dwelling						
Tenant	898 (81.6) **	546 (82.2)	352 (80.5)	389 (78.6)	172 (74.8)	217 (81.9)
Living with parents, family members or friends	149 (13.5)	89 (13.4)	60 (13.7)	60 (12.1)	36 (15.7)	24 (9.1)
Owner	45 (4.1)	22 (3.3)	23 (5.3)	42 (8.5)	21 (9.1)	21 (7.9)
At a hotel / motel / transition home	8 (0.7)	6 (0.9)	2 (0.5)	3 (0.6)	0 (0.0)	3 (1.1)
No fixed address	1 (0.1)	1 (0.2)	0 (0.0)	1 (0.2)	1 (0.4)	0 (0.0)
(n)	1101	664	437	495	230	265
Missing	4 (0.4)	4 (0.6)	0 (0.0)	2 (0.4)	1 (0.4)	1 (0.4)
Total	1105	668	437	497	231	266
Number of moves (last 12 months)						
≤1	983 (89.6) **	584 (88.5)	399 (91.3)	411 (83.7)	183 (80.3)	228 (86.7)
≥2	114 (10.4)	76 (11.5)	38 (8.7)	80 (16.3)	45 (19.7)	35 (13.3)
(n)	1097	660	437	491	228	263
Missing	8 (0.7)	8 (1.2)	0 (0.0)	6 (1.2)	3 (1.3)	3 (1.1)
Total	1105	668	437	497	231	266

* p<0.05 ** p<0.01 *** p<0.001

Summary of sociodemographic characteristics

A total of 1,602 participants, including 1,105 Montrealers from 36 sub-Saharan African countries, and 497 English-speaking Caribbeans from 12 countries meeting the selection criteria were recruited on the Island of Montreal from June 28, 2013 to February 8, 2014. A few more sub-Saharan African men were recruited (60.5%), and 53.5% of the English-speaking Caribbeans were women.

Almost all participants (98%) in both communities identified themselves as black. Although there are commonalities between black communities in general, there is also diversity in terms of ethnic, cultural, linguistic, religious² and sociodemographic characteristics within the sub-Saharan African communities themselves, and within the English- or French-speaking Caribbean communities. This is one reason for presenting the results for the two target communities separately. It also enabled the project to adapt intervention measures to the communities' specific characteristics.

The sociodemographic profile of participants in the SGS-AFCAR Survey is consistent with 2006 Census data, whose statistics on the Quebec population originating from these countries are similar in terms of age, level of education, employment status and income.²²

The average age for all participants was 30.7 years and the median age was 31.0 years. The distribution of age groups was different in the two target communities ($p<0.0001$), as was distribution by sex. One-third of the participants were in the 15–24 age group [30.5% (488/1,602)].

Most participants were heterosexual, 97.6% of sub-Saharan Africans vs. 95.0% of English-speaking Caribbeans. A significantly greater percentage of English-speaking Caribbeans than sub-Saharan Africans were single (60.7% vs. 51.2%) or couples not living together (18.1% vs 15.2%).

People from sub-Saharan Africa had a higher level of education than English-speaking Caribbeans. In fact, 70.1% vs. 37.9% had a high school education or higher ($p<0.0001$).

Of the participants who reported their family income, over half [51.7% (710/1,374)] had an annual income $< \$15,000$. English-speaking Caribbeans had higher incomes than sub-Saharan Africans ($p=0.001$). In addition, just over one-third of participants [41.3% (561/1,357)] reported they found it “somewhat difficult / very difficult” to meet their housing and food needs in the previous 12 months.

A greater percentage of sub-Saharan Africans had immigrated to Canada more recently (after 2000) than English-speaking Caribbeans (86.3% vs. 50.2%) ($p<0.0001$). Recent immigrant status often results in greater social and economic insecurity, which has to be taken into account when developing prevention programs.

3.3. Sexual behaviours

3.3.1. Overall sex life

Table 15 shows the number of lifetime sexual partners, men and women with vaginal or anal penetration, in each of the two target communities. Overall, 11.1% (167/1,498) of participants reported never having had sex in their lives [12.9% (133/1,031) of sub-Saharan Africans vs. 7.3% (34/467) of English-speaking Caribbeans].

English-speaking Caribbeans had more lifetime sexual partners ($p<0.0001$) than sub-Saharan Africans, and men had more lifetime sexual partners than women did in both communities ($p<0.0001$).

Although 13.1% (196/1,498) of all participants reported only one lifetime sexual partner, more than half reported having had four or more sexual partners [53.3% (798/1,498) or 47.4% (489/1,031) of sub-Saharan Africans vs. 66.2% (309/467) of English-speaking Caribbeans].

Table 15: Distribution of participants by number of lifetime sexual partners, men and women with vaginal or anal penetration

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
None	133 (12.9) ***	58 (9.4) ***	75 (18.0)	34 (7.3)	12 (5.5) ***	22 (8.8)
One	165 (16.0)	71 (11.6)	94 (22.5)	31 (6.6)	8 (3.7)	23 (9.2)
2 to 3	244 (23.7)	107 (17.4)	137 (32.9)	93 (19.9)	14 (6.5)	79 (31.6)
4 to 10	335 (32.5)	232 (37.8)	103 (24.7)	156 (33.4)	64 (29.5)	92 (36.8)
11	154 (14.9)	146 (23.8)	8 (1.9)	153 (32.8)	119 (54.8)	34 (13.6)
(n)	1031	614	417	467	217	250
Don't know	43 (3.9)	35 (5.2)	8 (1.8)	16 (3.2)	10 (4.3)	6 (2.3)
Refusal	31 (2.8)	19 (2.8)	12 (2.7)	14 (2.8)	4 (1.7)	10 (3.8)
Total	1105	668	437	497	231	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

The average age at first sexual intercourse with vaginal or anal penetration was higher among sub-Saharan Africans [18.4 years (range: 7 to 40 years)] than English-speaking Caribbeans [15.8 years (range: 6 to 40 years); $p < 0.0001$]. Men began having sexual relations earlier than women in both communities ($p < 0.0001$). (Table 16)

About 70% [69.9% (341/488)] of young women and men aged 15 to 24 had had sexual intercourse with vaginal or anal penetration, a third of them [30.5% (104/325)] before age 15.

Table 16: Distribution of participants by age at first sexual intercourse with vaginal or anal penetration

(ELIGIBLE: Respondents who had had a lifetime sexual relationship with vaginal or anal penetration)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Age at first sexual intercourse						
Average (SD)	18.4 (± 3.9) ***	18.0 (± 4.1) ***	19.1 (± 3.6)	15.8 (± 3.8)	14.4 (± 3.5) ***	17.0 (± 3.7)
Median	18	17	18	16	14	16
Range	7 - 40	7 - 40	9 - 35	6 - 40	6 - 40	8 - 36
(n)	918	572	346	444	208	236
Don't know	27 (2.8)	22 (3.6)	5 (1.4)	10 (2.2)	8 (3.7)	2 (0.8)
Refusal	27 (2.8)	16 (2.6)	11 (3.0)	9 (1.9)	3 (1.4)	6 (2.5)
Total	972	610	362	463	219	244

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$

More than half (57.7%) of sub-Saharan Africans and just under a third (32.2%) of English-speaking Caribbeans had had sexual partners from sub-Saharan Africa or the Caribbean. (Table 17)

Table 17: Number of lifetime sexual partners from sub-Saharan Africa, the Caribbean or Central/South Asia

(ELIGIBLE: Respondents who had had a lifetime sexual relationship with vaginal or anal penetration)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
None	60 (6.6) ***	41 (7.2) ***	19 (5.5)	43 (9.8)	22 (10.8) ***	21 (8.9)
Some partners	326 (35.7)	274 (48.3)	52 (15.0)	254 (58.0)	155 (76.4)	99 (42.1)
All partners	527 (57.7)	252 (44.4)	275 (79.5)	141 (32.2)	26 (12.8)	115 (48.9)
(n)	913	567	346	438	203	235
Missing	59 (6.1)	43 (7.0)	16 (4.4)	25 (5.4)	16 (7.3)	9 (3.7)
Total	972	610	362	463	219	244

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$

Most sexually active participants [84.5% (1,146/1,357)] reported having had vaginal or anal intercourse in the past year. The difference was significant between sub-Saharan African men and women (86.1% vs. 80.1%; $p=0.016$) and English-speaking Caribbean men and women (91.8% vs. 80.3%; $p=0.001$). However, 16.2% of sub-Saharan Africans and 14.3% of English-speaking Caribbeans reported not having had intercourse in the last 12 months. (Table 18)

Table 18: Sexual intercourse (vaginal or anal) in the past year

(ELIGIBLE: Respondents who had had a lifetime sexual relationship with vaginal or anal penetration)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Yes	767 (83.8)	490 (86.1) *	277 (80.1)	379 (85.7)	191 (91.8) **	188 (80.3)
No	148 (16.2)	79 (13.9)	69 (19.9)	63 (14.3)	17 (8.2)	46 (19.7)
(n)	915	569	346	442	208	234
Missing	57 (5.9)	41 (6.7)	16 (4.4)	21 (4.5)	11 (5.0)	10 (4.1)
Total	972	610	362	463	219	244

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Of the participants who had had sexual intercourse with vaginal or anal penetration in the past year, 55.3% (634/1,146) did not use a condom (male or female) during their last sexual encounter, regardless of the type of sexual partner. More women than men from sub-Saharan Africa were in this situation (64.6% vs. 52.9%; $p=0.002$). (Table 19)

Table 19: Condom use (male or female) during the last sexual encounter regardless of the type of sexual partner

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with a partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Yes	329 (42.9)	231 (47.1) **	98 (35.4)	183 (48.3)	96 (50.3)	87 (46.3)
No	438 (57.1)	259 (52.9)	179 (64.6)	196 (51.7)	95 (49.7)	101 (53.7)
(n)	767	490	277	379	191	188
Missing	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	767	490	277	379	191	188

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Just over one-third [36.7% (416/1,132)] of participants had multiple sexual partners (≥ 2 partners) in the past year. There was a significant difference between the two target communities. English-speaking Caribbeans had more sexual partners than Sub-Saharan Africans ($p < 0.0001$), and men had more sexual partners than women did in each community ($p < 0.0001$). (Table 20)

Table 20: Number of sexual partners with vaginal or anal penetration in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with a partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
One	516 (68.1) ***	275 (57.2) ***	241 (87.0)	200 (53.5)	65 (34.8) ***	135 (72.2)
2 to 3	164 (21.6)	133 (27.7)	31 (11.2)	100 (26.7)	63 (33.7)	37 (19.8)
4 to 10	71 (9.4)	66 (13.7)	5 (1.8)	55 (14.7)	41 (21.9)	14 (7.5)
≥ 11	7 (0.9)	7 (1.5)	0 (0.0)	19 (5.1)	18 (9.6)	1 (0.5)
(n)	758	481	277	374	187	187
Missing	9 (1.2)	9 (1.8)	0 (0.0)	5 (1.3)	4 (2.1)	1 (0.5)
Total	767	490	277	379	191	188

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$

3.3.2. Regular sexual partners

In the past year, 81.6% (620/760) of sub-Saharan Africans vs. 72.8% (273/375) of English-speaking Caribbeans had one regular sexual partner with vaginal or anal penetration, while 13.7% (104/760) of sub-Saharan Africans vs. 22.7% (85/375) of English-speaking Caribbeans had several ($p = 0.001$). In both target communities, men had more sexual partners than women ($p < 0.0001$). (Table 21)

Table 21: Number of sexual partners with vaginal or anal penetration in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with a partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
None	36 (4.7) **	32 (6.6) ***	4 (1.4)	17 (4.5)	9 (4.8) ***	8 (4.3)
One	620 (81.6)	357 (73.9)	263 (94.9)	273 (72.8)	115 (61.2)	158 (84.5)
Several	104 (13.7)	94 (19.5)	10 (3.6)	85 (22.7)	64 (34.0)	21 (11.2)
(n)	760	483	277	375	188	187
Missing	7 (0.9)	7 (1.4)	0 (0.0)	4 (1.1)	3 (1.6)	1 (0.5)
Total	767	490	277	379	191	188

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$

Nearly 80% (577/722) of sub-Saharan Africans and 78.5% (281/358) of English-speaking Caribbeans reported not always using condoms with their regular sexual partners (non-significant difference). (Table 22)

A higher percentage of sub-Saharan African women [86.0% (233/271)] than men [76.3% (344/451)] did not always use condoms with their regular sexual partners ($p=0.002$). More sub-Saharan African women in a relationship (whether or not cohabiting) were in this situation at 73.4% (171/233). (Data not shown)

Table 22: Condom use with regular sexual partners in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with at least one partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Always^a	145 (20.1)	107 (23.7) **	38 (14.0)	77 (21.5)	39 (21.8)	38 (21.2)
Not always^b	577 (79.9)	344 (76.3)	233 (86.0)	281 (78.5)	140 (78.2)	141 (78.8)
(n)	722	451	271	358	179	179
Missing	2 (0.3)	0 (0.0)	2 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)
Total	724	451	273	358	179	179

- Always: Condom was "Always" used

- Not always: Participants answered "Never," "Seldom," "Most of the time" or "Almost all the time."

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

3.3.3. Casual sex partners

Of those who had had intercourse with vaginal or anal penetration in the past year, more than a third had at least one casual sex partner, i.e. 32.5% (247/759) of sub-Saharan Africans vs. 46.0% (177/374) of English-speaking Caribbeans. Also, men in general had more casual sex partners and a greater proportion of English-speaking Caribbean men than women had more casual sex partners ($p<0.0001$). (Table 23)

Table 23: Number of casual sex partners with vaginal or anal penetration in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with a partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
None	512 (67.5) ***	278 (57.7) ***	234 (84.5)	202 (54.0)	69 (36.9) ***	133 (71.1)
One	125 (16.5)	91 (18.9)	34 (12.3)	71 (19.0)	41 (21.9)	30 (16.0)
Several	122 (16.1)	113 (23.4)	9 (3.2)	101 (27.0)	77 (41.2)	24 (12.8)
(n)	759	482	277	374	187	187
Missing	5 (0.7)	5 (1.0)	0 (0.0)	5 (1.3)	4 (2.1)	1 (0.5)
Total	764	487	277	379	191	188

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Just over half of English-speaking Caribbeans [51.4% (90/175)] vs. one-third [37.8% (95/251)] of sub-Saharan Africans did not always use a condom in the last 12 months during vaginal or anal intercourse with their casual sex partners ($p=0.005$). This applied to significantly more sub-Saharan African women than men (51.2% vs. 35.1%; $p=0.048$). (Table 24)

Table 24: Condom use with casual sex partners in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with at least one casual partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Always	156 (62.2) **	135 (64.9) *	21 (48.8)	85 (48.6)	63 (52.1)	22 (40.7)
Not always	95 (37.8)	73 (35.1)	22 (51.2)	90 (51.4)	58 (47.9)	32 (59.3)
(n)	251	208	43	175	121	54
Missing	1 (0.4)	1 (0.5)	0 (0.0)	2 (1.1)	1 (0.8)	1 (1.8)
Total	252	209	43	177	122	55

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Of the participants who had had casual sex partners in the last year, 40.0% (14/35) of sub-Saharan African women vs. 19.1% (40/209) of men ($p=0.006$), as well as 45.3% (24/53) of English-speaking Caribbean women vs. 23.1% (28/121) of men ($p=0.003$) did not use a condom during the last sexual encounter involving vaginal or anal penetration with a casual partner. (Table 25)

Table 25: Condom use during last sexual encounter with a casual sex partner

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with at least one casual partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Yes	190 (77.9)	169 (80.9) **	21 (60.0)	122 (70.1)	93 (76.9) **	29 (54.7)
No	54 (22.1)	40 (19.1)	14 (40.0)	52 (29.9)	28 (23.1)	24 (45.3)
(n)	244	209	35	174	121	53
Missing	8 (3.2)	0 (0.0)	8 (18.6)	3 (1.7)	1 (0.8)	2 (3.6)
Total	252	209	43	177	122	55

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

3.3.4. HIV status of last sexual partner

In total, 1.4% (13/938) of participants reported that their last regular or casual sex partner was HIV positive. (Table 26)

Table 26: HIV status of last regular or casual sex partner

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with a regular or casual sex partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Seronegative	622 (98.3)	387 (97.7)	235 (99.2)	303 (99.3)	149 (99.3)	154 (99.4)
Seropositive	11 (1.7)	9 (2.3)	2 (0.8)	2 (0.7)	1 (0.7)	1 (0.6)
(n)	633	396	237	305	150	155
Don't know	127 (16.6)	89 (18.3)	38 (13.7)	73 (19.3)	40 (20.9)	33 (17.6)
Missing	4 (0.5)	2 (0.4)	2 (0.7)	1 (0.3)	1 (0.5)	0 (0.0)
Total	764	487	277	379	191	188

* p<0.05 ** p<0.01 *** p<0.001

3.3.5. Sex in exchange for money, drugs or other goods and services

In total, 4.5% (52/1,145) of participants reported having given money, drugs or other goods or services in exchange for sex involving vaginal or anal penetration in the last year. In both target communities, more men had given money in exchange for sex than women, i.e. among sub-Saharan Africans, 6.3% of men vs. 0.4% of women (p<0.0001), and among English-speaking Caribbeans, 8.4% of men vs. 2.1% of women (p=0.007). (Table 27)

Also, 2.0% (23/1,142) of participants reported having received money, drugs or other goods or services in exchange for sex involving vaginal or anal penetration in the last year. A higher percentage of English-speaking Caribbeans were in this situation than sub-Saharan Africans, i.e. 4.2% vs. 0.9% (p<0.0001). More Caribbean men than women were in this situation (6.8% vs. 1.6%; p=0.012). (Table 27)

Table 27: Sex in exchange for goods or services in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Gave money, drugs or other goods in exchange for sex	32 (4.2)	31 (6.3) ***	1 (0.4)	20 (5.3)	16 (8.4) **	4 (2.1)
(n)	766	489	277	379	191	188
Missing	1 (0.1)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	767	490	277	379	191	188
Received money, drugs or other goods in exchange for sex	7 (0.9) ***	6 (1.2)	1 (0.4)	16 (4.2)	13 (6.8) *	3 (1.6)
(n)	763	487	276	379	191	188
Missing	4 (0.5)	3 (0.6)	1 (0.4)	0 (0.0)	0 (0.0)	0 (0.0)
Total	767	490	277	379	191	188

* p<0.05 ** p<0.01 *** p<0.001

3.3.6. Recreational alcohol or drug use

A third of participants had used alcohol while having sex in the last 12 months [30.3% (345/1,140), i.e. 19.7% (150/762) of sub-Saharan Africans vs. 51.6% (195/378) of English-speaking Caribbeans; $p<0.0001$]. (Table 28)

Higher percentages of English-speaking Caribbeans than sub-Saharan Africans and significantly more men than women in the two communities had used alcohol, marijuana or other drugs during vaginal or anal intercourse in the last 12 months. (Table 28)

Table 28: Use of alcohol or drugs during vaginal or anal intercourse in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with at least one partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Alcohol	150 (19.7) ***	123 (25.2) ***	27 (9.9)	195 (51.6)	120 (62.8) ***	75 (40.1)
(n)	762	488	274	378	191	187
Missing	5 (0.7)	2 (0.4)	3 (1.1)	1 (0.3)	0 (0.0)	1 (0.5)
Total	767	490	277	379	191	188
Marijuana/hashish	60 (7.9) ***	53 (10.9) ***	7 (2.6)	158 (41.8)	104 (54.5) ***	54 (28.9)
(n)	762	488	274	378	191	187
Missing	5 (0.7)	2 (0.4)	3 (1.1)	1 (0.3)	0 (0.0)	1 (0.5)
Total	767	490	277	379	191	188
Cocaine/crack/freebase	1 (0.1) ***	1 (0.2) ***	0 (0.0)	10 (2.7)	6 (3.2) **	4 (2.1)
(n)	762	488	274	377	190	187
Missing	5 (0.7)	2 (0.4)	3 (1.1)	2 (0.5)	1 (0.5)	1 (0.5)
Total	767	490	277	379	191	188
Other recreational drugs^a	3 (0.4) ***	2 (0.4)	1 (0.4)	30 (8.0)	16 (8.4)	14 (7.5)
(n)	762	488	274	377	190	187
Missing	5 (0.7)	2 (0.4)	3 (1.1)	2 (0.5)	1 (0.5)	1 (0.5)
Total	767	490	277	379	191	188

^a Popper; Viagra; Ecstasy; Crystal meth; GHB; Psychedelic drugs and other amphetamines.

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Four of the survey participants had an injection drug use history (i.e. three sub-Saharan Africans and one English-speaking Caribbean); only one of them had used in the last six months (a sub-Saharan African). (Table 29)

Table 29: Injection drug use history

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Injection of any type of drug						
Never	1098 (99.7)	662 (99.7)	436 (99.8)	494 (99.8)	229 (99.6)	265 (100.0)
Yes						
Over 6 months ago	2 (0.2)	1 (0.2)	1 (0.2)	1 (0.2)	1 (0.4)	0 (0.0)
Within the last 6 months	1 (0.1)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
(n)	1101	664	437	495	230	265
Missing	4 (0.4)	4 (0.6)	0 (0.0)	2 (0.4)	1 (0.4)	1 (0.4)
Total	1105	668	437	497	231	266

* p<0.05 ** p<0.01 *** p<0.001

3.3.7. Other sexual practices

Very few sexually active women practised vaginal drying in the last year, i.e. 2.6% (12/461). (Table 30)

Table 30: Vaginal drying in women during intercourse in the last 12 months

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with at least one partner in the last 12 months)

	Sub-Saharan Africa n=1105 n (%)	English-Speaking Caribbean n=497 n (%)
	Total	Total
Yes	5 (1.8)	7 (3.8)
No	270 (98.2)	179 (96.2)
(n)	275	186
Missing	2 (0.7)	2 (1.1)
Total	277	188

Overall, 6.3% (44/702) of African and Caribbean women were excised. The practice of excision was more common among sub-Saharan African women than English-speaking Caribbean women participants, with 9.2% vs. 1.5% ($p<0.0001$). (Table 31)

Table 31: Practice of excision in women

(ELIGIBLE: All women)

	Sub-Saharan Africa n=1105 n (%)	English-Speaking Caribbean n=497 n (%)
	Total	Total
Yes	40 (9.2) ***	4 (1.5)
No	397 (90.8)	261 (98.5)
(n)	437	265
Missing	0 (0.0)	1 (0.4)
Total	437	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

The vast majority of sub-Saharan African men were circumcised, i.e. 90.6%, compared to 35.4% of English-speaking Caribbeans ($p<0.0001$). (Table 32)

Table 32: Practice of circumcision in men

(ELIGIBLE: All men)

	Sub-Saharan Africa n=1105 n (%)	English-Speaking Caribbean n=497 n (%)
	Total	Total
Yes	599 (90.6) ***	81 (35.4)
No	62 (9.4)	148 (64.6)
(n)	661	229
Missing	7 (1.0)	2 (0.9)
Total	668	231

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

3.4. Travel history

3.4.1. Travel over the past five years in countries where HIV is highly endemic

Over the past five years, 36.5% (583/1,597) of participants had travelled to a country where HIV is highly endemic, i.e. 38.3% of sub-Saharan Africans vs. 32.5% of English-speaking Caribbeans ($p=0.024$). (Table 33)

Table 33: Travel over the past five years in a country where HIV is highly endemic

(ELIGIBLE: All respondents)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Yes	422 (38.3) *	256 (38.6)	166 (38.0)	161 (32.5)	69 (29.9)	92 (34.7)
No	679 (61.7)	408 (61.4)	271 (62.0)	335 (67.5)	162 (70.1)	173 (65.3)
(n)	1101	664	437	496	231	265
Missing	4 (3.2)	4 (0.0)	0 (0.0)	1 (0.2)	0 (0.0)	1 (0.4)
Total	1105	668	437	497	231	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Of the participants who had travelled in a country where HIV is highly endemic over the past five years, 33.7% (136/404) of sub-Saharan Africans vs. 27.2% (43/158) of English-speaking Caribbeans had engaged in sexual intercourse with a partner living in one of these countries. This applied more to sub-Saharan African ($p<0.0001$) and English-speaking Caribbean ($p=0.026$) men than women. (Table 34)

Table 34: Vaginal or anal intercourse during a trip, with a partner living in a country where HIV is highly endemic

(ELIGIBLE: All respondents who travelled in the past five years)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Yes	136 (33.7)	104 (42.8) ***	32 (19.9)	43 (27.2)	26 (38.2) *	17 (18.9)
No	233 (57.7)	129 (53.1)	104 (64.6)	104 (65.8)	38 (55.9)	66 (73.3)
Never had sexual intercourse	35 (8.7)	10 (4.1)	25 (15.5)	11 (7.0)	4 (5.9)	7 (7.8)
(n)	404	243	161	158	68	90
Missing	18 (4.3)	13 (5.1)	5 (3.0)	3 (1.9)	1 (1.4)	2 (2.2)
Total	422	256	166	161	69	92

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Slightly more than half [56.4% (101/179)] of participants who engaged in vaginal or anal intercourse during a trip in a country where HIV is highly endemic, with a sexual partner living there, did not always use a condom during sex. (Table 35)

A greater percentage of women from both communities were in this situation:

- 84.4% (27/32) of sub-Saharan African women vs. 52.9% (55/104) of men ($p=0.001$);
- 64.7% (11/17) of English-speaking Caribbean women vs. 30.8% (8/26) of men ($p=0.028$).

Table 35: Condom use during intercourse while travelling in a country where HIV is highly endemic

(ELIGIBLE: Respondents who reported having had vaginal or anal intercourse with at least one partner while travelling in an endemic country)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Always	54 (39.7)	49 (47.1) **	5 (15.6)	24 (55.8)	18 (69.2) *	6 (35.3)
Not always	82 (60.3)	55 (52.9)	27 (84.4)	19 (44.2)	8 (30.8)	11 (64.7)
(n)	136	104	32	43	26	17
Missing	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Total	136	104	32	43	26	17

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

3.5. History of testing for HIV or other STBBIs

3.5.1. HIV test history

A greater percentage of sub-Saharan Africans than English-speaking Caribbeans had undergone HIV testing, i.e. 80.4% (866/1,077) vs. 61.3% (302/493) ($p<0.0001$).

This applied to significantly more Caribbean women than men (68.9% vs. 52.4%; $p=0.0001$). (Table 36)

Also, just over half of those who had already had an HIV test had been tested in the past two years [67.8% of sub-Saharan Africans and 68.8% of English-speaking Caribbeans].

Table 36: HIV test history

(ELIGIBLE: All respondents)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Have already been tested for HIV						
Yes	866 (80.4) ***	513 (78.7)	353 (83.1)	302 (61.3)	120 (52.4) ***	182 (68.9)
No	211 (19.6)	139 (21.3)	72 (16.9)	191 (38.7)	109 (47.6)	82 (31.1)
(n)	1077	652	425	493	229	264
Missing	28 (2.5)	16 (2.4)	12 (2.7)	4 (0.8)	2 (0.9)	2 (0.8)
Total	1105	668	437	497	231	266
Have been tested for HIV in the last 2 years						
(ELIGIBLE: Respondents who have been tested)						
Yes	579 (67.8)	353 (66.3)	244 (69.9)	205 (68.8)	81 (68.6)	124 (68.9)
No	275 (32.2)	170 (33.7)	105 (30.1)	93 (31.2)	37 (31.4)	56 (31.1)
(n)	854	505	349	298	118	180
Missing	12 (1.4)	8 (1.6)	4 (1.1)	4 (1.3)	2 (1.7)	2 (1.1)
Total	866	513	353	302	120	182

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

At the time of the survey, 91.4% (202/221) of sub-Saharan African women participants had been tested for HIV during their last pregnancy compared to 83.5% (111/133) of English-speaking Caribbean women ($p=0.024$). Overall, 70.9% (222/313) of all women tested during their last pregnancy were tested in Canada. (Table 37)

Table 37: HIV test history during last pregnancy

(ELIGIBLE: Women)

	Sub-Saharan Africa n=1105 n (%)	English-Speaking Caribbean n=497 n (%)
	Total	Total
HIV testing during last pregnancy		
Yes	202 (91.4) *	111 (83.5)
No	19 (8.6)	22 (16.5)
(n)	221	133
Never been pregnant	98 (22.4)	45 (16.9)
No sexual intercourse	75 (17.2)	22 (8.3)
Missing	43 (9.8)	66 (24.8)
Total	437	266

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Table 38: Reasons for not having had an HIV test in the last two years

(ELIGIBLE: Respondents who reported never having had an HIV test)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Total	Men	Total	Women
You never thought about it	111 (40.4)	66 (38.8)	45 (42.9)	31 (33.3)	10 (27.0)	21 (37.5)
You're healthy, so you think you don't need to take the test	96 (34.9)	69 (40.6)	27 (25.7)	15 (16.1)	9 (24.3)	6 (10.7)
You think your risk of HIV infection is low	72 (26.2)	42 (24.7)	30 (28.6)	26 (28.0)	11 (29.7)	15 (26.8)
You think you know your HIV status	54 (19.6)	33 (19.4)	21 (20.0)	19 (20.4)	12 (32.4)	7 (12.5)
You think you can't get HIV	34 (12.4)	23 (13.5)	11 (10.5)	3 (3.2)	1 (2.7)	2 (3.2)
You're afraid to get tested	18 (6.5)	11 (6.5)	7 (6.7)	1 (1.1)	0 (0.0)	1 (1.8)
You think being HIV positive would affect your personal or professional life	6 (2.2)	4 (2.4)	2 (1.9)	2 (2.2)	0 (0.0)	2 (3.6)
You were tested during your immigration medical examination in Canada	13 (4.7)	7 (4.1)	6 (5.7)	13 (14.0)	3 (8.1)	10 (17.9)
You think nothing can be done if the test indicates you have HIV	3 (1.1)	2 (1.2)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)
You had trouble getting tested	2 (0.7)	2 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
You think the test is not reliable	4 (1.5)	3 (1.8)	1 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)
(n)	275	170	105	93	37	56

3.5.2. Self-reported HIV status and monitoring

Most participants who had already had an HIV test and knew their results reported that the results of their most recent HIV test were negative [97.9% (1,132/1,156)]. (Table 39)

However, **seven sub-Saharan Africans** (two men and five women) **reported that the results of their most recent HIV test were positive**. There were no self-reported HIV cases among English-speaking Caribbeans.

Table 39: Self-reported HIV status

(ELIGIBLE: Respondents who have already had an HIV test)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Seropositive	7 (0.8)	2 (0.4)	5 (1.4)	0 (0.0)	0 (0.0)	0 (0.0)
Seronegative	838 (97.9)	496 (98.0)	342 (97.7)	294 (98.0)	117 (98.3)	177 (97.8)
Unknown status ^a	11 (1.3)	8 (1.6)	3 (0.9)	6 (2.0)	2 (1.7)	4 (2.2)
(n)	856	506	350	300	119	181
Missing	10 (1.2)	7 (1.4)	3 (0.8)	2 (0.7)	1 (0.8)	1 (0.5)
Total	866	513	353	302	120	182

^a Includes respondents who reported they did not know the result of their last test, and respondents who had not received the results of their most recent test.

Six of the seven sub-Saharan Africans who reported they were HIV positive had known their status for over four years, and the circumstances of infection were specified for four of these seven individuals. Six of the seven individuals were infected outside Canada. (Table 40)

Table 40: Length of time since self-reported HIV positive status, circumstances and place of infection

(ELIGIBLE: Self-reported HIV-positive respondents)

	Sub-Saharan Africa n=1105 n (%)		
	Total	Men	Women
Length of time since self-reported HIV positive status			
<1 year	0 (0.0)	0 (0.0)	0 (0.0)
≥1 year but <4 years	1 (14.3)	0 (0.0)	1 (20.0)
≥ 4 years	6 (85.7)	2 (100.0)	4 (80.0)
(n)	7	2	5
Circumstances of infection			
Sexual intercourse with an HIV-positive man	2 (28.6)	0 (0.0)	2 (40.0)
Sexual intercourse with an HIV-positive woman	2 (28.6)	2 (100.0)	0 (0.0)
Not specified	3 (42.9)	0 (0.0)	3 (60.0)
(n)	7	2	5
Place of infection			
In Canada	1 (14.3)	0 (0.0)	1 (20.0)
In another country	6 (85.7)	2 (100.0)	4 (80.0)
(n)	7	2	5

Only one of the seven self-reported HIV-positive sub-Saharan Africans had stopped treatment for medical reasons. The other six were receiving antiretroviral therapy. (Table 40) Two out of five women had tried alternative or conventional treatments for HIV infection. (Table 41)

In addition, two self-reported HIV positive participants [one male and one female, 28.6% (2/7)] had regular HIV serodiscordant or seroconcordant sexual partners with whom they did not always use condoms. (Results not shown)

Table 41: HIV monitoring and treatment

(ELIGIBLE: Self-reported HIV-positive respondents)

	Sub-Saharan Africa n=1105 n (%)		
	Total	Men	Women
Time between confirmation of diagnosis and medical consultation			
Within 3 months of diagnosis	3 (42.9)	0 (0.0)	3 (60.0)
Between 4 and 6 months after diagnosis	1 (14.3)	0 (0.0)	1 (20.0)
Between 7 and 12 months after diagnosis	1 (14.3)	1 (50.0)	0 (0.0)
More than a year after diagnosis	2 (28.6)	1 (50.0)	1 (20.0)
(n)	7	2	5
Currently receiving treatment for HIV			
Yes	7 (100.0)	2 (100.0)	5 (100.0)
No	0 (0.0)	0 (0.0)	0 (0.0)
(n)	7	2	5
Has already taken HIV medications			
Yes	6 (85.7)	2 (100.0)	4 (80.0)
No	1 (14.3)	0 (0.0)	1 (20.0)
(n)	7	2	5
Currently taking HIV medications			
Yes	6 (85.7)	2 (100.0)	4 (80.0)
No	1 (14.3)	0 (0.0)	1 (20.0)
(n)	7	2	5
Initial treatment location			
In Canada	5 (71.4)	1 (50.0)	4 (80.0)
In another country	1 (14.3)	1 (50.0)	0 (0.0)
(n)	6	2	4
Missing	1	0	1 (14.3)
Total	7	2	5

3.5.3. History of other reported STBBIs and tuberculosis

Table 38 provides a detailed STBBI history. The percentages of participants who reported having had various STBBIs depended on the type of infection. For all participants, 11.9% (59/497) of English-speaking Caribbeans vs. 3.4% (38/1,105) of sub-Saharan Africans had reported a history of chlamydia, gonorrhea or syphilis ($p < 0.0001$). This was the case in 5.3% (26/488) of those aged 15–24. (Table 42)

Table 42: Reported history of diagnosed STBBIs and tuberculosis

(ELIGIBLE: All respondents)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Hepatitis C						
No	1078 (98.7)	649 (98.8)	429 (98.6)	485 (98.8)	223 (98.2)	262 (99.2)
Yes, in the last six months	7 (0.6)	2 (0.3)	5 (1.1)	1 (0.2)	0 (0.0)	1 (0.4)
Yes, over six months ago	7 (0.6)	6 (0.9)	1 (0.2)	5 (1.0)	4 (1.8)	1 (0.4)
(n)	1092	657	435	491	227	264
Missing	13 (1.2)	11 (1.6)	2 (0.5)	6 (1.2)	4 (1.7)	2 (0.8)
Total	1105	668	437	497	231	266
Hepatitis B						
No	1067 (97.4)	642 (97.3)	425 (97.7)	486 (99.0)	224 (98.7)	262 (99.2)
Yes, in the last six months	7 (0.6)	3 (0.5)	4 (0.9)	2 (0.4)	0 (0.0)	2 (0.8)
Yes, over six months ago	21 (1.9)	15 (2.3)	6 (1.4)	3 (0.6)	3 (1.3)	0 (0.0)
(n)	1095	660	435	491	227	264
Missing	10 (0.9)	8 (1.2)	2 (0.5)	6 (1.2)	4 (1.7)	2 (0.8)
Total	1105	668	437	497	231	266
Syphilis						
No	1084 (98.8)	652 (98.6)	432 (99.1)	486 (98.8)	225 (98.7)	261 (98.9)
Yes, in the last six months	5 (0.5)	3 (0.5)	2 (0.5)	1 (0.2)	0 (0.0)	1 (0.4)
Yes, over six months ago	8 (0.7)	6 (0.9)	2 (0.5)	5 (1.0)	3 (1.3)	2 (0.8)
(n)	1097	661	436	492	228	264
Missing	8 (0.7)	7 (1.0)	1 (0.2)	5 (1.0)	3 (1.3)	2 (0.8)
Total	1105	668	437	497	231	266
Gonorrhea						
No	1077 (98.3) ***	648 (98.2)	429 (98.4)	467 (95.1)	215 (94.3)	252 (95.8)
Yes, in the last six months	4 (0.4)	2 (0.3)	2 (0.5)	1 (0.2)	0 (0.0)	1 (0.4)
Yes, over six months ago	15 (1.4)	10 (1.5)	5 (1.1)	23 (4.7)	13 (5.7)	10 (3.8)
(n)	1096	660	436	491	228	263
Missing	9 (0.8)	8 (1.2)	1 (0.2)	6 (1.2)	3 (1.3)	3 (1.1)
Total	1105	668	437	497	231	266

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Genital or anal warts						
No	1088 (99.2) *	657 (99.4)	431 (98.9)	484 (98.4)	222 (97.4)	262 (99.2)
Yes, in the last six months	5 (0.5)	1 (0.2)	4 (0.9)	1 (0.2)	0 (0.0)	1 (0.4)
Yes, over six months ago	4 (0.4)	3 (0.5)	1 (0.2)	7 (1.4)	6 (2.6)	1 (0.4)
(n)	1097	661	436	492	228	264
Missing	8 (0.7)	7 (1.0)	1 (0.2)	5 (1.0)	3 (1.3)	2 (0.8)
Total	1105	668	437	497	231	266
Chlamydia trachomatis infection						
No	1065 (97.2) ***	645 (97.7)	420 (96.3)	444 (90.4)	214 (93.9)	230 (87.5)
Yes, in the last six months	8 (0.7)	4 (0.6)	4 (0.9)	4 (0.8)	1 (0.4)	3 (1.1)
Yes, over six months ago	23 (2.1)	11 (1.7)	12 (2.8)	43 (8.8)	13 (5.7)	30 (11.4)
(n)	1096	660	436	491	228	263
Missing	9 (0.8)	8 (1.2)	1 (0.2)	6 (1.2)	3 (1.3)	3 (1.1)
Total	1105	668	437	497	231	266
Genital herpes						
No	1084 (98.9)	654 (99.1)	430 (98.6)	483 (98.4)	223 (98.2)	260 (98.5)
Yes, in the last six months	4 (0.4)	2 (0.3)	2 (0.5)	1 (0.2)	0 (0.0)	1 (0.4)
Yes, over six months ago	8 (0.7)	4 (0.6)	4 (0.9)	7 (1.4)	4 (1.8)	3 (1.1)
(n)	1096	660	436	491	227	264
Missing	9 (0.8)	8 (1.2)	1 (0.2)	6 (1.2)	4 (1.7)	2 (1.2)
Total	1105	668	437	497	231	266
Human papilloma virus						
No	1088 (99.3)	657 (99.4)	431 (99.1)	485 (98.6)	225 (98.7)	260 (98.5)
Yes, in the last six months	3 (0.3)	1 (0.2)	2 (0.5)	1 (0.2)	0 (0.0)	1 (0.4)
Yes, over six months ago	5 (0.5)	3 (0.5)	2 (0.5)	6 (1.2)	3 (1.3)	3 (1.1)
(n)	1096	661	435	492	228	264
Missing	9 (0.8)	7 (1.0)	2 (0.5)	5 (1.0)	3 (1.3)	2 (0.8)
Total	1105	668	437	497	231	266
Lymphogranuloma venereum (LGV)						
No	1090 (99.5)	656 (99.4)	434 (99.5)	487 (99.0)	225 (98.7)	262 (99.2)
Yes, in the last six months	2 (0.2)	1 (0.2)	1 (0.2)	1 (0.2)	0 (0.0)	1 (0.4)
Yes, over six months ago	4 (0.4)	3 (0.5)	1 (0.2)	4 (0.8)	3 (1.3)	1 (0.4)
(n)	1096	660	436	492	228	264
Missing	9 (0.8)	8 (1.2)	1 (0.2)	5 (1.0)	3 (1.3)	2 (0.8)
Total	1105	668	437	497	231	266
Tuberculosis						
No	1069 (97.6)	644 (97.6)	425 (97.7)	483 (98.4)	222 (98.2)	261 (98.5)
Yes, in the last six months	6 (0.5)	3 (0.5)	3 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)
Yes, over six months ago	20 (1.8)	13 (2.0)	7 (1.6)	8 (1.6)	4 (1.8)	4 (1.5)
(n)	1095	660	435	491	226	265
Missing	10 (0.9)	8 (1.2)	2 (0.5)	6 (1.2)	5 (2.2)	1 (0.4)
Total	1105	668	437	497	231	266

* p<0.05 ** p<0.01 *** p<0.001

3.6. HIV prevalence in the SGS-AFCAR Survey

Of the 1,602 participants recruited, 96.5% (1,546/1,602) provided a DBS sample for biological analysis, and only one sample was insufficient to perform the HIV test. (Table 43)

Overall, HIV prevalence among participants in the SGS-AFCAR Survey who provided an adequate biological sample for laboratory tests was **1.4% [95% CI: 0.9–2.1%], for a total of 22 HIV-positive samples.**

For all participants recruited from the **sub-Saharan African community**, HIV prevalence was **1.7% [95% CI: 1.1–2.6%], for a total of 18 HIV-positive tests.** HIV prevalence rates by sex were:

- In men: **1.2% [95% CI: 0.6–2.4%], for a total of 8 HIV-positive tests.**
- In women: **2.4% [95% CI: 1.2–4.2%], for a total of 10 HIV-positive tests.**
- Also, just over one-third of HIV-positive participants [**38.9% (7/18)**] knew their HIV status. (Table 43)

For all participants recruited from the **English-speaking Caribbean community**, prevalence was **0.8% [95% CI: 0.3–2.1%], for a total of 4 HIV-positive tests.** HIV prevalence rates by sex were:

- In men: **0.9% [95% CI: 0.2–3.2%], for a total of 2 HIV-positive tests.**
- In women: **0.8% [95% CI: 0.2–2.8%], for a total of 2 HIV-positive tests.**
- There were no self-reported HIV-positive participants. (Table 43)

Table 43: HIV seroprevalence and known HIV status

(ELIGIBLE WITH RESPECT TO SEROPREVALENCE: Respondents who provided an adequate biological sample for a laboratory test)

(ELIGIBLE WITH RESPECT TO KNOWLEDGE OF STATUS: Respondents who provided an adequate biological sample for a laboratory test and indicated their HIV status)

	Total SGS-AFCAR	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
		Total	Men	Women	Total	Men	Women
HIV Seroprevalence ^a							
Seropositivity	22 (1.4)	18 (1.7)	8 (1.2)	10 (2.4)	4 (0.8)	2 (0.9)	2 (0.8)
[95% CI]	[0.9-2.1]	[1.1-2.6]	[0.6-2.4]	[1.2-4.2]	[0.3-2.1]	[0.2-3.2]	[0.2-2.8]
Seronegativity	1523 (98.6)	1055 (98.3)	642 (98.8)	413 (97.6)	468 (99.2)	215 (99.1)	253 (99.2)
(n)	1545	1073	650	423	472	217	255
No DBS	56 (3.5)	31 (2.8)	18 (2.7)	13 (3.0)	25 (5.0)	14 (6.1)	11 (4.1)
Insufficient quantity	1 (0.1)	1 (0.1)	0 (0.0)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Total	1602	1105	668	437	497	231	266
Knowledge of their seropositivity ^b							
Knew	7 (31.8)	7 (38.9)	2 (25.0)	5 (50.0)	0 (0.0)	0 (0.0)	0 (0.0)
Didn't know	15 (68.2)	11 (61.1)	6 (75.0)	5 (50.0)	4 (100.0)	2 (100.0)	2 (100.0)
(n)	22	18	8	10	4	2	2

^a A positive result indicated a current HIV infection (based on DBS test results).

^b Respondents who had positive DBS (biological sample) lab test results and reported that their most recent test was positive were included in the "Knew their HIV status" category. Respondents who had never undergone testing or had not received the results of their most recent test, as well as respondents who did not know the result of their last test or reported that their last test was negative, but had positive DBS (biological sample) test results were included in the "Didn't know their HIV status" category.

For sub-Saharan Africa, the distribution of percentages of HIV-positive tests (n=18) (Figure 10) and HIV prevalence based on geographic area of birth was:

- **Central Africa** (n=427): **44.4% (8/18)** of HIV-positive tests; HIV prevalence: **1.9% [95% CI: 0.8–3.6%]**.
- **West Africa** (n=398): **22.2% (4/18)** of HIV-positive tests; HIV prevalence: **1.0% [95% CI: 0.4–2.6%]**.
- **East Africa** (n=222): **22.2% (4/18)** of HIV-positive tests; HIV prevalence: **1.9% [95% CI: 0.7–4.6%]**.
- **Southern Africa** (n=26): **11.1% (2/18)** of HIV-positive tests; HIV prevalence: **7.7% [95% CI: 2.1–24.1%]**.




Figure 10: Distribution of the number of HIV-positive cases by geographical area of birth

The majority of HIV-positive tests were found in the 40–49 age group [34.9% (12/22)], followed by the 30–39 age group [31.8% (7/22)]. All HIV-positive women were in these two age groups.

Age was significantly associated with HIV [OR=1.10; 95% CI: 1.05–1.16; $p<0.0001$]. Compared to the 15–29 reference age group, the 40–49 age group was most at risk for HIV. (Figure 11)

In the two target communities, single women accounted for 58.3% (7/12) of HIV-positive tests.



Figure 11: Percentage of HIV-positive test results by sex, age group and marital status

3.7. Seroprevalence of HCV infection

Prevalence of HCV infection among participants in the SGS-AFCAR Survey who provided an adequate biological sample for laboratory tests was **0.5% [95% CI: 0.2–1.01%], for a total of 8 positive tests for HCV.** (Table 44)

For all participants recruited from the **sub-Saharan African community**, HCV prevalence was **0.6% [95% CI: 0.2–1.2%], for a total of 6 positive tests for HCV**, including 4 in women. (Table 44)

For all participants recruited from the **English-speaking Caribbean community**, prevalence was **0.4% [95% CI: 0.1–1.5%], for a total of 2 positive tests for HCV.**

One case of HIV and HCV coinfection was identified in a sub-Saharan African woman.

Table 44: HCV seroprevalence, HIV and HCV coinfection

(ELIGIBLE: Respondents who provided an adequate biological sample for testing)

	Total SGS-AFCAR	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
		Total	Men	Women	Total	Men	Women
HCV seroprevalence ^a							
Seropositivity ^a	8 (0.5)	6 (0.6)	2 (0.3)	4 (0.9)	2 (0.4)	1 (0.5)	1 (0.4)
[95% CI]	[0.2-1.01]	[0.2-1.2]	[0.08-1.1]	[0.3-2.4]	[0.1-1.5]	[0.08-2.5]	[0.06-2.1]
Seronegativity	1538 (99.5)	1068 (99.4)	648 (99.7)	420 (99.1)	470 (99.6)	216 (99.5)	254 (99.6)
(n)	1545	1073	650	423	472	217	255
No DBS	56 (3.5)	31 (2.8)	18 (2.7)	13 (3.0)	25 (5.0)	14 (6.1)	11 (4.1)
Insufficient quantity	1 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	1 (0.4)	0 (0.0)
Total	1602	1105	668	437	497	231	266
HIV and HCV coinfection							
Yes	1 (12.5)	1 (16.7)	0 (0.0)	1 (25.0)	0 (0.0)	0 (0.0)	0 (0.0)
(n)	8	6	2	4	2	1	1

^a A positive result indicates past or present HCV infection and does not discriminate acute from chronic or resolved infections.

3.8. Sources of information, knowledge and beliefs about HIV infection

Information on HIV/AIDS tended to come more from the mass media (television, newspapers, radio, etc.) than the medical and public health community. (Table 45)

Table 45: Sources of information on HIV/AIDS

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Television	802 (73.2) ***	494 (74.8)	308 (70.6)	320 (64.5)	154 (67.0)	166 (62.4)
Newspapers	579 (52.8) ***	360 (54.5)	219 (50.2)	202 (40.7)	99 (43.0)	103 (38.7)
Radio	527 (51.9) ***	357 (54.1)	212 (48.6)	192 (38.7)	97 (42.2)	95 (35.7)
Friends	565 (51.6) *	357 (54.1) *	208 (47.7)	226 (45.6)	111 (48.3)	115 (43.2)
School	520 (47.4) ***	309 (46.8)	211 (48.4)	156 (31.5)	76 (33.0)	80 (30.1)
Social media	480 (43.8)	300 (45.5)	180 (41.3)	205 (41.3)	95 (41.3)	110 (41.4)
Advertising	466 (42.5) ***	281 (42.6)	185 (42.4)	137 (27.6)	61 (26.5)	76 (28.6)
Family	463 (42.2)	282 (42.7)	181 (41.5)	196 (39.5)	101 (43.9)	95 (35.7)
Website	457 (41.7) *	276 (41.8)	181 (41.5)	178 (35.9)	81 (35.2)	97 (36.5)
Doctors / Other health professionals	439 (40.1)	251 (38.0)	188 (43.1)	141 (28.4)	56 (24.3)	85 (32.0)
Clinic/Hospital/CLSC	344 (31.4) ***	210 (31.8)	134 (30.7)	104 (21.0)	46 (20.0)	58 (21.8)
Work	325 (29.7) ***	203 (30.8)	122 (28.0)	95 (19.2)	42 (18.3)	53 (19.9)
Conference/Seminars	243 (22.2) ***	158 (23.9)	85 (19.5)	43 (8.7)	19 (8.3)	24 (9.0)
Local or provincial public health services	210 (19.2) ***	136 (20.6)	74 (17.0)	43 (8.7)	20 (8.7)	23 (8.6)
Community organization	200 (18.2) ***	128 (19.4)	72 (16.5)	43 (8.7)	19 (8.3)	24 (9.0)
Churches	190 (17.3) **	111 (16.8)	79 (18.1)	54 (10.9)	27 (11.7)	27 (10.2)
Public Health Agency of Canada	147 (13.4)	96 (14.5)	51 (11.7)	54 (10.9)	27 (11.7)	27 (10.2)
No information	38 (3.5)	21 (3.2)	17 (3.9)	24 (4.8)	11 (4.8)	13 (4.9)

* p<0.05 ** p<0.01 *** p<0.001

Most participants correctly answered questions on HIV prevention and transmission. Participants from sub-Saharan Africa had a higher level of knowledge about HIV transmission modes than English-speaking Caribbeans. (Table 46 and Figure 12)

However, 31.1% (496/1,595) of participants [25.0% of sub-Saharan Africans vs. 44.6% of English-speaking Caribbeans] answered incorrectly or did not know the answer to the question on the risk of HIV infection from mosquito bites. The same applied to 17.7% (282/1,596) of participants [10.4% of sub-Saharan Africans vs. 33.9% of English-speaking Caribbeans] regarding the question on the risk of HIV infection by sharing a meal with an infected person. While almost all respondents understood that people with HIV could appear healthy, 6% (96/1,596) [5.1% of sub-Saharan Africans vs. 8.1% of English-speaking Caribbeans] did not answer the question or did not know the answer. (Table 46)

Table 46: Knowledge and beliefs about HIV

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
Having sex with one faithful, uninfected partner can reduce the risk of HIV transmission						
Yes ^a	957 (87.0) ***	584 (88.0)	373 (85.6)	375 (75.6)	171 (74.3) **	204 (76.7)
No	105 (9.5)	64 (9.6)	41 (9.4)	90 (18.1)	36 (15.7)	54 (20.3)
Don't know	38 (3.5)	16 (2.4)	22 (5.0)	31 (6.3)	23 (10.0)	8 (3.0)
(n)	1100	664	436	496	230	266
Missing	5 (0.5)	4 (0.6)	1 (0.2)	1 (0.2)	1 (0.4)	0 (0.0)
Total	1105	668	437	497	231	266
Condom use reduces the risk of HIV transmission						
Yes ^a	1001 (91.2) ***	623 (94.1) ***	378 (86.9)	416 (83.9)	191 (83.0)	225 (84.6)
No	66 (6.0)	25 (3.8)	41 (9.4)	56 (11.3)	23 (10.0)	33 (12.4)
Don't know	30 (2.7)	14 (2.1)	16 (3.7)	24 (4.8)	16 (7.0)	8 (3.0)
(n)	1097	662	435	496	230	266
Missing	8 (0.7)	6 (0.9)	2 (0.5)	1 (0.2)	1 (0.4)	0 (0.0)
Total	1105	668	437	497	231	266
A person who looks healthy can have HIV						
Yes ^a	1044 (94.9) *	632 (95.2)	412 (94.5)	456 (91.9)	210 (91.3)	246 (92.5)
No	37 (3.4)	21 (3.2)	16 (3.7)	31 (6.3)	13 (5.7)	18 (6.8)
Don't know	19 (1.7)	11 (1.7)	8 (1.8)	9 (1.8)	7 (3.0)	2 (0.8)
(n)	1100	664	436	496	230	266
Missing	5 (0.5)	4 (0.6)	1 (0.2)	1 (0.2)	1 (0.4)	0 (0.0)
Total	1105	668	437	497	231	266

	Sub-Saharan Africa n=1105 n (%)			English-Speaking Caribbean n=497 n (%)		
	Total	Men	Women	Total	Men	Women
A person can become infected with HIV from mosquito bites						
Yes	156 (14.2) ***	99 (14.9)	57 (13.1)	124 (25.1)	70 (30.6) *	54 (20.3)
No ^a	825 (75.0)	500 (75.3)	325 (74.5)	274 (55.4)	113 (49.3)	161 (60.5)
Don't know ^b	119 (10.8)	65 (9.8)	54 (12.4)	97 (19.6)	46 (20.1)	51 (19.2)
(n)	1100	664	436	495	229	266
Missing	5 (0.5)	4 (0.6)	1 (0.2)	2 (0.4)	2 (0.9)	0 (0.0)
Total	1105	668	437	497	231	266
A person can become infected with HIV by sharing a meal with someone who is infected						
Yes	60 (5.5) ***	45 (6.8)	15 (3.4)	112 (22.6)	48 (20.9)	64 (24.1)
No ^a	986 (89.6)	585 (88.1)	401 (92.0)	328 (66.1)	153 (66.5)	175 (65.8)
Don't know ^b	54 (4.9)	34 (5.1)	20 (4.6)	56 (11.3)	29 (12.6)	27 (10.2)
(n)	1100	664	436	496	230	266
Missing	5 (0.5)	4 (0.6)	1 (0.2)	1 (0.2)	1 (0.4)	0 (0.0)
Total	1105	668	437	497	231	266
Currently there is a cure for HIV, meaning a treatment that can completely eliminate it						
Yes	47 (4.3) ***	30 (4.5)	17 (3.9)	61 (12.3)	36 (15.7) **	25 (9.4)
No ^a	943 (85.7)	572 (86.1)	371 (85.1)	350 (70.6)	145 (63.0)	205 (77.1)
Don't know ^b	110 (10.0)	62 (9.3)	48 (11.0)	85 (17.1)	49 (21.3)	36 (13.5)
(n)	1100	664	436	496	230	266
Missing	5 (0.5)	4 (0.6)	1 (0.2)	1 (0.2)	1 (0.4)	0 (0.0)
Total	1105	668	437	497	231	266

^a: Correct answers

^b The answer "Don't know" (DK) must be recorded as an incorrect answer.

* p<0.05 ** p<0.01 *** p<0.001

The distribution of knowledge and assumptions regarding HIV transmission, by age group and highest level of education in both communities, are presented in Figures 12 and 13.




Figure 12: Percentage of knowledge and assumptions about HIV transmission in both communities




Figure 13: Percentage of knowledge and assumptions about HIV transmission in both communities by age group

The level of education appears to correlate with the level of knowledge about modes of HIV transmission. This highlights the knowledge gaps that must be addressed taking into account the social and cultural diversity of the target populations. (Figure 14)




Figure 14: Percentage of knowledge and assumptions regarding HIV transmission in both communities by highest level of education

3.9. Access to and use of health systems and services

In total, 60.7% (963/1,587) of participants recruited in the two communities did not have a family doctor, i.e. 67.1% of sub-Saharan Africans vs. 46.5% of English-speaking Caribbeans ($p<0.0001$). However, women in both communities had greater access to a family doctor than men did ($p<0.0001$).

There was a discrepancy in results regarding the question about accessing the needle exchange or harm reduction service location. Participants may have misunderstood the question. However, it seems that some people go there to get condoms. Particular attention will be paid to this question in the next second generation HIV surveillance survey. (Table 47)

Table 47: Type of access to health care systems

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n (%) N=1105			English-Speaking Caribbean n (%) N=497		
	Total	Men	Women	Total	Men	Women
Do you have a family doctor or primary health care provider?						
Yes	360 (32.9) ***	180 (27.3) ***	180 (41.5)	264 (53.5)	102 (44.3) ***	162 (61.6)
No	734 (67.1)	480 (72.7)	254 (58.5)	229 (46.5)	128 (55.7)	101 (38.4)
(n)	1094	660	434	493	230	263
Missing	11 (1.0)	8 (1.2)	3 (0.7)	4 (0.8)	1 (0.4)	3 (1.1)
Total	1105	668	437	497	231	266
Medical care or consultations in the last 12 months						
Yes	743 (67.4)	404 (60.8) ***	339 (77.6)	323 (65.5)	128 (55.7) ***	195 (74.1)
No	359 (32.6)	261 (39.2)	98 (22.4)	170 (34.5)	102 (44.3)	68 (25.9)
(n)	1102	665	437	493	230	263
Missing	3 (0.3)	3 (0.4)	0 (0.0)	4 (0.8)	1 (0.4)	3 (1.1)
Total	1105	668	437	497	231	266
Location of consultations in the last 12 months						
Family doctor or care provider	223 (30.2) ***	113 (28.3)	110 (32.5)	155 (48.4)	48 (37.8) **	107 (55.4)
Hospital	414 (55.9)	212 (52.7)	202 (59.8)	178 (55.6)	79 (62.2)	99 (51.3)
Medical clinic or walk-in clinic	377 (51.1)	197 (49.3)	180 (53.3)	157 (49.1)	56 (44.1)	101 (52.3)
CLSC	247 (33.5)	107 (26.7) ***	140 (41.5)	123 (38.6)	50 (39.4)	73 (38.0)
Community organization	34 (4.6)	16 (4.0)	18 (5.3)	13 (4.1)	7 (5.5)	6 (3.1)
Alternative or traditional medicine	14 (1.9)	6 (1.5)	8 (2.4)	2 (0.6)	1 (0.8)	1 (0.5)
Needle exchange or harm reduction program	139 (18.8)	71 (17.7)	68 (20.1)	45 (14.0)	17 (13.4)	28 (14.4)
Detox centre	2 (0.3)	2 (0.5)	0 (0.0)	2 (0.6)	1 (0.8)	1 (0.5)
Mental health or addiction treatment centre	7 (0.9)	4 (1.0)	3 (0.9)	7 (2.2)	4 (3.1)	3 (1.5)
Sexual health centre or facility	3 (0.4)	3 (0.7)	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.5)
Prison / Corrections system	3 (0.4) *	3 (0.7)	0 (0.0)	5 (1.6)	3 (2.4)	2 (1.0)
Other	3 (0.4)	3 (0.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

In all, 66.8% (1,066/1,595) of survey participants had received medical treatment or consulted a health professional in the last 12 months; more women than men in both communities were in this group ($p<0.0001$). (Table 48)

Of those requiring medical care in the 12 previous months, 15.1% (223/1,473) reported having difficulty accessing necessary health care services during this period [15.7% of sub-Saharan Africans and 10.1% of English-speaking Caribbeans]. The main types of difficulties they encountered are presented in Table 48.

Table 48: Lack of access to health care systems

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n (%) N=1105			English-Speaking Caribbean n (%) N=497		
	Total	Men	Women	Total	Men	Women
Have you had difficulties accessing the health care you needed in the last 12 months?						
Yes	173 (15.7) *	87 (13.1) **	86 (19.7)	50 (10.1)	23 (10.0)	27 (10.2)
No	842 (76.5)	515 (77.7)	327 (74.8)	408 (82.4)	189 (82.2)	219 (82.4)
No need	85 (7.7)	61 (9.2)	24 (5.5)	37 (7.5)	18 (7.8)	19 (7.2)
(n)	1100	663	437	495	230	265
Missing	5 (0.5)	5 (0.7)	0 (0.0)	2 (0.4)	1 (0.4)	1 (0.4)
Total	1105	668	437	497	231	266
Type of difficulties encountered						
Long wait times	120 (69.4) *	57 (65.5)	63 (73.3)	25 (50.0)	11 (47.8)	14 (51.9)
Difficulty getting an appointment, diagnosis or referral	86 (49.7)	37 (42.5)	49 (57.0)	19 (38.0)	5 (21.7) *	14 (51.9)
Service not covered by insurance, lack of financial means	31 (17.9)	17 (19.5)	14 (16.3)	12 (24.0)	4 (17.4)	8 (29.6)
No health care provider available in your area	15 (8.7) *	6 (6.6)	9 (10.5)	10 (20.0)	2 (8.7)	8 (29.6)
Don't know how to find health care	11 (6.4)	4 (4.6)	7 (8.1)	5 (10.0)	2 (8.7)	3 (11.1)
Specific service not available in the area	10 (5.8)	6 (6.9)	4 (4.7)	7 (14.0)	2 (8.7)	5 (18.5)
Transportation problem	4 (2.3)	4 (4.6) *	0 (0.0)	2 (4.0)	1 (4.3)	1 (3.7)
Inability to leave home because of a health problem	4 (2.3)	2 (2.3)	2 (2.3)	1 (2.0)	1 (4.3)	0 (0.0)
No health insurance card	4 (2.3)	2 (2.3)	2 (2.3)	1 (2.0)	1 (4.3)	0 (0.0)
Personal or family responsibilities preventing access to health care	3 (1.7)	2 (2.3)	1 (1.2)	3 (6.0)	0 (0.0)	3 (11.1)
Health care provider was not culturally sensitive or was racist	3 (1.7) **	2 (2.3)	1 (1.2)	6 (12.0)	2 (8.7)	4 (14.8)
Health care provider was of the opposite sex	2 (1.2)	0 (0.0)	2 (2.3)	1 (2.0)	0 (0.0)	1 (3.7)
Language barrier	1 (0.6)	1 (1.1)	0 (0.0)	2 (4.0)	0 (0.0)	2 (7.4)
Other	13 (7.5)	7 (8.0)	6 (7.0)	4 (8.0)	3 (13.0)	1 (3.7)

* $p<0.05$ ** $p<0.01$ *** $p<0.001$

Table 49: History of immunization against diseases and diagnosis

(ELIGIBLE: All respondents surveyed)

	Sub-Saharan Africa n (%) N=1105			English-Speaking Caribbean n (%) N=497		
	Total	Men	Women	Total	Men	Women
Hepatitis B immunization						
No	414 (43.8) ***	276 (49.1) ***	138 (35.9)	252 (57.7)	126 (63.6)	126 (52.7)
Yes, in the last six months	68 (7.2)	31 (5.5)	37 (9.6)	24 (5.5)	9 (4.5)	15 (6.3)
Yes, over six months ago	464 (49.0)	255 (45.4)	209 (54.4)	161 (36.8)	63 (31.8)	98 (41.0)
(n)	946	562	384	437	198	239
Missing	156 (14.4)	106 (15.9)	53 (12.1)	60 (12.1)	33 (14.3)	27 (10.2)
Total	1105	668	437	497	231	266
Immunization against human papillomavirus						
No	835 (87.6)	541 (92.5) ***	294 (79.9)	376 (85.3)	181 (90.5) *	195 (80.9)
Yes, in the last six months	20 (2.1)	6 (1.0)	14 (3.8)	10 (2.3)	3 (1.5)	7 (2.9)
Yes, over six months ago	98 (10.3)	38 (6.5)	60 (16.3)	55 (12.5)	16 (8.0)	39 (16.2)
(n)	953	585	368	441	200	241
Missing	152 (13.8)	83 (12.4)	69 (15.8)	56 (11.3)	31 (13.4)	25 (9.4)
Total	1105	668	437	497	231	266
Have you been tested for tuberculosis?						
No	534 (52.2) ***	326 (52.6)	208 (51.6)	327 (72.0)	169 (80.9) ***	158 (64.5)
Yes	489 (47.8)	294 (47.4)	195 (48.4)	127 (28.0)	40 (19.1)	87 (35.5)
(n)	1023	620	403	454	209	245
Missing	82 (7.4)	48 (7.2)	34 (7.8)	43 (8.7)	22 (9.5)	21 (7.9)
Total	1105	668	437	497	231	266

* p<0.05 ** p<0.01 *** p<0.001

3.10. HIV prevalence rate based on certain characteristics

Table 50: HIV prevalence based on certain characteristics and behaviours

(ELIGIBLE: Respondents who provided an adequate biological sample for a laboratory test)

	Sub-Saharan Africa						English-Speaking Caribbean					
	Total n=1073		Men n=650		Women n=423		Total n=472		Men n=217		Women n=255	
	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]
HIV (Positive)	1073	18 (1.7 [1.1-2.6])	650	8 (1.2 [0.6-2.4])	423	10 (2.4 [1.2-4.2])	472	4 (0.8 [0.3-2.1])	217	2 (0.9 [0.2-3.2])	255	2 (0.8 [0.2-2.8])
Sexual orientation												
Heterosexual	1048 (97.7)	18 (1.7 [1.1-2.6])	634 (97.5)	8 (1.2 [0.6-2.4])	414 (97.9)	10 (2.4 [1.3-4.3])	450 (95.3)	4 (0.9 [0.3-2.2])	212 (97.7)	2 (0.9 [0.2-3.2])	238 (93.3)	2 (0.8 [0.2-3.0])
Other	25 (2.3)	0	16 (2.5)	0	9 (2.1)	0	22 (4.7)	0	5 (2.3)	0	17 (6.7)	0
Age in years		**				***		**				*
Average (±SD)	30.4 (9.1)	37.5 (9.5)	31.09 (9.2)	32.8 (11.8)	29.4 (8.9)	41.2 (5.3)	31.3 (10.2)	44.7 (4.7)	30.9 (10.6)	41.5 (4.9)	31.5 (9.9)	48 (0.0)
[Min-Max]	[15-49]	[19-49]	[15-49]	[19-47]	[15-49]	[33-49]	[15-49]	[38-48]	[15-49]	[38-45]	[15-49]	[48]
Age group		**				***						
15–19 years	178 (16.6)	2 (1.1 [0.3-4.0])	106 (16.3)	2 (1.9 [0.5-6.6])	72 (17.0)	0	85 (18.0)	0	42 (19.4)	0	43 (16.9)	0
20–29 years	308 (28.7)	1 (0.3 [0.05-1.8])	161 (24.8)	1 (0.6 [0.1-3.4])	147 (34.8)	0	116 (24.6)	0	57 (26.3)	0	59 (23.1)	0
30–39 years	399 (37.2)	6 (1.5 [0.7-3.2])	258 (39.7)	2 (0.7 [0.2-2.7])	141 (33.3)	4 (2.8 [1.1-7.06])	148 (31.4)	1 (0.7 [0.1-3.8])	65 (30.0)	1 (1.5 [0.2-8.2])	83 (32.5)	0
40–49 years	188 (17.5)	9 (4.7 [2.5-8.8])	125 (19.2)	3 (2.4 [0.8-6.8])	63 (14.9)	6 (9.5 [4.4-19.2])	123 (26.1)	3 (2.4 [0.8-7.2])	53 (24.4)	1 (1.9 [0.3-9.9])	70 (27.5)	2 (2.8 [0.7-9.8])
Region												
Central Africa	427 (39.8)	8 (1.9 [0.8-3.6])	261 (40.2)	2 (0.8 [0.2-2.7])	166 (39.2)	6 (3.6 [1.6-7.6])	—	—	—	—	—	—
West Africa	398 (37.1)	4 (1.04 [0.4-2.6])	246 (37.8)	3 (1.2 [0.4-3.5])	152 (35.9)	1 (0.7 [0.1-3.6])	—	—	—	—	—	—
East Africa	222 (20.7)	4 (1.9 [0.7-4.6])	126 (19.4)	2 (1.6 [0.4-5.6])	96 (22.7)	2 (2.1 [0.5-7.2])	—	—	—	—	—	—
Southern Africa	26 (2.4)	2 (7.7 [2.1-24.1])	17 (2.6)	1 (5.9 [1.04-26.9])	9 (2.1)	1 (11.1 [1.9-43.5])	—	—	—	—	—	—
Level of education												
≤ High school education	322 (30.1)	6 (1.9 [0.8-4.1])	175 (27.1)	2 (1.1 [0.3-4.0])	147 (34.8)	4 (2.7 [1.06-6.7])	294 (62.4)	2 (0.7 [0.1-2.5])	154 (71.0)	1 (0.6 [0.1-3.5])	140 (55.1)	1 (0.7 [0.1-3.9])
> High school education	746 (69.9)	12 (1.6 [0.9-2.8])	470 (72.9)	6 (1.3 [0.5-2.7])	276 (65.2)	6 (2.2 [1-4.6])	177 (37.6)	2 (1.1 [0.3-4.2])	63 (29.0)	1 (1.6 [0.2-8.4])	114 (44.9)	1 (0.9 [0.1-4.8])
Annual family income							*					
<\$15,000	514 (47.9)	9 (1.7 [0.9-3.3])	308 (47.4)	6 (1.9 [0.8-4.1])	206 (48.7)	4 (1.9 [0.7-4.8])	173 (36.6)	0	85 (39.2)	0	88 (34.5)	0
≥\$15,000	452 (42.1)	9 (2.0 [1.06-3.7])	285 (43.8)	3 (1.1 [0.3-3.04])	167 (39.5)	6 (3.6 [1.6-7.6])	184 (39.0)	4 (2.2 [0.8-5.6])	78 (35.9)	2 (2.6 [0.7-8.8])	106 (41.6)	2 (1.9 [0.5-6.6])
Not reported	107 (10.0)	0	57 (8.8)	0	50 (11.8)	0	115 (24.4)	0	54 (24.9)	0	61 (23.9)	0
Residency status						*						
Citizen / Permanent resident	910 (84.8)	14 (1.5 [0.9-2.6])	546 (84.0)	6 (1.1 [0.5-2.3])	364 (86.1)	8 (2.2 [1.1-4.2])	428 (91.6)	4 (0.9 [0.3-2.3])	195 (90.3)	2 (1.0 [0.2-3.6])	233 (92.8)	2 (0.9 [0.2-3.07])
Visa / Work permit	100 (9.3)	1 (1.0 [0.1-5.5])	59 (9.1)	1 (1.7 [0.2-8.9])	41 (9.7)	0	12 (2.6)	0	6 (2.8)	0	6 (2.4)	0
Refugee/Other	63 (5.9)	3 (4.8 [1.8-14.6])	45 (6.9)	1 (2.2 [0.3-11.5])	18 (4.3)	2 (11.1 [3.1-32.8])	27 (5.8)	0	15 (6.9)	0	12 (4.8)	0
Immigration date												
<1990	18 (1.7)	1 (5.6 [0.9-25.7])	12 (1.9)	0	6 (1.4)	1 (16.7 [3.0-56.3])	107 (23.0)	1 (1.0 [0.1-5.2])	60 (27.8)	1 (1.7 [0.2-8.8])	47 (18.7)	0
1990–1999	125 (11.7)	3 (2.4 [0.8-6.8])	64 (9.9)	1 (1.6 [0.2-8.3])	61 (14.5)	2 (3.3 [0.9-11.2])	123 (26.3)	0	51 (23.6)	0	72 (28.7)	0
≥2000	927 (86.6)	14 (1.5 [0.9-2.5])	572 (88.3)	7 (1.2 [0.6-2.5])	355 (84.1)	7 (2.0 [0.9-4.0])	237 (50.7)	3 (1.3 [0.4-3.8])	105 (48.6)	1 (1.0 [0.1-5.1])	132 (52.6)	2 (1.5 [0.4-5.3])
Marital status												
Single	551 (51.5)	8 (1.5 [0.7-2.9])	343 (53.0)	3 (0.9 [0.2-2.5])	208 (49.3)	5 (2.4 [1.03-5.5])	281 (59.7)	3 (1.1 [0.3-3.2])	126 (58.1)	1 (0.8 [0.1-4.3])	155 (61.0)	2 (1.3 [0.3-4.5])
Living together as a couple	358 (33.5)	9 (2.5 [1.3-4.7])	204 (31.5)	4 (2.0 [0.7-4.9])	154 (36.5)	5 (3.2 [1.3-7.3])	104 (22.1)	0	49 (22.6)	0	55 (21.7)	0
Couple not living together	160 (15.0)	1 (0.6 [0.1-3.5])	100 (15.5)	1 (1.0 [0.1-5.4])	60 (14.2)	0	86 (18.3)	1 (1.2 [0.2-6.6])	42 (19.4)	1 (2.4 [0.4-12.3])	44 (17.3)	0
Employment status		*	**									
Unemployed	132 (12.4)	6 (4.5 [2.1-9.5])	74 (11.5)	4 (5.4 [2.1-13.09])	58 (13.9)	2 (3.4 [0.9-11.7])	103 (22.0)	0	47 (21.6)	0	56 (22.3)	0
Student	460 (43.3)	4 (0.9 [0.3-2.2])	265 (41.1)	2 (0.8 [0.2-2.7])	195 (46.8)	2 (1.0 [0.2-3.6])	138 (29.5)	1 (0.7 [0.1-4.1])	54 (24.9)	1 (1.9 [0.3-9.7])	84 (33.5)	0
Full-time or part-time employee	470 (44.3)	8 (1.7 [0.8-3.3])	306 (47.4)	2 (0.7 [0.1-2.3])	164 (39.3)	6 (3.7 [1.6-7.7])	227 (48.5)	3 (1.3 [0.4-4.0])	116 (53.5)	1 (0.9 [0.1-4.7])	111 (44.2)	2 (1.8 [0.5-6.3])

*STBBI history: Chlamydia, gonorrhea and syphilis

* p<0.05 ** p<0.01 *** p<0.001

	Sub-Saharan Africa						English-Speaking Caribbean					
	Total n=1073		Men n=650		Women n=423		Total n=472		Men n=217		Women n=255	
	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]	n (%)	HIV positive n (%) [95% CI]
Number of sexual partners (last 12 months)									**			
None	126 (12.6)	1 (0.8 [0.1-4.9])	55 (9.2)	1 (1.8 [0.3-9.6])	71 (17.5)	0	33 (7.4)	0	11 (5.4)	0	22 (9.1)	0
1 partner	160 (16.0)	1 (0.6 [0.1-3.4])	69 (11.6)	0	91 (22.5)	1 (1.1 [0.1-5.9])	31 (7.0)	1 (3.2 [0.6-16.1])	8 (3.9)	1 (12.5 [2.2-47.09])	23 (9.5)	0
2-3 partners	237 (23.7)	4 (1.7 [0.6-4.2])	104 (17.4)	1 (1.0 [0.1-5.2])	133 (32.8)	3 (2.2 [0.7-6.4])	89 (20.0)	0	13 (6.3)	0	76 (31.5)	0
≥4 partners	479 (47.8)	12 (2.5 [1.3-4.3])	369 (61.8)	6 (1.6 [0.7-3.5])	110 (27.2)	6 (5.4 [2.5-11.4])	293 (65.7)	3 (1.0 [0.3-2.9])	173 (84.4)	1 (0.6 [0.1-3.2])	120 (49.8)	2 (1.6 [0.4-6.4])
Condom use during last sexual encounter (No)	425 (57.0)	10 (2.4 [1.2 -4.2])	250 (52.4)	4 (1.6 [0.6-4.04])	175 (64.8)	6 (3.4 [1.5-7.2])	191 (52.6)	2 (1.0 [0.3-3.9])	93 (51.7)	1 (1.1 [0.2-5.8])	99 (53.6)	1 (1.0 [0.1-5.5])
≥2 sexual partners (last 12 months) (Yes) and condom use during last sexual encounter (No)	335 (54.3)	10 (3.0 [1.6-5.4])	209 (50.0)	4 (1.9 [0.7-4.8])	126 (63.3)	6 (4.8 [2.2-10])	175 (53.0)	2 (1.1 [0.3-4.0])	91 (53.5)	1 (1.1 [0.1-5.9])	84 (52.5)	1 (1.2 [0.2-6.4])
Casual partner (last 12 months) (Yes)	247 (33.2)	3 (1.2 [0.4-3.6])	205 (43.2)	2 (1.0 [0.2-3.4])	42 (15.6)	1 (2.4 [0.4-12.3])	168 (46.3)	2 (1.2 [0.3-4.5])	114 (63.3)	1 (0.9 [0.1-4.8])	54 (29.5)	1 (1.9 [0.3-9.7])
Condom use during last sexual encounter with a casual partner (last 12 months) (No)	52 (21.8)	1 (1.9 [0.3-10.1])	38 (18.5)	1 (2.6 [0.4-13.4])	14 (41.2)	0	49 (29.7)	1 (2.0 [0.3-10.7])	26 (23.0)	0	23 (44.2)	1 (4.3 [0.7-20.9])
Casual partner (last 12 months) (Yes) and condom use during last sexual encounter with a casual partner (No)	52 (7.0)	1 (1.9 [0.3-10.1])	38 (8.0)	1 (2.6 [0.4-13.5])	14 (5.2)	0	49 (13.5)	1 (2.0 [0.3-10.7])	26 (14.4)	0	23 (12.6)	1 (4.3 [0.7-20.9])
Sexual intercourse when travelling in an endemic country (Yes)	135 (34.3)	2 (1.5 [0.4-5.3])	103 (43.3)	1 (1.0 [0.1-5.3])	32 (20.5)	1 (3.1 [0.5-15.7])	41 (27.0)	1 (2.4 [0.4-12.6])	24 (37.5)	1 (4.2 [0.7-20.2])	17 (19.3)	0
Condom use (Not always) when travelling in an endemic country, with a sexual partner living there	81 (60.0)	2 (2.5 [0.6-8.5])	54 (52.4)	1 (1.9 [0.3-9.7])	27 (84.4)	1 (3.7 [0.6-18.2])	19 (46.3)	1 (5.3 [0.9-24.6])	8 (33.3)	1 (12.5 [2.2-47.09])	11 (64.7)	0
Last regular or casual HIV-positive sex partner (Yes)	*** 11 (1.8)	2 (18.2 [5.1-47.7])	** 9 (2.3)	1 (11.1 [1.9-43.5])	*** 2 (0.9)	1 (50 [9.4-90.5])	1 (0.3)	0	0	0	1(0.7)	0
Sex in exchange for money, drugs or other goods (Yes)	33 (4.4)	0	31 (6.5)	0	2 (0.7)	0	25 (6.9)	0	21 (11.7)	0	4 (2.2)	0
Lifetime IDU	3 (0.3)	0	2 (0.3)	0	1 (0.2)	0	1 (0.2)	0	1 (0.5)	0	0	0
Circumcision (Man)												
No	—	—	61 (9.5)	1 (1.6 [0.3-8.7])	—	—	—	—	140 (64.8)	2 (1.4 [0.3-5.05])	—	—
Yes	—	—	583 (90.5)	7 (1.2 [0.5-2.4])	—	—	—	—	76 (35.2)	0	—	—
History of STBBI [§] (Yes)	38 (3.5)	0	21 (3.2)	0	17 (4.0)	0	*** 59 (12.5)	3 (5.1 [1.7-13.9])	23 (10.6)	1 (4.3 [0.7-20.9])	*** 36 (14.1)	2 (5.6 [1.5-18.1])
HIV test history												
≤2 years	565 (68.0)	7 (1.2 [0.6-2.5])	327 (66.3)	2 (0.6 [0.16-2.2])	238 (70.4)	5 (2.1 [0.9-4.8])	198 (69.5)	0	77 (68.8)	0	121 (69.9)	0
>2 years	266 (32.0)	6 (2.2 [1.03-4.8])	166 (33.7)	2 (1.2 [0.3-4.2])	100 (29.6)	4 (4.0 [1.5-9.8])	87 (30.5)	1 (1.1 [0.2-6.2])	35 (31.3)	0	52 (30.1)	1 (1.9 [0.3-10.1])
Level of knowledge about HIV transmission mode							*					
No incorrect answers	752 (70.1)	11 (1.5 [0.8-2.6])	458 (70.5)	4 (0.9 [0.3-2.2])	294 (69.5)	7 (2.4 [1.1-4.8])	201 (42.6)	0	88 (40.6)	0	113 (44.3)	0
1 to 2 incorrect answers	294 (27.4)	7 (2.4 [1.1-4.8])	178 (27.4)	4 (2.2 [0.8-5.6])	116 (27.4)	3 (2.6 [0.8-7.3])	226 (47.9)	2 (0.9 [0.2-3.1])	106 (48.8)	2 (1.9 [0.5-6.6])	120 (47.1)	0
3 to 5 incorrect answers	27 (2.5)	0	14 (2.2)	0	13 (3.1)	0	45 (9.5)	2 (4.4 [1.2-14.8])	23(10.6)	0	22 (8.6)	2 (9.1 [2.5-27.8])
Difficulty accessing care (Yes)	169 (15.8)	3 (1.8 [0.6-5.1])	87 (13.5)	0	82 (19.4)	3 (3.7 [1.2-10.2])	47 (10.0)	1 (2.1 [0.3-11.1])	22 (10.2)	1 (4.5 [0.8-21.8])	25 (9.8)	0

In Table 50, the sociodemographic and economic profile, as well as sexual risk behaviours, history of STBBI and testing, and level of knowledge about HIV transmission mode are presented based on HIV-positive status, by sex and target community.

All results are presented, despite the small number of HIV-positive participants in certain categories and wide confidence intervals.

In both target communities, the 22 HIV-positive cases occurred among heterosexuals. Most HIV-positive tests were detected in 30–49 year-old [86.4% (19/22)]. In sub-Saharan Africans, the average age of HIV-positive individuals was 32.8 years [range 19–47 years] among men vs. 41.2 years [range 33–49 years] among women. The average age among Caribbeans was 41.5 years [range 38–45 years] among men vs. 48 years among women (n=2). Age was significantly associated with HIV ($p<0.0001$) [OR=1.10; 95% CI: 1.05–1.16].

HIV prevalence was higher in certain subgroups based on:

- **Area of origin:** Among people from Southern Africa (7.7% [2.1–24.1%]) for all sub-Saharan Africans, followed by East Africa for men (1.6% [0.4–5.6%]) and Central Africa for women (3.6% [1.6–7.6%]).
- **Level of education:** In sub-Saharan African women who had a high school education or less (2.7% [1.06–6.7%]), whereas prevalence was higher in the other three subgroups among those who had more than a high school education (African men: 1.3% [0.5–2.7%]; Caribbean men: 1.6% [0.2–8.4%]; Caribbean women: 0.9% [0.1–4.8%]).
- **Income:** Among African men whose income was <\$15,000 (1.9% [0.8–4.1%]), while prevalence was higher in the other three subgroups whose income was ≥\$15,000 (African women: 3.6% [1.6–7.6%]; Caribbean men: 2.6% [0.7–8.8%]; Caribbean women: 1.9% [0.5–6.6%]).
- **Residency status:** Among Africans who had refugee status / other (men: 2.2% [0.3–11.5%]; women: 11.1% [3.1–32.8%]).
- **Immigration year:** ≥2000 among English-speaking Caribbean women (1.5% [0.4–5.3%]) and ≤1999 for the other three subgroups (Caribbean men: 1.7% [0.2–8.8%]; African men: 1.6% [0.2–8.3%]; African women: 16.7% [3.0–56.3%]).
- **Marital status:** Among Africans living together as a couple (men: 2.0% [0.7–4.9%]; women: 3.2% [1.3–7.3%]); whereas it was higher among English-speaking Caribbean men in a relationship but not living with their partner (1.0% [0.1–5.4%]) and among single English-speaking Caribbean women (1.3% [0.3–4.5%]).
- **Employment status:** Among unemployed sub-Saharan Africans (men: 5.4% [2.1–13.0%]; women: 3.4% [0.9–11.7%]).
- **High-risk sexual behaviours:**
 - Among women who had had four or more sexual partners in the last 12 months (sub-Saharan African women: 5.4% [2.5–11.4%]; English-speaking Caribbean women: 1.6% [0.4–6.4%]).
 - Among female participants who did not always use a condom during the last sexual encounter (sub-Saharan African women: 3.4% [1.5–7.2%]; English-speaking Caribbean women: 1.0% [0.1–5.5%]).
 - Despite the small numbers, among those who had casual sex partners in the last 12 months and had not used a condom during the last sexual encounter, those who did not always use a condom when travelling in an endemic country with a sexual partner living there, and those whose last regular or casual sex partner was HIV positive.
- **History of STBBIs:** Among all English-speaking Caribbeans (5.1% [1.7–13.9%]).
- **HIV test history >2 years:** Among sub-Saharan Africans (men: 1.2% [0.3–4.2%]; women: 4.0% [1.5–9.8%]).
- **HIV transmission modes:** In those who had one or two incorrect answers among sub-Saharan Africans (2.4% [1.1–4.8%]) and English-speaking Caribbean men (1.9% [0.5–6.6%]); English-speaking Caribbean women who had three or four incorrect answers (9.1% [2.5–27.8%]).
- **Difficulty accessing health care services:** For sub-Saharan African women (3.7% [1.2–10.2%]).

4. STRENGTHS/LIMITATIONS

These findings could be used as a benchmark for monitoring trends in HIV and HCV prevalence, screening habits and behaviours associated with the acquisition and transmission of these infections among people from HIV-endemic countries, in general, but more specifically in sub-Saharan African and English-speaking Caribbean communities. Moreover, the findings should help guide the development of policies, programs and interventions to prevent and control infection with HIV and other STBBIs by targeting the needs of ethno-cultural communities on the Island of Montreal.

The SGS-AFCAR Survey is a voluntary, anonymous, cross-sectional survey; participants are recruited mainly using “time-location sampling” methods. One of the challenges of this surveillance project was related to sampling and recruitment strategies to obtain a representative sample of members of the sub-Saharan African and English-speaking Caribbean communities living on the Island of Montreal. While recognizing that there are commonalities between sub-Saharan African and Caribbean communities in general, the results highlighted differences in demographics, lifestyle and sexual behaviours.

Compared to 2006 Census data, there were more men than women in the two target communities among the 1,602 individuals recruited; the same applied to members of the 15–24 and 25–34 age groups, who are generally difficult to recruit. Also, there were fewer individuals in Montreal from certain countries in East and Southern Africa, whereas there were greater numbers of them in the other two sentinel sites (Ottawa, Toronto) selected for the E-Track national surveillance system.

The target population was difficult to access. However, the field team from the two target communities was competent and unwavering in its commitment. Through careful planning, networking and active searches, the team found the social gathering places of the various associations grouped by country of origin and community organizations, which contributed to the success of the SGS-AFCAR Survey. The recruitment strategy, which involved using a vehicle to find recruits, was most effective, and enabled the team to recruit 51.7% of participants in various social gathering places on the Island of Montreal. However, this required adequate financial and logistical resources.

As shown by the 81.8% participation rate, the team was able to access people from both communities during promotion and recruitment activities. The main reasons for refusal were related to DBS sampling, mistrust and the feeling of stigmatization experienced by some participants, especially in the English-speaking Caribbean community. Also, a connection was established between the screening tests involved in the process of immigrating to Canada and the need to get retested.

Compared to sub-Saharan African communities, English-speaking Caribbean communities consisted of newcomers and people who had been in Canada for many generations. As a result, the first generation of some subgroups is aging (age >49 years), a situation that needs to be taken into account in subsequent phases of the second generation E-Track surveillance system in Montreal. It should also be noted that approximately 23% (111/484) of newly diagnosed cases of HIV in individuals from an endemic country occurred in those 50 years of age and older since the HIV infection surveillance program was introduced in Quebec, i.e. from April 2002 to December 2013.⁵ Future research should compare HIV risk factors between first and second generations of people from HIV-endemic countries to better target at-risk subgroups and prevent the acquisition and transmission of new infections.

Regarding the methodological limitations of the SGS-AFCAR Survey, note that the data were collected through the voluntary participation of members of both communities living on the Island of Montreal. We believe they were no different from those who decided not to participate. The sample was constructed based on a rational choice to make it reflect the target population (non-probability sampling). The surveillance results are therefore not representative of the entire target population in Quebec. In addition, results presented in the report are based on self-reported data, which are subject to memory and social desirability biases. Socially undesirable behaviours may therefore have been under-reported, and socially desirable behaviours over-reported. However, compliance with recruitment and questionnaire administration guidelines (self-administered or not), and monitoring of the quality of the interviewers' work allowed us to minimize certain biases. Lastly, the methods chosen, which are reproducible and easy to implement, produced high-quality data.

The data collected should serve as a benchmark for making comparisons over time and performing more complex analyses in the future, which will focus on risk factors associated with HIV, screening, sexual risk behaviours in each community and the vulnerability of women. These results will provide a better understanding of target STBBIs and factors that make people from HIV-endemic countries vulnerable.

Other research initiatives (qualitative and quantitative) could focus on the way they perceive the risk of infection and how they adapt to it. Questionnaires should therefore provide a better characterization of sexual risk behaviours (anal, vaginal, oral) with or without condom use based on the type of sexual partner.

5. SUMMARY OF RESULTS, RISK PROFILE AND POTENTIAL INTERVENTIONS

Based on the results, this chapter outlines an HIV infection risk profile for each of the two target communities.

5.1. Vulnerabilities common to both communities

Based on this second generation HIV surveillance survey, certain subgroups in both communities appeared more vulnerable to HIV/AIDS. These subgroups should be priority targets for measures promoting HIV/AIDS prevention and protection.

5.1.1. Level of knowledge of HIV status

It is disturbing to note that 68.2% (15/22) of participants infected with HIV did not know they were HIV positive. This percentage is about 25% in Canada.⁴ In total, 80.4% (866/1,077) of sub-Saharan Africans vs. 61.3% (302/493) of English-speaking Caribbeans had already undergone HIV testing, with just over half having undergone testing in the previous two years [67.8% of sub-Saharan Africans and 68.8% of English-speaking Caribbeans]. These results suggest that screening programs do not reach individuals most likely to be infected with HIV. This hypothesis is supported by the fact that the INSPQ's HIV infection surveillance program reported that in 2013, 18.6% (41/220) of new cases diagnosed among heterosexuals from HIV-endemic countries had never before been tested for HIV.⁵

5.1.2. Sexual risk behaviours

In the past year, the majority of sexually active 15–49-year-olds [84.5% (1,146/1,357)] reported having had vaginal or anal sex with penetration, and over a third of them [36.7% (416/1,132)] had multiple sexual partners (≥ 2). During the same period, 33.0% (252/764) of sub-Saharan Africans vs. 46.7% (177/379) of English-speaking Caribbeans had at least one sexual partner with whom they had had vaginal or anal sex with penetration, and overall 43.4% (185/426) did not always use a condom (male or female) with these partners. This applied to significantly more sub-Saharan African women than men (51.2% vs. 35.1%; $p=0.048$).

In addition, 40.0% (14/35) of sub-Saharan African women vs. 19.1% (49/209) of men ($p=0.006$), as well as 45.3% (24/53) of English-speaking Caribbean women vs. 23.1% (28/121) of men ($p=0.003$) had not used a condom the last time they had vaginal or anal sex with penetration with their casual sex partners. Some of them had regular concomitant sexual partners and lived either as a couple with their partner or separately. The frequency of high-risk vaginal or anal sex with penetration was therefore significant in both communities. These risk behaviours have a significant impact on the spread of the HIV epidemic, as people affected by these practices are at greater risk of acquiring or transmitting HIV sexually.²⁷

5.1.3. Vulnerability of women

The survey results highlight women's vulnerability to HIV infection. A greater number of women in the two communities did not use a condom (male or female) the last time they had vaginal or anal sex with penetration with a casual sex partner, and did not use a condom during a sexual encounter when travelling in a country where HIV is highly endemic.

It has been shown in other contexts that women's greater vulnerability to HIV is, among other things, due to social, cultural and economic pressures that do not allow them to take preventive measures.²⁸ This vulnerability was more pronounced in the sub-Saharan African community.

5.1.4. 30–49 age group

The majority of HIV-positive tests occurred in the 40–49 age group [34.9% (12/22)], followed by the 30–39 age group [31.8% (7/22)]. All HIV-positive women were in these two age groups. Age was significantly associated with HIV [OR=1.10; 95% CI: 1.05–1.16; $p<0.0001$]. Compared to the 15–29 reference age group, the 40–49 age group was most at risk for HIV.

5.1.5. Risk taking while travelling in a country where HIV is highly endemic

Slightly more than half [56.4% (101/179)] of participants who had vaginal or anal sex with penetration during a trip in a country where HIV is highly endemic, with a sexual partner living there, did not always use a condom (male or female) during sex. A greater percentage of women from both communities were in this situation.

This result had been previously observed in members of the Haitian community living in Montreal.⁸⁻¹⁰

These surveillance data do not explain the reason(s) for this type of risk-taking while travelling. But, the fact that it occurs in countries where HIV is highly endemic clearly indicates that preventive measures must be taken to protect travellers.

5.1.6. More limited access to primary care

In total, 60.7% (963/1,587) of participants recruited in the two communities did not have a family doctor [67.1% of sub-Saharan Africans vs. 46.5% of English-speaking Caribbeans ($p<0.0001$)], whereas the 2012 TOPO survey reported that 35% of all Montrealers did not have a family doctor.²⁹

Among all participants who required medical care in the 12 previous months, 15.1% (223/1,473) reported having difficulty accessing the health care services they needed during this period [15.7% of Sub-Saharan Africans and 10.1% of English-speaking Caribbeans].

Significant differences observed between the living conditions of Montrealers and survey participants, particularly in terms of income, housing, employment, education, nutrition and access to health care, have a direct effect on their health and life expectancy.²⁹ Factors associated with social inequalities in health should be countered by effective preventive measures to ensure living conditions that promote the health and development of target communities.

5.2. Communities from sub-Saharan Africa

5.2.1. High prevalence of HIV infection

HIV prevalence in the community was 1.7% [1.0–2.6%]. It is comparable to that among Quebecers of Haitian origin in 1994: 1.3% [1.0–1.6%].⁸ In Canada, estimated HIV prevalence was 0.21% [0.17–0.24%] in 2011.⁴

The high prevalence of HIV infection among Montrealers from sub-Saharan Africa clearly highlights the need for rigorous, effective programs. This include measures to encourage a greater percentage of sexually active individuals to get tested for HIV when they change or acquire new sexual partners, or return to a previous sexual partner.³⁰

5.2.2. African women and heterosexual transmission of HIV

In Quebec and elsewhere in Canada, the fact that the majority of heterosexual transmission of HIV infection occurs in people from sub-Saharan Africa and the Caribbean, as well as the preponderance of infection among women, are constants. In the SGS-AFCAR Survey, all HIV positive cases (n=22) occurred among heterosexuals, and just over half the cases [54.5% (12/22)] occurred in women.

According to the SGS-AFCAR Survey, sub-Saharan African women had the highest HIV prevalence at 2.4% [1.2–4.2%], compared to men in the same community (1.2% [0.6–2.4%]). In contrast, among Quebecers of Haitian origin, HIV prevalence was higher in men than women: 1.6% [1.1–2.2%] vs. 1.1% [0.7–1.5%].¹⁰

Sub-Saharan African (and Caribbean) women are at high risk for HIV/AIDS because of their socioeconomic status and their attitudes about condom use, and because some of them and some of their partners have multiple partners.¹⁰ Other studies have highlighted their biological susceptibility to HIV/AIDS² and their inability to require that their partner use condoms.

A better understanding of the decision-making dynamics between men and women, specifically the role of African women in deciding to use or oblige their sexual partners (especially casual) to use condoms, is required to counter heterosexual transmission of HIV in this high-risk subgroup.

Also, concrete steps must be taken to empower African women and reinforce their preventive behaviours by emphasizing the meaning of risk taking, and promoting condom use and routine screening for STBBIs.

5.3. Communities from the English-speaking Caribbean

5.3.1. Sexual risk behaviours for HIV infection

A higher frequency of sexual risk behaviours was observed among members of this community. They had a higher number of casual sex partners and did not always use a condom (male or female) while having vaginal or anal sex with penetration, when travelling in HIV-endemic countries. These are all areas on which stakeholders working with this community should focus.

Also, the average age at first intercourse was earlier among English-speaking Caribbeans (15.8 years; SD: ± 3.8) compared to sub-Saharan Africans (18.4 years; SD: ± 3.9), who were closer to the Quebec average, which according to population survey results appears stable at around 18 years.²⁶ One-third of young women and men aged 15 to 24 had engaged in sexual intercourse before age 15. This highlights the need to increase prevention and educate young people very early. Some evidence suggests that when the first sexual encounter occurs at a later age, the risk of infection through sexual intercourse is reduced, at least among women.²⁷ Preventive measures should be developed to facilitate dialogue and communication between young people and their parents or guardians in order to promote healthy sexual attitudes and prevent STBIs.³⁰

5.3.2. Level of education

The percentage of English-speaking Caribbean men who had a high school education or higher was lower than among women from the same community. In total, 37.9% (188/496) of English-speaking Caribbeans [29.9% (69/231) of men vs. 44.9% (119/265) of women; $p=0.001$] compared to 70.1% (771/1,100) of sub-Saharan Africans [73.2% (485/663) of men vs. 65.4% (286/437) of women; $p=0.006$] involved in the SGS-AFCAR Survey had a high school education or higher.

Among other things, the level of education influences health because it enhances people's ability to learn and understand information that can improve their health.² This underscores the need to focus on subgroups with low levels of literacy through adapted community programs.

5.3.3. Knowledge about HIV transmission modes

Compared to sub-Saharan Africans, English-speaking Caribbeans knew less about HIV transmission modes and prevention factors. Also, 8.1% of English-speaking Caribbeans vs. 5.1% of sub-Saharan Africans believed that someone who looked healthy could not be HIV positive. This is a common misconception that can lead to unprotected sex with infected sexual partners.²⁷

Also, 44.6% of English-speaking Caribbeans vs. 25.0% of sub-Saharan Africans believed that HIV is transmitted through mosquito bites or did not know the answer to that question. This belief could reduce a person's willingness to adopt risk-free sexual behaviours.²⁷

Lastly, 33.9% of English-speaking Caribbeans vs. 10.4% of sub-Saharan Africans believed that the virus can be transmitted by sharing a meal with an infected person or did not know the answer to that question. This belief would further stigmatize people living with AIDS.²⁷

The level of education appears to correlate with the level of knowledge about modes of HIV transmission. It is as important to correct major misconceptions about HIV transmission as it is to provide accurate information on actual modes of HIV transmission.²⁷ The gaps identified in knowledge about HIV transmission modes need to be addressed, taking into account the social and cultural diversity of target populations.

5.3.4. Prenatal screening

Only 83.5% (111/133) of English-speaking Caribbean women were tested for HIV during their last pregnancy. All kinds of barriers can hinder this population's access to screening. The benefits of early detection for mother and child are becoming increasingly obvious. This makes it all the more important to identify these barriers and create partnerships in the community that promote the development of measures to eliminate them.

5.3.5. Substance use during sexual intercourse

Higher percentages of English-speaking Caribbeans than sub-Saharan Africans and significantly more men than women in the two communities had used alcohol [51.6% (195/378) of English-speaking Caribbeans vs. 19.7% (150/762) of sub-Saharan Africans], marijuana or other drugs while having vaginal or anal sex with penetration in the last 12 months.

Holistic, non-stigmatizing and non-judgemental steps should be taken to prevent HIV among these individuals.³⁰

6. OTHER INTERVENTIONS AND RECOMMENDATIONS

Possible approaches had been identified previously to help improve or develop HIV/AIDS prevention and protection strategies for the most vulnerable subgroups. Also, as part of the regional STBBI prevention program, professionals and decision-makers involved at various levels should take into account the main findings of the SGS-AFCAR Survey and implement these potential approaches, based on the target communities' specific context and environment.

A feasibility study²¹ was conducted to identify factors that facilitated or hindered implementation of second generation HIV surveillance survey in Montreal communities from sub-Saharan Africa and the English-speaking Caribbean, and effective implementation of the SGS-AFCAR Survey. The study provided us with a better understanding of the situation in these communities and their STBBI prevention needs. The following potential approaches were not based on the quantitative results of the SGS-AFCAR Survey. They were developed through discussions, consultations and interactions we had with the Community Advisory Committee, organizations and some community and religious leaders, and members of both communities before the start of the survey, but also through observations made while the participants were recruited.

6.1. Reality and needs of the communities involved

Compared to Montreal's Haitian community, there are very few African and English-speaking Caribbean community organizations working on preventing STBBIs and more generally on other health issues in Montreal.

The community participation approach that helped the target communities to become involved in all stages of the development and implementation of the SGS-AFCAR Survey should be bolstered. This would encourage the target communities to use the survey results to mount a timely, effective HIV/AIDS prevention campaign.

These communities would have to get organized to create opportunities and mechanisms to facilitate contacts and cooperation between members in response to HIV/AIDS-related issues. They will have to work with not only community leaders, religious institutions such as churches and mosques, and groups of nationals from the same country, but also with PHA from these communities in order to develop a community response that recognizes diversity within the African and English-speaking Caribbean communities.

These emerging Montreal community leadership would identify channels for promoting HIV/AIDS awareness in order to minimize further stigmatization, improve communications and cooperation (community media, bilingual brochures, etc.), and follow up on the recommendations in this report.

6.2. Commitment and awareness of health professionals and managers

Our colleagues in the community highlighted the need to make health professionals and managers aware of the specific issues involved in providing medical attention for members of ethno-cultural communities (different beliefs regarding HIV and other STBBIs, etc.), and the difficulties ethno-cultural populations are having with access to the health care system. The potential solutions they recommend include promoting local services so that everyone has access to quality health care, and involving ethno-cultural communities in response program development, implementation and delivery. They also suggest that partnerships be established and that we collaborate with community organizations working in the field of HIV/AIDS (COCQSIDA, etc.) and other partners (community organizations that help new immigrant and refugee families, agencies such as food banks that provide support in times of crisis, organizations that offer French-language and literacy courses, etc.). To improve their ethno-cultural awareness and competence and ensure the continuity of STBBI care and programs, these stakeholders should use approaches that take into account the broader social context that influences the health of individuals and communities, that is, culture, gender, socioeconomic status, social support networks, and social environments.³⁰

6.3. Political responsibility

Concrete steps must be taken in Montreal because people from HIV-endemic countries are over-represented in the heterosexual HIV transmission category, the SGS-AFCAR Survey has indicated that the prevalence rate in both target communities is high, and certain subgroups including sub-Saharan African women are vulnerable. HIV/AIDS response should therefore be a funding priority in ethno cultural communities. Funding is needed to increase the number of programs, services and service providers, and to enhance the availability, accessibility and quality of HIV/AIDS programs and services.³¹ This response should include the prevention of HIV and other STBBIs, and other issues affecting ethno-cultural communities, such as immigration, poverty and housing.³¹ It could also adapt the ministerial guidelines outlined in the document entitled “Une passerelle vers un avenir en santé” [Path to a healthy future]³² to Montreal realities.

CONCLUSION

People from sub-Saharan Africa and the Caribbean are more vulnerable to HIV/AIDS and should receive special attention in the fight against HIV/AIDS. In Quebec and elsewhere in Canada, these subgroups have seldom been the focus of HIV/AIDS research programs and prevention policies. In the past, little information was available to develop appropriate measures to help them. The SGS-AFCAR Survey helped meet this need.

Second generation HIV surveillance surveys can therefore be implemented for people from HIV-endemic countries. The SGS-AFCAR Survey should be resumed to analyze trends and improve the targets of culturally appropriate strategies for enhancing the level of HIV/AIDS knowledge, reducing risk behaviours and improving access to screening for people from HIV-endemic countries. These data will also be used to evaluate response programs.

The SGS-AFCAR Survey results confirmed that HIV prevalence remains high among people from HIV-endemic countries living on the Island of Montreal. High-risk sexual practices were highlighted and the risk of transmitting HIV and other STBBIs persists in the most vulnerable subgroups, namely women. Swift action on these results needs to be taken to develop and improve policies, programs, interventions and services at the community, local, provincial and national levels for English- or French-speaking sub-Saharan African and Caribbean populations in Quebec or elsewhere in Canada.

REFERENCES

- 1 Public Health Agency of Canada (2013). HIV and AIDS in Canada: Surveillance Report to December 31, 2012. Surveillance and Epidemiology Division, Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada, 2013. Available at: http://publications.gc.ca/collections/collection_2014/aspc-phac/HP37-2-2012-eng.pdf [Accessed February 2, 2014]
- 2 Public Health Agency of Canada (2009). Population-Specific HIV/AIDS Status Report: People from Countries where HIV is Endemic – Black people of African and Caribbean descent living in Canada. Available at: <http://www.phac-aspc.gc.ca/aids-sida/publication/ps-pd/africacaribbe/index-eng.php> [Accessed March 12, 2013]
- 3 UNAIDS (2014). The Gap report: UNAIDS report on the global AIDS epidemic 2013. Available at: http://www.unaids.org/en/media/unaids/contentassets/documents/unaidspublication/2014/UNAIDS_Gap_report_en.pdf [Accessed July 19, 2014]
- 4 Public Health Agency of Canada (2012). Summary: Estimates of HIV Prevalence and Incidence in Canada, 2011. Centre for Communicable Diseases and Infection Control, 2012. ISSN: 2291-045X. Available at: <http://www.phac-aspc.gc.ca/aids-sida/publication/survreport/assets/pdf/estimat2011-eng.pdf> (Accessed April 28, 2014).
- 5 Bitera R. (2014). Programme de surveillance de l'infection par le virus de l'immunodéficience humaine (VIH) au Québec : cas cumulatifs 2002-2013. (Rapport préliminaire), Québec, Institut national de santé publique du Québec.
- 6 Ministère de la santé et des services sociaux (2013). Surveillance du sida au Québec. Publication no.: 12-268-09W. Available at: <http://publications.msss.gouv.qc.ca/acrobat/f/documentation/preventioncontrole/sidapage2.pdf> [Accessed May 20, 2014].
- 7 Ogunnaiké-Cooke S, et al. (2013). A Framework for the E-Track: An enhanced surveillance system designed to monitor trends in the prevalence of HIV and associated factors and infections among people in Canada from countries where HIV is endemic. Poster presentation at the 22nd Annual Canadian Conference on HIV/AIDS Research, Vancouver, April 11–14, 2013 (P131). <http://www.cahr-acrv.ca/content/uploads/2012/10/CAHR-2013-Abstract-Book-FINAL.pdf>
- 8 Adrien A, Leane V, Remis RS, et al. (1999). Migration and HIV: an epidemiological study of Montrealers of Haitian origin. *Int J STD AIDS* 1999; 10(4):237–42.
- 9 Joseph G, Adrien A, Tremblay C, Leclerc P (2010). Les infections transmises sexuellement et par le sang chez les Québécois d'origine haïtienne / Surveillance biologique et comportementale – 2e édition Available at: <http://collections.banq.qc.ca/ark:/52327/bs1990144> [Accessed June 12, 2013].
- 10 Adrien A, Cox J, Leclerc P, et al. (2010). Behavioral risks for HIV infection among Quebec residents of Haitian origin. *J Immigr Minor Health* 2010; 12:894–9.

- 11 Public Health Agency of Canada (2014). Questions and answers: Inclusive practice in the prevention of sexually transmitted and blood borne infections among ethnocultural minorities. Available at: <http://www.catie.ca/sites/default/files/Inclusive-practice-in-the-prevention-of-STBBIs-among-ethnocultural-minorities.pdf> [Accessed June 16, 2014]
- 12 Yang Q, Boulos D, Yan P, et al. (2010). Estimates of the number of prevalent and incident human immunodeficiency virus (HIV) infections in Canada, 2008. *Can J Public Health*, 101(6):486–
- 13 Brenner BG, Roger M, Routy JP, et al. (2007). High rates of forward transmission events after acute/early HIV-1 infection. *J Infect Dis* 195:951–9.
- 14 Pillay D, Fisher M (2007). Primary HIV Infection, phylogenetics, and antiretroviral prevention. *J Infect Dis* 195:924–6.
- 15 UNAIDS (2013). Global Report: UNAIDS report on the global AIDS epidemic 2013. Available at: http://www.unaids.org/en/media/unaids/contentassets/documents/epidemiology/2013/gr2013/UNAIDS_Global_Report_2013_en.pdf [Accessed January 11, 2014]
- 16 UNAIDS (2014). Issues Brief: Local Epidemics. Available at: http://www.unaids.org/en/media/unaids/contentassets/documents/unaidspublication/2014/JC2559_local-epidemics_en.pdf [Accessed July 6, 2014]
- 17 Institut national de santé publique du Québec (2014). Nombre de cas déclarés pour Sida, pour la Province de Québec. Available at: <https://www.extranetmado.ca/mado04/MADO/asp/rapportmadopublic.asp?regso=0&grp=16>
- 18 Fleury E, et al. (2014). Guide québécois de dépistage des infections transmissibles sexuellement et par le sang, Québec, Service de lutte contre les ITSS Direction générale de la santé publique, Ministère de la santé et des services sociaux. 239 pp. Available at: <http://publications.msss.gouv.qc.ca/acrobat/f/documentation/2014/14-308-01W.pdf>
- 19 Ministère de la santé et des services sociaux (2013). Formulaire SP-100. Available at: <http://publications.msss.gouv.qc.ca/acrobat/f/documentation/preventioncontrole/SP-100.pdf> [Accessed May 20, 2014]
- 20 World Health Organization, UNAIDS (2000). Second generation surveillance for HIV: the next decade. Geneva, World Health Organization. 40 pp (WHO/CDS/CSR/EDC/2000.5). Available at: http://whqlibdoc.who.int/hq/2000/WHO_CDS_CSR_EDC_2000.5.pdf?ua=1
- 21 Butoyi J, Hot A, Bernier M, Adrien A, Ogunnaike-Cooke S, et al. (2012). Faisabilité d'un système de surveillance de deuxième génération du VIH auprès des personnes originaires de pays où le VIH est endémique. Poster presentation at the 21st Annual Canadian Conference on HIV/AIDS Research, Montreal, April 2012 (P244).
- 22 Ministère de l'Immigration et des Communautés culturelles (MICC) 2010. Portrait statistique de la population recensée au Québec en 2006. Available at: <http://www.micc.gouv.qc.ca/fr/publications/etudes-recherches-statistiques/statistiques-population/recensement-2006.html> [Accessed March 25, 2012]
- 23 INSPQ (2003). Secrétariat du Comité d'éthique de santé publique. La surveillance continue de l'état de santé de la population une synthèse; pp. 4–5.

-
- 24 Unité d'éthique de la recherche du MSSS (2007). Note de clarification relative aux compétences matérielles et territoriales des comités d'éthique de la recherche.
- 25 Comité d'éthique de santé publique (2012). Avis sur le Plan régional de surveillance de la région de Montréal 2011-2015.
- 26 Blais M, et al. (2009). La sexualité des jeunes Québécois et Canadiens : Regard critique sur le concept l'hypersexualisation. *Globe : Revue internationale d'études québécoises*, vol. 12, no. 2, pp. 27–8.
- 27 UNAIDS (2014). Global Report: Global AIDS Response Progress Reporting 2014. Construction of Core Indicators for monitoring the 2011 United Nations Political Declaration on HIV and AIDS. Available at: http://www.unaids.org/en/media/unaids/contentassets/documents/document/2014/GARPR_2014_guidelines_en.pdf [Accessed May 11, 2014]
- 28 Criton C, Fener P (2007). Facteurs de risque de l'infection à VIH/sida chez la femme. Centre National de la Recherche Scientifique (CNRS), France.
- 29 Direction de santé publique de l'Agence de la santé et des services sociaux de Montréal (2014). Enquête TOPO sur les maladies chroniques et leurs déterminants, 2012. Available at: http://emis.santemontreal.qc.ca/fileadmin/emis/Sant%C3%A9_des_Montr%C3%A9alais/%C3%89tat_de_sant%C3%A9/sant%C3%A9_physique/TOPO_synthese_infographique/TOPO-synthese-MTL-201405-fr.pdf [Accessed June 15, 2014]
- 30 LLana J (2006). HIV Prevention Guidelines and Manual: A Tool for Service Providers serving African and African Caribbean Communities in Canada. African and Caribbean Council on HIV/AIDS in Ontario (ACCHO) and Women's Health in Women's Hands Community Health Centre (WHIWH). Toronto, Canada. ISBN 0-9736431-3-9.(PDF).53 pp. Available at: http://www.accho.ca/pdf/hiv_manual-part_1_english.pdf
- 31 DA Falconer & Associates Inc. (2008). Taking Action on HIV and AIDS in Black Communities in Canada: A Resource for Moving Ahead. Ottawa, Interagency Coalition on AIDS and Development. Available at: http://icad-cisd.com/pdf/Taking_Action_EN.pdf [Accessed June 2014].
- 32 Ministère de la santé et des services sociaux (2012). Une passerelle vers un avenir en santé. La planification et l'organisation des services à l'intention des réfugiés et des demandeurs d'asile : évaluer le bien-être et l'état de santé physique pour mieux desservir : orientations ministérielles. La Direction des communications du ministère de la Santé et des Services sociaux. ISBN: 978-2-550-63958-9 (version PDF). 27 pp. Available at: <http://publications.msss.gouv.qc.ca/acrobat/f/documentation/2011/11-522-01W.pdf> [Accessed August 2014]

APPENDICES

Appendix 1: List of HIV-endemic countries

Africa		Caribbean, Bermuda and Central America / South America
▪ Angola ▪ Benin ▪ Botswana ▪ Burkina Faso ▪ Burundi ▪ Cameroon ▪ Cape Verde ▪ Central African Republic ▪ Chad ▪ Democratic Republic of the Congo (formerly Zaire)–Kinshasa ▪ Republic of the Congo–Brazzaville ▪ Djibouti ▪ Equatorial Guinea ▪ Eritrea ▪ Ethiopia ▪ Gabon ▪ Gambia ▪ Ghana ▪ Guinea ▪ Guinea-Bissau ▪ Côte-d'Ivoire ▪ Kenya	▪ Lesotho ▪ Liberia ▪ Malawi ▪ Mali ▪ Mozambique ▪ Namibia ▪ Niger ▪ Nigeria ▪ Rwanda ▪ Senegal ▪ Sierra Leone ▪ Somalia ▪ South Africa ▪ Sudan ▪ South Sudan ▪ Swaziland ▪ Tanzania ▪ Togo ▪ Uganda ▪ Zambia ▪ Zimbabwe	▪ Anguilla ▪ Antigua and Barbuda ▪ Bahamas ▪ Barbados ▪ Bermuda ▪ British Virgin Islands ▪ Cayman Islands ▪ Dominica ▪ Dominican Republic ▪ French Guiana ▪ Grenada ▪ Guadeloupe ▪ Guyana ▪ Haiti[§] ▪ Honduras ▪ Jamaica ▪ Martinique ▪ Montserrat ▪ Netherlands Antilles ▪ St. Lucia ▪ Saint Kitts and Nevis ▪ Saint Vincent and the Grenadines ▪ Surinam ▪ Trinidad and Tobago ▪ Turks and Caicos Islands ▪ US Virgin Islands

[§] Montrealers of Haitian descent were excluded from the SGS-AFCAR Survey.

Appendix 2: E-Track Questionnaire

E-Track

Effectuer le suivi des tendances en matière de
prévalence du VIH et des facteurs comportementaux
et sociodémographiques connexes chez les Canadiens
nés dans des pays où le VIH est endémique

PHASE 1 – Montreal 2013

**CONFIDENTIAL
QUESTIONNAIRE**

Administrative questions (to be completed by the interviewer)

Eligibility screening

Assess and check: (both statements must be checked to continue)

Have you checked whether the candidate is eligible?

☐ Participant is eligible

Did the participant give consent to participate?

☐ Yes

Did the participant consent to the storage and additional testing of their finger-prick blood sample in the future?

☐ Yes

☐ No, DBS sample to be destroyed after testing

MTL(1). Recruitment organization: _____

1. Write in the last four digits of the Site ID Code:

E-13-MTL-_____

This survey is confidential and anonymous. Your name will not be put on the questionnaire. I will ask you to enter your initials and date of birth into a special computer program which will make up a unique scrambled code. After the scrambled code is generated and recorded on the questionnaire, your date of birth and initials will be deleted immediately from the computer program.

2. Enter the Encrypt ID code (always 10 characters long):



3. Select the name of the city:

☐ Montreal

4. Please enter the site code of the location where the interview took place::

Site code: _____

5. Enter today's date using the format: DD/MM/YYYY (e.g., 15/04/2013):

_____/_____/_____
DD MM YYYY

6. Enter your initials (from 2 to 5 letters): _____

7. Enter the interview start time. Please indicate in 24-hour time. For example, enter 16:15 for 4:15 p.m.

We are now ready to start the questionnaire portion of this interview.

)

How did you hear about the E-Track survey? *(Check ALL that apply)*

- ☐ Word of mouth
- ☐ Poster/flyer
- ☐ Recommendation
- ☐ When you arrived
- ☐ Media/event
- ☐ Social media (e.g., Facebook, Twitter) (specify): _____
- ☐ Community organizations (specify): _____
- ☐ Other (specify): _____

Your answers will now be put into the questionnaire. Would you like to complete the questionnaire yourself, or would you like me to read the questions and fill in the answers that you provide? Either way, I will stay in the room with you until the questionnaire is completed and I am happy to help you with any questions that you may have.

Are you ready to start?

SECTION ONE: Socio-demographics

1.1 How old are you now?

Number of years (specify): _____

If the participant is less than 15 years of age, end interview here.

1.2 Do you live in Montréal right now?

- ☐ Yes
☐ No

- ☐ Don't know
☐ Prefer not to answer

If the answer is No, Don't know, or the person Prefer not to answer, end the interview.

MTL1 (1.2) What are the first three characters of your postal code? If you live in a shelter, enter the first three characters of the shelter's postal code. Please note that the three characters of your postal code cannot be used to identify your residence.

- ☐ *Don't know*
☐ *Prefer not to answer*

MTL2 (1.2) What is your CLSC (local community service centre)? (See the list of CLSCs to enter your CLSC's code)

- ☐ *Don't know*
☐ *Prefer not to answer*

1.3 What is your country of birth? (Select only ONE)

Africa		Caribbean, Bermuda and Central America / South America
☐ Angola ☐ Benin ☐ Botswana ☐ Burkina Faso ☐ Burundi ☐ Cameroon ☐ Cape Verde ☐ Central African Republic ☐ Chad ☐ Democratic Republic of the Congo (formerly Zaire)–Kinshasa ☐ Republic of the Congo–Brazzaville ☐ Djibouti ☐ Equatorial Guinea ☐ Eritrea ☐ Ethiopia ☐ Gabon ☐ Gambia ☐ Ghana ☐ Guinea ☐ Guinea-Bissau ☐ Côte-d'Ivoire ☐ Kenya	☐ Lesotho ☐ Liberia ☐ Malawi ☐ Mali ☐ Mozambique ☐ Namibia ☐ Niger ☐ Nigeria ☐ Rwanda ☐ Senegal ☐ Sierra Leone ☐ Somalia ☐ South Africa ☐ Sudan ☐ South Sudan ☐ Swaziland ☐ Tanzania ☐ Togo ☐ Uganda ☐ Zambia ☐ Zimbabwe	☐ Anguilla ☐ Antigua and Barbuda ☐ Bahamas ☐ Barbados ☐ Bermuda ☐ British Virgin Islands ☐ Cayman Islands ☐ Dominica ☐ Dominican Republic ☐ French Guiana ☐ Grenada ☐ Guadeloupe ☐ Guyana ☐ Honduras ☐ Jamaica ☐ Martinique ☐ Montserrat ☐ Netherlands Antilles ☐ St. Lucia ☐ Saint Kitts and Nevis ☐ Saint Vincent and the Grenadines ☐ Surinam ☐ Trinidad and Tobago ☐ Turks and Caicos Islands ☐ US Virgin Islands

☐ Other (specify): _____

☐ Don't know

☐ Prefer not to answer

} If the answer is Other, Don't know, or the person Prefer not to answer, end the interview.

1.4 What year did you come to live in Canada?

Year (specify): _____

- ☐ *Don't know*
☐ *Prefer not to answer*

1.5 What is your residency status in Canada? (Check ALL answers that apply)

- ☐ Canadian citizen
☐ Landed immigrant / permanent resident
☐ Refugee / protected person (means you have been officially accepted as a refugee)
☐ Refugee claimant / person in need of protection (means you have applied for refugee protection status, and are waiting for a decision on your claim)
☐ Temporary work permit (temporary foreign worker)
☐ Temporary resident visa (visitor)
☐ Study permit (international student)
☐ Parent and grandparent super visa
☐ Other (specify): _____
- ☐ *Don't know*
☐ *Prefer not to answer*

1.6 In which country did you spend most of your childhood (0 to 18 years old)?

- ☐ Your country of birth
☐ Other country (specify): _____
- ☐ *Don't know*
☐ *Prefer not to answer*

1.7 People living in Canada are of very diverse cultural and racial backgrounds. Do you think you belong to one of the following groups? (Check ALL answers that apply up to a MAXIMUM of four)

- ☐ Black
- ☐ White
- ☐ South Asian
- ☐ Far East Asian
- ☐ Southeast Asian
- ☐ Arab or West Asian
- ☐ Latin American / Hispanic
- ☐ Aboriginal
- ☐ Other (specify): _____

- ☐ Don't know
- ☐ Prefer not to answer

1.8 What was your sex at birth? (Check ONE answer only)

- ☐ Man
- ☐ Woman
- ☐ Other (specify): _____

- ☐ Don't know
- ☐ Prefer not to answer

1.9 Today you consider yourself a? (Check ONE answer only)

- ☐ Man
- ☐ Woman
- ☐ Transsexual or transgender
- ☐ Other (specify): _____

- ☐ Don't know
- ☐ Prefer not to answer

1.10 1.10 Which of the following best describes your sexual orientation? (Check ONE only)

- ☐ Heterosexual or Straight
☐ Homosexual or Gay or Lesbian
☐ Bisexual
☐ Questioning
☐ Other (specify): _____

☐ *Don't know*
☐ *Prefer not to answer*

1.11 What is your CURRENT relationship or marital status? (Check all that apply to current relationship or marital status)

- ☐ Single (you are not in a relationship and have never been married)
☐ In a relationship with a man, but you are not living together and are not married.
☐ In a relationship with a woman, but you are not living together and are not married.
☐ In a relationship with a man; you are living together as a couple, but are not legally married.
☐ In a relationship with a woman; you are living together as a couple, but are not legally married.
☐ Married and living with my husband.
☐ Married and living with my wife.
☐ Married, but my husband does not live in Canada.
☐ Married, but my wife does not live in Canada.
☐ Married, but separated from my husband.
☐ Married, but separated from my wife.
☐ Divorced.
☐ Widower/Widow
☐ Other (specify): _____

☐ *Don't know*
☐ *Prefer not to answer*

1.12 What are your beliefs or religion, if any? (Check ONE answer only)

- ☐ Sunni Muslim
- ☐ Shiite Muslim
- ☐ Catholic
- ☐ Anglican
- ☐ Lutheran
- ☐ Baptist
- ☐ Seventh-day Adventist
- ☐ Jehovah's Witness
- ☐ Member of the Evangelical Church
- ☐ Member of a traditional African religion
- ☐ Hindu
- ☐ None or Atheist or Atheist
- ☐ Other (specify): _____

- ☐ *Don't know*
- ☐ *Prefer not to answer*

1.13 What is the highest level of education or training that you have completed? (Check ONE only)

- ☐ None
- ☐ Some elementary school
- ☐ Completed elementary school
- ☐ Some high school
- ☐ Completed high school
- ☐ Some college, CEGEP, vocational school, trade school, or apprenticeship training
- ☐ Completed college, CEGEP, vocational school, trade school, or apprenticeship training
- ☐ Some university
- ☐ Completed university certificate or diploma
- ☐ Completed undergraduate university degree (for example B.A., B.Sc., B.Ed.)
- ☐ Completed graduate university degree (for example M.A., M.Sc., PhD, D.Sc., M.D., J.D)
- ☐ Other (specify): _____

- ☐ *Don't know*
- ☐ *Prefer not to answer*

1.14 What was your employment status in the last 12 months? (Check ALL answers that apply)

- ☐ Employed full time (including self-employment)
- ☐ Employed part time (including self-employment)
- ☐ Full-time student
- ☐ Part-time student
- ☐ Unpaid work at home (doing housework or caring for family members)
- ☐ Volunteering or providing care (for people other than members of your family)
- ☐ Retired
- ☐ Received disability benefits
- ☐ Received social security benefits
- ☐ Unemployed
- ☐ Other (specify): _____
- ☐ Don't know
- ☐ Prefer not to answer

1.15 In the last 12 months, what was your approximate total household income? That is to say, what was the total income of all household members, from all sources before taxes and other deductions? (Check ONE answer only)

- ☐ Less than \$5,000
- ☐ \$5,000 or more but less than \$10,000
- ☐ \$10,000 or more but less than \$15,000
- ☐ \$15,000 or more but less than \$20,000
- ☐ \$20,000 or more but less than \$30,000
- ☐ \$30,000 or more but less than \$40,000
- ☐ \$40,000 or more but less than \$50,000
- ☐ \$50,000 or more but less than \$60,000
- ☐ \$60,000 or more but less than \$70,000
- ☐ \$70,000 or more but less than \$80,000
- ☐ \$80,000 or more but less than \$90,000
- ☐ \$90,000 or more but less than \$100,000
- ☐ \$100,000 or more but less than \$150,000
- ☐ \$150,000 and over

- ☐ Don't know
- ☐ Prefer not to answer

} If the person replied Don't know or Prefer not to answer, go to question 1.19.

1.16 How many people people living in Canada (including yourself) depend on your household income?**Number of individuals in Canada:** Specify: _____

- ☐ *Don't know*
☐ *Prefer not to answer*

1.17 How many people living outside Canada depend on your household income?**Number of individuals outside of Canada:** Specify: _____

- ☐ *Don't know*
☐ *Prefer not to answer*

1.18 In the past 12 months, considering your household income, how easy or difficult was it for you to meet basic food and shelter costs? (Check ONE only)

- ☐ Very easy
☐ Somewhat easy
☐ Somewhat difficult
☐ Very difficult

☐ *Don't know*
☐ *Prefer not to answer*

1.19 What type of place do you currently live in? (Check ONE only)

- ☐ Your house, your apartment, your condo (owner)
☐ Your house, your apartment, your condo (rented)
☐ Your parents' house
☐ Another family member's house
☐ Friend's house
☐ Hotel or motel room
☐ Shelter or hostel
☐ Transition house or halfway house
☐ Rooming or boarding house
☐ Public place (street, park, squats, subway, underpass, washroom, stairwell, etc.)

☐ Refugee camp or immigration detention facility
☐ Other (specify): _____

☐ *Don't know*
☐ *Prefer not to answer*

1.20 How many times have you moved in the last 12 months?

Number of times: _____

- ☐ *Don't know*
☐ *Prefer not to answer*

SECTION TWO: Sexual behaviours

2.1 How old were you the first time you had vaginal or anal sex?

_____ (Age in years)

☐ Never had sex

→ If you are a man and have never had sex, go to question 2.27.

☐ Don't know

→ If you are a woman and have never had sex, go to question 2.28.

☐ Prefer not to answer

2.2 How many people in your lifetime, including men and women, have you had vaginal or anal sex with?

This includes people you have had sex with only one time. (Check ONE only)

☐ 1 partner

☐ 2 partners

☐ 3 partners

☐ 4 to 5 partners

☐ 6 to 10 partners

☐ 11 to 19 partners

☐ 20 or more partners

☐ Don't know

☐ Prefer not to answer



2.3 How many of these sex partners were born in Sub-Saharan Africa, the Caribbean, Central / South America or Asia? (See countries highlighted on map above for reference. Check *ONE* only)

- ☐ None of them
☐ Some of them
☐ All of them

☐ *Don't know*
☐ *Prefer not to answer*

2.4 Have you had vaginal or anal sex in the last 12 months?

- ☐ Yes
☐ No

☐ *Don't know*
☐ *Prefer not to answer*

If the answer is No, Don't know, or Prefer not to answer go to question 2.27 if it's a man or to question 2.28 if it's a woman.

2.4 a Did you or your partner use a condom the last time you had sex? Including male and female condoms.

- ☐ Yes
☐ No

☐ *Don't know*
☐ *Prefer not to answer*

2.5 In the last 12 months, with how many MALE partners have you had vaginal or anal sex with? (Check ONE answer only)

- ☐ None
- ☐ 1 partner
- ☐ 2 partners
- ☐ 3 partners
- ☐ 4 to 5 partners
- ☐ 6 to 10 partners
- ☐ 11 to 19 partners
- ☐ 20 or more partners

→ If the answer is None, go to question 2.13.

- ☐ Don't know
- ☐ Prefer not to answer

2.6 In the last 12 months, how many REGULAR MALE sex partners have you had vaginal or anal sex with?
(Check ONE only)

- ☐ None
- ☐ 1 partner
- ☐ 2 partners
- ☐ 3 partners
- ☐ 4 to 5 partners
- ☐ 6 to 10 partners
- ☐ 11 to 19 partners
- ☐ 20 or more partners

→ If the answer is None, go to question 2.10.

- ☐ Don't know
- ☐ Prefer not to answer

2.7 In the last 12 months, how often did you use condoms with your REGULAR MALE sex partners? (Check ONE answer only)

- ☐ Wanted to but never had a condoms
- ☐ Never (0%)
- ☐ Rarely (less than 25% of the time)
- ☐ Sometimes (25 to 49% of the time)
- ☐ Most of the time (50 to 74% of the time)
- ☐ Almost every time (75 to 99% of the time)
- ☐ All the time (100% of the time)

} If Wanted to or Never, go to question 2.8.

- ☐ Don't know
- ☐ Prefer not to answer

2.7a The last time you had sex with your REGULAR MALE sex partner, was a condom used?

- ☐ Yes
- ☐ No

- ☐ Don't know
- ☐ Prefer not to answer

2.8 In the last 12 months, months, did any of your REGULAR MALE sex partners have any other sexual partners while in a sexual relationship with you?

- ☐ Yes
- ☐ No
- ☐ *Don't know*
- ☐ *Prefer not to answer*

2.9 What is the HIV status of your most recent REGULAR MALE sex partner? (Check ONE only)

- ☐ Seropositive
- ☐ Seronegative
- ☐ You don't know the HIV status of your last REGULAR MALE sexual partner.
- ☐ *Prefer not to answer*

2.10 In the past 12 months, with how many MALE CASUAL sex partners have you had vaginal or anal sex with penetration? (Check ONE answer only)

- ☐ None → If the answer is None, go to question 2.13.
- ☐ 1 partner
- ☐ 2 partners
- ☐ 3 partners
- ☐ 4 to 5 partners
- ☐ 6 to 10 partners
- ☐ 11 to 19 partners
- ☐ 20 or more partners
-
- ☐ *Don't know*
- ☐ *Prefer not to answer*

2.11 In the past 12 months, , how often did you use condoms with your CASUAL MALE sex partner(s)?
(Check ONE only)

- ☐ Wanted to but never had
- ☐ Never (0%)
- ☐ Rarely (less than 25% of the time)
- ☐ Sometimes (25 to 49% of the time)
- ☐ Most of the time (50 to 74% of the time)
- ☐ Almost every time (75 to 99% of the time)
- ☐ All the time (100% of the time)

- ☐ Don't know
- ☐ Prefer not to answer



Condoms If Wanted to or Never, go to question 2.12.

2.11a The last time you had sex with your CASUAL MALE sex partner, was a condom used?

- ☐ Yes
- ☐ No

- ☐ Don't know
- ☐ Prefer not to answer

2.12 What is the HIV status of your most recent CASUAL MALE sex partner? (Check ONE only)

- ☐ Seropositive
- ☐ Seronegative

- ☐ You don't know the HIV status of your most recent CASUAL MALE sex partner.

- ☐ Prefer not to answer

2.13 In the last 12 months, how many FEMALE sex partners have you had vaginal or anal sex with? (Check ONE only)

- ☐ None → If the answer is None, go to question 2.21.
- ☐ 1 partner
- ☐ 2 partners
- ☐ 3 partners
- ☐ 4 to 5 partners
- ☐ 6 to 10 partners
- ☐ 11 to 19 partners
- ☐ 20 or more partners

- ☐ Don't know
- ☐ Prefer not to answer

2.14 In the last 12 months, how many REGULAR FEMALE sex partners have you had vaginal or anal sex with? (Check ONE only)

- ☐ None → If the answer is None, go to question 2.18.
- ☐ 1 partner
- ☐ 2 partners
- ☐ 3 partners
- ☐ 4 to 5 partners
- ☐ 6 to 10 partners
- ☐ 11 to 19 partners
- ☐ 20 or more partners

- ☐ Don't know
- ☐ Prefer not to answer

2.15 In the last 12 months, how often did you use condoms with your REGULAR FEMALE sex partner(s)?
(Check ONE only)

- ☐ Wanted to but never had
- ☐ Never (0%)
- ☐ Rarely (less than 25% of the time)
- ☐ Sometimes (25 to 49% of the time)
- ☐ Most of the time (50 to 74% of the time)
- ☐ Almost every time (75 to 99% of the time)
- ☐ All the time (100% of the time)

If Wanted to or Never, go to question 2.16.

- ☐ Don't know
- ☐ Prefer not to answer

2.15a The last time you had sex with your REGULAR FEMALE sex partner, was a condom used?

- ☐ Yes
- ☐ No

- ☐ Don't know
- ☐ Prefer not to answer

2.16 In the past 12 months, did any of your REGULAR FEMALE sexual partners have any other sexual partners while in a sexual relationship with you?

- ☐ Yes
- ☐ No

- ☐ Don't know
- ☐ Prefer not to answer

2.17 What is the HIV status of your most recent REGULAR FEMALE sex partner? (Check ONE only)

- ☐ Seropositive
- ☐ Seronegative

☐ You don't know the HIV status of your most recent REGULAR FEMALE sex partner.

☐ Prefer not to answer

2.18 In the past 12 months, with how many FEMALE CASUAL sex partners have you had vaginal or anal sex with penetration? (Check ONE answer only)

- ☐ None → If the answer is None, go to question 2.21.
- ☐ 1 partner
- ☐ 2 partners
- ☐ 3 partners
- ☐ 4 to 5 partners
- ☐ 6 to 10 partners
- ☐ 11 to 19 partners
- ☐ 20 or more partners
- ☐ Don't know
- ☐ Prefer not to answer

2.19 In the last 12 months, how many CASUAL FEMALE sex partners have you had vaginal or anal sex with? (Check ONE only)

- ☐ Wanted to but never had
- ☐ Never (0%)
- ☐ Rarely (less than 25% of the time)
- ☐ Sometimes (25 to 49% of the time)
- ☐ Most of the time (50 to 74% of the time)
- ☐ Almost every time (75 to 99% of the time)
- ☐ All the time (100% of the time)
- ☐ Don't know
- ☐ Prefer not to answer
- } If the answer is Planned to or Never, go to question 2.20.

2.19 In the last 12 months, how often did you use condoms with your CASUAL FEMALE sex partner(s)? (Check ONE only)

- ☐ Yes
- ☐ No
- ☐ Don't know
- ☐ Prefer not to answer

2.20 What is the HIV status of your last FEMALE CASUAL sex partner? (Check ONE answer only)

- ☐ Seropositive
☐ Seronegative

☐ You don't know the HIV status of your most recent CASUAL FEMALE sex partner.

☐ *Don't know*
☐ *Prefer not to answer*

2.21 In the last 12 months, to how many sexual partners have you GIVEN money, drugs, goods or services in exchange for vaginal or anal sex with penetration? (Check ONE answer only)

- ☐ None
☐ 1 partner
☐ 2 partners
☐ 3 partners
☐ 4 to 5 partners
☐ 6 to 10 partners
☐ 11 to 19 partners
☐ 20 or more partners

☐ *Don't know*
☐ *Prefer not to answer*
- If the answer is None, go to question 2.23.

2.22 In the last 12 months, how often did you use condoms with sex partners who you GAVE money, drugs, or other goods or services in exchange for vaginal or anal sex? (Check ONE only)

- ☐ Wanted to but never had
☐ Never (0%)
☐ Rarely (less than 25% of the time)
☐ Sometimes (25 to 49% of the time)
☐ Most of the time (50 to 74% of the time)
☐ Almost every time (75 to 99% of the time)
☐ All the time (100% of the time)

☐ *Don't know*
☐ *Prefer not to answer*
- } If wanted to or Never, go to question 2.23.

2.22 a The last time you had vaginal or anal sex with penetration with a sexual partner to whom you GAVE money, drugs, goods or services in exchange for sex, did you or your partner use a condom?

- ☐ Yes
☐ No
☐ Don't know
☐ Prefer not to answer

2.23 In the last 12 months, from how many sexual partners have you RECEIVED money, drugs, goods or services in exchange for vaginal or anal sex with penetration? (Check ONE answer only)

- ☐ None
☐ 1 partner
☐ 2 partners
☐ 3 partners
☐ 4 to 5 partners
☐ 6 to 10 partners
☐ 11 to 19 partners
☐ 20 or more partners
☐ Don't know
☐ Prefer not to answer

→ If the answer is None, go to question 2.25.

2.24 In the last 12 months, how often did you use condoms with sex partners from whom you RECEIVED money, drugs, or other goods or services in exchange for vaginal or anal sex? (Check ONE only)

- ☐ Wanted to but never had
☐ Never (0%)
☐ Rarely (less than 25% of the time)
☐ Sometimes (25 to 49% of the time)
☐ Most of the time (50 to 74% of the time)
☐ Almost every time (75 to 99% of the time)
☐ All the time (100% of the time)

} If wanted to or Never, go to question 2.25.

- ☐ Don't know
☐ Prefer not to answer

2.24 a last time you had vaginal or anal sex with a partner from whom you RECEIVED money, drugs or other goods or services in exchange for vaginal or anal sex, did you or your partner use a condom?

☐ Yes

☐ No

☐ Don't know

☐ Prefer not to answer

2.25 In the past 12 months, have you been drunk, high or buzzed on any of the following substances while you were having vaginal or anal sex? (Check ONE response for each item)

	Yes	No	Don't know	Prefer not to answer
Alcohol	☐	☐	☐	☐
Marijuana or hash or pot or grass	☐	☐	☐	☐
Poppers	☐	☐	☐	☐
Special K or ketamine	☐	☐	☐	☐
Ecstasy or MDMA	☐	☐	☐	☐
Other amphetamines or stimulants	☐	☐	☐	☐
Cocaine or crack or free base	☐	☐	☐	☐
Heroin (smack)	☐	☐	☐	☐
Other opioids (percocet, dilaudid, oxycontin or oxyneo)	☐	☐	☐	☐
Crystal meth	☐	☐	☐	☐
GHB ("G")	☐	☐	☐	☐
Tranquilizers or benzodiazepine (benzos)	☐	☐	☐	☐
Psychedelics (LSD, acid, mescaline or mushrooms)	☐	☐	☐	☐
Other (specify): _____	☐	☐	☐	☐

2.26 In the past 12 months, have you engaged in dry sex?

- ☐ Yes
☐ No


☐ *Don't know*
☐ *Prefer not to answer*

2.27 Have you been circumcised (cut)?

- ☐ Yes
☐ No

☐ Does not apply

☐ *Don't know*
☐ *Prefer not to answer*



Go to question 3.1.

2.28 Have you had any kind of female excision?

- ☐ Yes
☐ No

☐ *Don't know*
☐ *Prefer not to answer*

Section THREE: Travel History



3.1 In the past five years, have you travelled to any of the highlighted countries in Sub-Saharan Africa, the Caribbean, Central / South America or Asia? (See countries highlighted on map above for reference)

☐ Yes

☐ No

☐ Don't know

☐ Prefer not to answer

Prefer not to answer, go to question 4.1.

MTL3 (3.1) Specify the country or countries that you have travelled to in the past five years and the number of times that you have travelled there during that period.

Country: _____	Frequency: _____
Country: _____	Frequency: _____
Country: _____	Frequency: _____
Country: _____	Frequency: _____
Country: _____	Frequency: _____

☐ Don't know

☐ Prefer not to answer

3.1 a During these trips, did you have vaginal or anal sex with anyone who lives in the specified countries in Sub-Saharan Africa, the Caribbean, Central / South America or Asia?

☐ Yes

☐ No

☐ Never had sex

☐ Don't know

☐ Prefer not to answer

*If the answer is No, Never had sex,
Don't know, or Prefer not to answer, go to
question 4.1.*

3.1 b How often did you use condoms with these sex partners? (Check ONE answer only)

☐ Wanted to but never had

☐ Never (0%)

☐ Rarely (less than 25% of the time)

☐ Sometimes (25 to 49% of the time)

☐ Most of the time (50 to 74% of the time)

☐ Almost every time (75 to 99% of the time)

☐ All the time (100% of the time)

☐ Don't know

☐ Prefer not to answer

Section FOUR: Drug Use

4.1 Have you ever injected drugs (shot up) for non-medical reasons? This includes self-injection or injection by someone else. (Check ONE only)

- ☐ Never
- ☐ Yes, but not in the past six months
- ☐ Yes, in the past six months

} If the answer is Never or Yes, but not in the last six months, go to question 5.1.

- ☐ Don't know
- ☐ Prefer not to answer

} If the person replied Don't know or Prefer not to answer, go to question 5.1.

4.1 a In the past 6 months, have you used needles and / or syringes that had already been used by someone else? This includes all used needles and syringes, whether they had been cleaned or not.

- ☐ Yes
- ☐ No
- ☐ Don't know
- ☐ Prefer not to answer

Section FIVE: Access to and Use of Health Systems and Services**NOTE!**

5.1 Do you have a family doctor or a primary health care provider?

- ☐ Yes
☐ No

☐ *Don't know*
☐ *Prefer not to answer*

5.2 In the past 12 months, did you visit a health care provider or a health facility for medical attention, consultation, health information or to take part in a program?

- ☐ Yes
☐ No

☐ *Don't know*
☐ *Prefer not to answer*

} *If the answer is No, Don't know, or Prefer not to answer, go to question 5.4.*

5.3 In the past 12 months which of the following health providers, facilities or services have you used or visited for medical attention, health information or to take part in a program? (Check ONE response for each item)

	Yes	No	Don't know	Prefer not to answer
	☐	☐	☐	☐
Hospital	☐	☐	☐	☐
	☐	☐	☐	☐
Medical clinic or walk-in clinic	☐	☐	☐	☐
	☐	☐	☐	☐
Community organization (Community day centre)	☐	☐	☐	☐
	☐	☐	☐	☐
Needle exchange or harm reduction program:				
In a fixed location in a community organization like Cactus, Spectre de rue, Dopamine, etc.	☐	☐	☐	☐
In a CLSC	☐	☐	☐	☐
In a drug store	☐	☐	☐	☐
Other (specify): _____	☐	☐	☐	☐
	☐	☐	☐	☐
Mental health or addiction centre	☐	☐	☐	☐
Prison / Corrections facility	☐	☐	☐	☐
	☐	☐	☐	☐
Other (specify):	☐	☐	☐	☐

5.4 In the past 12 months, have you experienced any difficulties getting the health care you needed?*(Check ONE only)*☐ Yes☐ No☐ You have not required any health care in the past 12☐ Don't know☐ Prefer not to answer

} If the answer is Yes, go to
question 5.5.

5.4a What type of difficulties did you experience? (Check ALL answers that apply)☐ No health care providers were available in your area.☐ The specific service you needed was not available in your area.☐ You didn't know how to find the health care you needed.☐ You found it difficult to get an appointment, diagnosis or referral.☐ The wait time was too long.☐ You had problems getting there.☐ There was a language barrier (i.e. you weren't able to communicate with the health care service provider).☐ You couldn't pay for the health care service or the service wasn't covered by your insurance.☐ Personal or family responsibilities prevented you from receiving the health care you needed.☐ You weren't able to leave home because of a health problem.☐ The health care provider was not culturally sensitive or was racist.☐ The health care provider was of the opposite sex.☐ Other (specify): _____☐ Don't know☐ Prefer not to answer**5.5 Have you ever had a vaccine against the following? (Check ONE response for each item)**

	Yes, in the past 6 months	Yes, more than 6 months ago	No	Don't know	Prefer not to answer
Hepatitis B (e.g. Engerix-B, Pediatrix, Recombivax HB)	☐	☐	☐	☐	☐
Human papillomavirus (HPV) (e.g, Gardasil, Cervarix)	☐	☐	☐	☐	☐

5.6 Have you ever been tested for tuberculosis?☐ Yes☐ No☐ *Don't know*☐ *Prefer not to answer***5.7 Have you ever been told by a health professional (e.g. doctor or nurse) that you have or had any of the following infections? (Check ONE response for each item)**

	Yes, in the In the past 6 months	Yes, than 6 months	No	<i>Don't know</i>	<i>Prefer not to answer</i>
Tuberculosis (Tb)	☐	☐	☐	☐	☐
Hepatitis B	☐	☐	☐	☐	☐
Hepatitis C	☐	☐	☐	☐	☐
Chlamydia	☐	☐	☐	☐	☐
Gonorrhoea	☐	☐	☐	☐	☐
Syphilis	☐	☐	☐	☐	☐
Genital or anal warts	☐	☐	☐	☐	☐
Human Papillomavirus	☐	☐	☐	☐	☐
Genital Herpes	☐	☐	☐	☐	☐
Lymphogranuloma venereum	☐	☐	☐	☐	☐

Section SIX: HIV Testing, Care and Treatment

6.1 Have you ever been tested for HIV?

☐ Yes

☐ No

☐ Don't know

☐ Prefer not to answer


If the answer is No, Don't know, or Prefer not to answer, go to question 6.17.

6.2 Why did you take your last HIV test? (Check ALL answers that apply)

☐ You were tested during your immigration medical examination.

☐ You get tested routinely for HIV and this was a routine test

☐ You took the test after starting a new relationship.

☐ You were (or are) pregnant or plan to have a child.

☐ Your friends, family or partner encouraged you to take the test.

☐ You got tested after an event that might have put you at risk for HIV

☐ You had symptoms or illness that you thought might be related to HIV

☐ You found out that a sex partner was or might be HIV positive

☐ A health worker recommended that you get tested for HIV

☐ You were being tested for another sexually transmitted infection

☐ You saw an ad, poster, or article about HIV

☐ A Public Health Branch told you that you may have been exposed to HIV.

☐ You were notified by a public health unit that you could have been exposed to HIV

☐ Other reason (specify): _____

☐ No particular reason

☐ Don't know

☐ Prefer not to answer

6.3 Were you tested for HIV during your last pregnancy? (Check ONE only)

- ☐ You have never been pregnant → *If the person has never been pregnant, go to question 6.4.*
- ☐ Yes, you were tested for HIV during your last pregnancy
- ☐ No, you were not tested for HIV during your last pregnancy
- ☐ *Don't know*
- ☐ *Prefer not to answer*
- } *If the person replied Don't know or Prefer not to answer, go to question 6.4.*

6.3 a Were you living in Canada during your last pregnancy?

- ☐ Yes
- ☐ No
- ☐ *Don't know*
- ☐ *Prefer not to answer*

6.4 When was your most recent HIV test? (Check ONE only)

- ☐ Within the past month
- ☐ 1 to 6 months ago
- ☐ 7 to 12 months ago
- ☐ More than 1 year ago, but less than 2 years ago
- } *Go to question 6.5.*
- ☐ More than 2 years ago, but less than 4 years ago.
- ☐ More than 4 years ago
- ☐ *Don't know*
- ☐ *Prefer not to answer*
- } *If the person replied Don't know or Prefer not to answer, go to question 6.5.*

6.4 a The following following are some of the reasons people give for not getting an HIV test. Have you not been tested for HIV in the past 2 years because: (Check ALL that apply)

- ☐ You've never thought of it.
- ☐ You're afraid to get tested.
- ☐ You think your risk of HIV infection is low.
- ☐ You had trouble getting tested.
- ☐ You're healthy, so you think you don't need to take the test.
- ☐ You think the test is not always reliable.
- ☐ You think you can't get HIV.
- ☐ You think nothing can be done if the test indicates you have HIV.
- ☐ You think being HIV positive would affect your personal or professional life.
- ☐ You think you know your HIV status.
- ☐ You were tested during your immigration medical examination in Canada.
- ☐ Other (specify): _____
- ☐ Don't know
- ☐ Prefer not to answer

6.5 What was the result of your most recent HIV test? (Check ONE answer only)

- ☐ HIV positive: you have the virus
 - ☐ HIV negative: you do NOT have the virus
 - ☐ Indeterminate result: the result was inconclusive.
 - ☐ You didn't understand the result.
 - ☐ You are still waiting for the result.
 - ☐ The result is known, but you haven't received it yet.
 - ☐ Don't know
 - ☐ Prefer not to answer
- } If the answer is HIV negative or indeterminate, or if the result was not understood, could not be accessed or was not received, go to question 7.1.
- } If the person replied Don't know or Prefer not to answer, go to question 7.1.

6.6 When did you get your first HIV positive result? (Check ONE answer only)

- ☐ Within the last month
 - ☐ 1 to 6 months ago
 - ☐ 7 to 12 months ago
 - ☐ More than 1 year ago, but less than 2 years ago
 - ☐ More than 2 years ago, but and less than 4 years ago
 - ☐ 4 years ago or more
 - ☐ Don't know
 - ☐ Prefer not to answer
- } If the person replied Don't know or Prefer not to answer, go to question 6.8.

6.7 Before your first HIV positive test, when was the last time you had a negative HIV test result? (Check ONE answer only)

- ☐ 1 to 6 months before the first positive test
- ☐ 7 to 12 months before the first positive test
- ☐ More than 12 months before the first positive test
- ☐ I was never tested for HIV before the first positive test

- ☐ Don't know
- ☐ Prefer not to answer

6.8 Do you know how you got infected?

- ☐ Yes
- ☐ No
- ☐ Prefer not to answer

} If the answer is No or Prefer not to answer, go to question 6.9.

6.8 To the best of your knowledge, how did you become infected with HIV? (Check ONE only)

- ☐ Sex with an HIV-positive man
- ☐ Sex with an HIV-positive woman
- ☐ Blood transfusion
- ☐ Occupational exposure, e.g., accidental puncture with a syringe
- ☐ Drug injections with used syringes
- ☐ Mother-to-child transmission during birth or breastfeeding
- ☐ Razors or syringes that were not disinfected, e.g., scarification or traditional healing technique
- ☐ Other (specify): _____
- ☐ Don't know
- ☐ Prefer not to answer

6.9 To the best of your knowledge, were you infected with HIV in Canada or in another country? (Check ONE only)

- ☐ In Canada
- ☐ In another country
- ☐ Don't know
- ☐ Prefer not to answer

6.10 How long after your first HIV diagnosis did you start seeing a doctor for your HIV infection? (Check ONE only)

- ☐ You have never seen a doctor for your HIV infection →
- ☐ 3 months or less after your diagnosis
- ☐ 4 months to 6 months after your
- ☐ 7 months to one year after your
- ☐ More than a year after your diagnosis

- ☐ Don't know
- ☐ Prefer not to answer

If the answer is you have never seen a doctor for your HIV infection, go to question 7.1.

6.11 Are you currently under the care of a doctor or other health care professional for your HIV infection? This means a single visit or more to a doctor or other health professional in the past six months for HIV monitoring (e.g., viral load, CD4 cell count), treatment, counselling, etc.

- ☐ Yes
- ☐ No

- ☐ Don't know
- ☐ Prefer not to answer

6.12 Have you EVER taken prescription drugs to treat your HIV infection?

- ☐ Yes
- ☐ No

- ☐ Don't know
- ☐ Prefer not to answer

→ *If the answer is No, go to question 6.15.*

} *If the person replied Don't know or Prefer not to answer, go to question 6.16.*

6.13 Where

- ☐ In Canada
- ☐ In another country
- ☐ Don't know
- ☐ Prefer not to answer

6.14 Are you CURRENTLY taking prescription drugs to treat your HIV infection?

- ☐ Yes
☐ No

→ If the answer is Yes, go to
question 6.16.

- ☐ Don't know
☐ Prefer not to answer



If the person replied Don't know or
Prefer not to answer, go to question
6.16.

6.14 a When did you stop taking prescription drugs to treat your HIV infection? (Check ONE answer only)

- ☐ Within the last month
☐ 1 to 6 months ago
☐ 7 to 12 months ago
☐ More than 1 year ago, but less than 2 years ago
☐ More than 2 years ago, but and less than 4 years ago
☐ 4 years ago or more than
- ☐ Don't know
☐ Prefer not to answer

6.14 b Why are you not taking prescription drugs to treat your HIV infection? (Check ALL answers that apply)

- ☐ You don't have a doctor.
- ☐ You had far too many logistical challenges to get your HIV medications, e.g., transportation, distance between home and the drug store, opening hours of the clinic or pharmacy.
- ☐ You tried taking them and stopped because you couldn't afford them.
- ☐ You tried taking them and stopped because you had side effects.
- ☐ You tried taking them and stopped because the prescription was too complicated or it was too difficult to take the medications.
- ☐ Your doctor told you that you didn't need them yet for medical reasons.
- ☐ Your doctor told you that you were not ready to take them, for non-medical reasons.
- ☐ You were never prescribed any.
- ☐ They are not covered by your insurance policy.
- ☐ Your doctor recommended that you stop treatment because of side effects, another illness, travel restrictions or surgery.
- ☐ Other (specify): _____
- ☐ Don't know
- ☐ Prefer not to answer

6.15 Why have you never taken prescription drugs to treat your HIV infection? (Check ALL answers that apply)

- ☐ You never started taking them because you couldn't afford them.
- ☐ Your doctor never discussed treatment with you or didn't offer you the medications.
- ☐ Your doctor told you that you didn't need them yet, for medical reasons.
- ☐ Your doctor told you that you were not ready to take them, for non-medical reasons.
- ☐ They are not covered by your insurance policy.
- ☐ Other (specify): _____
- ☐ Don't know
- ☐ Prefer not to answer

6.16 Have you ever used an alternative or traditional medical treatment for your HIV infection?

- ☐ Yes
 - ☐ No

 - ☐ *Don't know*
 - ☐ *Prefer not to answer*
- 

6.17 The following statements provide some reasons people give for never having taken an HIV test. You have never taken an HIV test because: (Check ALL answers that apply)

- ☐ You never thought about it
- ☐ You're afraid to get tested
- ☐ You think your risk of HIV infection is low
- ☐ You had trouble getting tested
- ☐ You're healthy, so you think you don't need to take the test
- ☐ You think the test is not always reliable
- ☐ You think you can't get an HIV infection
- ☐ You think nothing can be done if the test indicates you have HIV
- ☐ You think being HIV positive would affect your personal or professional life
- ☐ You think you know your HIV status
- ☐ Other (specify): _____

- ☐ *Don't know*
- ☐ *Prefer not to answer*

Section SEVEN: Knowledge and Beliefs about HIV

7.1 In the last 12 months, apart from this survey, where have you heard about, seen or read about HIV or AIDS?

(Check ALL answers that apply)

- ☐ Friends
- ☐ Family
- ☐ Television
- ☐ Radio
- ☐ Newspapers or magazines
- ☐ Social media (Twitter, Facebook, etc.)
- ☐ Websites or the Internet (Google, Wikipedia, etc.)
- ☐ Advertising (flyers, posters, etc.)
- ☐ Doctors, nurses, social workers, pharmacists and other health professionals
- ☐ At school
- ☐ At work
- ☐ At seminars, conferences or workshops
- ☐ At a medical clinic or hospital
- ☐ At local or provincial public health service locations
- ☐ Through Health Canada or the Public Health Agency of Canada
- ☐ Through an HIV/AIDS service organization for community organizations
- ☐ Through churches or other religious institutions
- ☐ Other (specify): _____
- ☐ You haven't seen or heard anything about HIV or AIDS
- ☐ *Don't know*
- ☐ *Prefer not to answer*

7.2 Based on what you know, answer Yes or No to the following questions. There is nothing wrong with saying you don't know. This is not a test. (Check ONLY ONE box for each option)

	Yes	No	Don't know	Prefer not to answer
Can having sex with only one, faithful, uninfected partner reduce the risk of HIV transmission?	☐	☐	☐	☐
Can using condoms reduce the risk of HIV transmission?	☐	☐	☐	☐
Can a healthy-looking person have HIV?	☐	☐	☐	☐
Can a person get HIV from mosquito bites?	☐	☐	☐	☐
Can a person get HIV by sharing a meal with someone who is infected?	☐	☐	☐	☐
Is there currently a cure for HIV?	☐	☐	☐	☐

If you are completing it on your own, please inform the research assistant that you are ready to have your blood sample taken.

Enter the interview end time. Please indicate in 24-hour time. For example, 5:45pm is recorded as 17:45. .

Do you have any comments about the questionnaire? For example: Were any of the questions confusing? Did you object to answering any of the questions? How did you find the length of the questionnaire?

DBS collection

Was the DBS collected?

☐ Yes

☐ No (indicate the reason): _____

Record time taken for DBS collection: _____ (minutes)

Do you have any general comments about the survey?
