

**Geometric-arithmetic index and
minimum degree of connected graphs**

M. Aouchiche, I. El Hallaoui,
P. Hansen

G-2019-58

August 2019

La collection *Les Cahiers du GERAD* est constituée des travaux de recherche menés par nos membres. La plupart de ces documents de travail a été soumis à des revues avec comité de révision. Lorsqu'un document est accepté et publié, le pdf original est retiré si c'est nécessaire et un lien vers l'article publié est ajouté.

Citation suggérée : M. Aouchiche, I. El Hallaoui, P. Hansen (Août 2019). Geometric-arithmetic index and minimum degree of connected graphs, Rapport technique, Les Cahiers du GERAD G-2019-xx, GERAD, HEC Montréal, Canada.

Avant de citer ce rapport technique, veuillez visiter notre site Web (<https://www.gerad.ca/fr/papers/G-2019-xx>) afin de mettre à jour vos données de référence, s'il a été publié dans une revue scientifique.

La publication de ces rapports de recherche est rendue possible grâce au soutien de HEC Montréal, Polytechnique Montréal, Université McGill, Université du Québec à Montréal, ainsi que du Fonds de recherche du Québec – Nature et technologies.

Dépôt légal – Bibliothèque et Archives nationales du Québec, 2019
– Bibliothèque et Archives Canada, 2019

The series *Les Cahiers du GERAD* consists of working papers carried out by our members. Most of these pre-prints have been submitted to peer-reviewed journals. When accepted and published, if necessary, the original pdf is removed and a link to the published article is added.

Suggested citation: M. Aouchiche, I. El Hallaoui, P. Hansen (August 2019). Geometric-arithmetic index and minimum degree of connected graphs, Technical report, Les Cahiers du GERAD G-2019-xx, GERAD, HEC Montréal, Canada.

Before citing this technical report, please visit our website (<https://www.gerad.ca/en/papers/G-2019-xx>) to update your reference data, if it has been published in a scientific journal.

The publication of these research reports is made possible thanks to the support of HEC Montréal, Polytechnique Montréal, McGill University, Université du Québec à Montréal, as well as the Fonds de recherche du Québec – Nature et technologies.

Legal deposit – Bibliothèque et Archives nationales du Québec, 2019
– Library and Archives Canada, 2019

Geometric-arithmetic index and minimum degree of connected graphs

Mustapha Aouchiche ^a

Issmail El Hallaoui ^{b, c}

Pierre Hansen ^{b, d}

^a Department of Mathematical Sciences, COS, UAEU, Al-Ain, UAE

^b GERAD, Montréal (Québec), Canada

^c Department of Mathematics and Industrial Engineering, Polytechnique Montréal, Montréal (Québec) Canada

^d Department of Decision Sciences, HEC Montréal, Montréal (Québec), Canada

maouchiche@uaeu.ac.ae

issmail.elhallaoui@gerad.ca

pierre.hansen@gerad.ca

August 2019

Les Cahiers du GERAD

G–2019–58

Copyright © 2019 GERAD, Aouchiche, El Hallaoui, Hansen

Les textes publiés dans la série des rapports de recherche *Les Cahiers du GERAD* n'engagent que la responsabilité de leurs auteurs. Les auteurs conservent leur droit d'auteur et leurs droits moraux sur leurs publications et les utilisateurs s'engagent à reconnaître et respecter les exigences légales associées à ces droits. Ainsi, les utilisateurs:

- Peuvent télécharger et imprimer une copie de toute publication du portail public aux fins d'étude ou de recherche privée;
- Ne peuvent pas distribuer le matériel ou l'utiliser pour une activité à but lucratif ou pour un gain commercial;
- Peuvent distribuer gratuitement l'URL identifiant la publication.

Si vous pensez que ce document enfreint le droit d'auteur, contactez-nous en fournissant des détails. Nous supprimerons immédiatement l'accès au travail et enquêterons sur votre demande.

The authors are exclusively responsible for the content of their research papers published in the series *Les Cahiers du GERAD*. Copyright and moral rights for the publications are retained by the authors and the users must commit themselves to recognize and abide the legal requirements associated with these rights. Thus, users:

- May download and print one copy of any publication from the public portal for the purpose of private study or research;
- May not further distribute the material or use it for any profit-making activity or commercial gain;
- May freely distribute the URL identifying the publication.

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract: In the present paper, we prove lower and upper bounds for each of the ratios GA/δ , as well as a lower bound on $GA/\sqrt{\delta}$, in terms of the order n , over the class of connected graphs on n vertices, where GA and δ denote the geometric-arithmetic index and the minimum degree, respectively. We also characterize the extremal graphs corresponding to each of those bounds. In order to prove our results, we provide a modified statement of a well-known lower bound on the geometric-arithmetic index in terms of minimum degree.

Keywords: Geometric-arithmetic index, minimum degree, extremal graph, molecular descriptor

1 Introduction and definitions

We begin by recalling some definitions. In this paper, we consider only simple, undirected and finite graphs, *i.e.*, undirected graphs on a finite number of vertices without multiple edges or loops. A graph is (usually) denoted by $G = G(V, E)$, where V is its vertex set and E its edge set. The *order* of G is the number $n = |V|$ of its vertices and its *size* is the number $m = |E|$ of its edges. For two vertices u and v ($u, v \in V$), if $uv \in E$, we say u and v are *adjacent* in G . The *degree* of a vertex u , denoted d_u , is the number of vertices adjacent to it in G . A graph G is said to be *regular* of degree d , or *d-regular* if $d_u = d$ for every vertex u in G . The minimum degree in a graph G is denoted by δ . As usual, we denote by S_n the star and by K_n the complete graph, each on n vertices.

Molecular descriptors play a very important role in mathematical chemistry especially in QSAR (quantitative structure-activity relationship) and/or QSPR (quantitative structure-property relationship) related studies. Among those descriptors, a special interest is devoted to so-called topological indices. They are used to understand physicochemical properties of chemical compounds in a simple way, since they sum up some of the properties of a molecule in a single number. During the last decades, a legion of topological indices were introduced and found some applications in chemistry, see e.g., [7, 8, 16]. The study of topological indices goes back to the seminal work by Wiener [18] in which he used the sum of all shortest-path distances, nowadays known as the *Wiener index*, of a (molecular) graph for modeling physical properties of alkanes.

Another very important molecular descriptor, was introduced by Randić [13]. It is called the *Randić (connectivity) index* and defined as

$$Ra = Ra(G) = \sum_{uv \in E} \frac{1}{\sqrt{d_u d_v}}$$

where d_u denotes the degree (number of neighbors) of u in G . The Randić index is probably the most studied molecular descriptor in mathematical chemistry. Actually, there are more than two thousand papers and five books devoted to this index (see, e.g., [6, 9, 10, 11, 12] and the references therein).

Motivated by the definition of Randić connectivity index, Vukičević and Furtula [17] proposed the *geometric-arithmetic index*. It is so-called since its definition involves both the geometric and the arithmetic means of the endpoints degrees of the edges in a graph. For a simple graph G , the geometric-arithmetic index $GA(G)$ of a graph G is defined as in [17] by

$$GA = GA(G) = \sum_{uv \in E} \frac{2\sqrt{d_u d_v}}{d_u + d_v}.$$

It is noted in [17] that the predictive power of GA for physico-chemical properties is somewhat better than the predictive power of the Randić connectivity index. In [17], Vukičević and Furtula gave lower and upper bounds for GA , identified the trees with the minimum and the maximum GA indices, which are the star S_n and the path P_n , respectively. In [19] Yuan, Zhou and Trinajstić gave the lower and upper bounds for GA index of molecular graphs using the numbers of vertices and edges. They also determined the n -vertex molecular trees with the minimum, the second, and the third minimum, as well as the second and the third maximum GA indices. The chemical applicability of the geometric-arithmetic index was highlighted in [3, 5, 17].

Lower and upper bounds on the geometric-arithmetic index in terms of order n , size m , minimum degree δ and/or maximum degree were proved in [14]. Also in [14], GA was compared to other well known topological indices such as the Randić index, the first and second Zagreb indices, the harmonic index and the sum connectivity index. Other lower and upper bounds, on the geometric-arithmetic index, involving the order n the size m , the minimum and the maximum degrees and the second Zagreb index were proved in [2]

In [1], several bounds and comparisons, involving the geometric-arithmetic index and several other graph parameters, were proved.

In the present paper we deal with the problem of finding upper and lower bounds, with the characterization of the corresponding extremal graphs, for the geometric-arithmetic index of a connected graph with given number of vertices n and minimum degree δ . Earlier study of this problem can be found in [4, 15].

2 Main results

In this section, we first prove upper bounds on the ratios GA/δ in terms of the order n , over the class of connected graphs on n vertices. We also characterize the corresponding extremal graphs. Thereafter, we prove lower bounds on the ratios $GA/\sqrt{\delta}$ and GA/δ in terms of the order n .

To prove our first bound, namely an upper bound on GA/δ , we need the following preliminary result.

Lemma 2.1 For $n \geq 3$,

$$\frac{(n-2)(n-3)}{2} + \frac{2(n-2)\sqrt{(n-1)(n-2)}}{2n-3} + \frac{2\sqrt{n-1}}{n} > \frac{(n-1)(n-2)}{2}.$$

Proof. Considering the ratio r_n of the left hand side to the right hand side of the inequality we have

$$\begin{aligned} r_n &= \left(\frac{(n-2)(n-3)}{2} + \frac{2(n-2)\sqrt{(n-1)(n-2)}}{2n-3} + \frac{2\sqrt{n-1}}{n} \right) \cdot \left(\frac{2}{(n-1)(n-2)} \right) \\ &= \frac{n-3}{n-1} + \frac{4}{2n-3} \sqrt{\frac{n-2}{n-1}} + \frac{4}{n(n-2)\sqrt{n-1}} \\ &= 1 - \frac{2}{n-1} + \frac{4\sqrt{n-2}}{(2n-3)\sqrt{n-1}} + \frac{4}{n(n-2)\sqrt{n-1}} \\ &= 1 + \frac{2 \left(2\sqrt{(n-2)(n-1)} - (2n-3) \right)}{(2n-3)(n-1)} + \frac{4}{n(n-2)\sqrt{n-1}} \\ &= 1 + \frac{2 \left(4(n-2)(n-1) - (2n-3)^2 \right)}{(2n-3)(n-1) \left(2\sqrt{(n-2)(n-1)} + (2n-3) \right)} + \frac{4}{n(n-2)\sqrt{n-1}} \\ &= 1 - \frac{2}{(2n-3)(n-1) \left(2\sqrt{(n-2)(n-1)} + (2n-3) \right)} + \frac{4}{n(n-2)\sqrt{n-1}} > 1. \end{aligned}$$

This shows the inequality. □

Theorem 2.2 For any connected graph on $n \geq 3$ with minimum degree δ and geometric-arithmetic index GA ,

$$\frac{GA}{\delta} \leq \frac{(n-2)(n-3)}{2} + \frac{2(n-2)\sqrt{(n-1)(n-2)}}{2n-3} + \frac{2\sqrt{n-1}}{n}$$

with equality if and only if G is the kite $Ki_{n,n-1}$.

Proof. If $\delta = 1$, then the maximum number of edges in G is $(n-1)(n-2)/2 + 1$ which is attained if and only if G is the kite $Ki_{n,n-1}$. In this case equality holds.

If $\delta = 1$ and the number of edges is not maximum, *i.e.*, $m \leq (n-1)(n-2)/2$, then using Lemma 2.1 and the fact that $GA \leq m$, we have

$$GA \leq \frac{(n-1)(n-2)}{2} < \frac{(n-2)(n-3)}{2} + \frac{2(n-2)\sqrt{(n-1)(n-2)}}{2n-3} + \frac{2\sqrt{n-1}}{n}.$$

Therefore, in this case, the inequality is strict.

If $\delta \geq 2$, then

$$\frac{GA}{\delta} \leq \frac{m}{2} \leq \frac{n(n-1)}{4} < \frac{(n-2)(n-3)}{2} + \frac{2(n-2)\sqrt{(n-1)(n-2)}}{2n-3} + \frac{2\sqrt{n-1}}{n}.$$

Therefore, in this case also, the inequality is strict. \square

The following theorem is proved in [14].

Theorem 2.3 ([14]) *Let G be a graph on n vertices with minimum degree δ such that $\delta \geq k \geq 2$, for some integer k .*

1. If $n \leq 10$, then

$$GA \geq \frac{nk}{2}.$$

2. If $n \geq 11$, then

$$GA \geq \min \left\{ \frac{nk}{2}, \frac{(k+1)\sqrt{k}(n-1)^{\frac{3}{2}}}{n-1+k} \right\}.$$

We are going to make a more precise statement of the above theorem. For that purpose, we need the following lemma.

Lemma 2.4 *For an integer $k \geq 4$, let*

$$f_k(t) = \frac{(k+1)\sqrt{k}(t-1)^{\frac{3}{2}}}{t-1+k} - \frac{kt}{2} \quad \text{for } t \geq k+1.$$

We have

$$f_k(t) \begin{cases} \geq 0 & \text{if } k+1 \leq t \leq k+8; \\ < 0 & \text{if } t \geq k+9. \end{cases}$$

Proof. To do the calculations related to the function in this lemma, we used the software *WolframAlpha* (available at wolframalpha.com).

The second derivative of f_k is

$$f_k''(t) = -\frac{(k+1)\sqrt{k}(t^3 + (7k-3)t^2 + (3k^2 - 14k + 3)t - 3k^3 - 3k^2 + 7k - 1)}{4(t-1+k)^4\sqrt{t-1}}$$

which is negative for all $t \geq k+1$ with $k \geq 4$. Thus the graph of f_k is concave down for all $t \geq k+1$ with $k \geq 4$. Therefore, the equation $f_k(t) = 0$ has at most 2 solutions.

We have $f_k(k+1) = 0$, that is $t_1 = k+1$ is a solution of $f_k(t) = 0$. Also

$$f_k(k+8) = \frac{(k+1)\sqrt{k}(k+7)^{\frac{3}{2}}}{2k+7} - \frac{k(k+8)}{2} > 0 \quad \text{for all } k \geq 4$$

and

$$f_k(k+9) = \frac{(k+1)\sqrt{k}(k+8)^{\frac{3}{2}}}{2k+8} - \frac{k(k+9)}{2} < 0 \quad \text{for all } k \geq 4,$$

that is $f_k(t) = 0$ has a solution t_2 satisfying $k+8 < t_2 < k+9$. \square

In the statement of the above lemma, there is the condition that $k \geq 4$. For $k = 2$ and $k = 3$, the conclusions are slightly different, namely

$$f_2(t) = \frac{3\sqrt{2}(t-1)^{\frac{3}{2}}}{t+1} - t \begin{cases} \geq 0 & \text{if } 3 \leq t \leq 11; \\ < 0 & \text{if } t \geq 12; \end{cases}$$

and

$$f_3(t) = \frac{4\sqrt{3}(t-1)^{\frac{3}{2}}}{t+2} - \frac{3t}{2} \begin{cases} \geq 0 & \text{if } 3 \leq t \leq 12; \\ < 0 & \text{if } t \geq 13. \end{cases}$$

In view of the above lemma and discussion, we can state Theorem 2.3 as follows.

Theorem 2.3' *Let G be a graph on n vertices with minimum degree δ such that $\delta \geq 2$.*

1. *If $\delta = 2$, then*

$$GA \geq \begin{cases} n & \text{if } 3 \leq n \leq 11; \\ \frac{3\sqrt{2}(n-1)^{\frac{3}{2}}}{n+1} & \text{if } n \geq 12. \end{cases}$$

2. *If $\delta = 3$, then*

$$GA \geq \begin{cases} \frac{3n}{2} & \text{if } 4 \leq n \leq 12; \\ \frac{4\sqrt{3}(n-1)^{\frac{3}{2}}}{n+2} & \text{if } n \geq 13. \end{cases}$$

3. *If $\delta \geq 4$, then*

$$GA \geq \begin{cases} \frac{\delta n}{2} & \text{if } \delta + 1 \leq n \leq \delta + 8; \\ \frac{(\delta+1)\sqrt{\delta}(n-1)^{\frac{3}{2}}}{n+\delta-1} & \text{if } n \geq \delta + 9. \end{cases}$$

Note that the cases $\delta = 2$ and $\delta = 3$ in the above theorem were already stated in [14].

We next prove a lower bound on the ratio $GA/\sqrt{\delta}$ and characterize the corresponding extremal graphs.

Proposition 2.5 *For any connected graph on $n \geq 3$ vertices with minimum degree δ and geometric-arithmetic index GA ,*

$$\frac{GA}{\sqrt{\delta}} \geq \frac{2(n-1)^{\frac{3}{2}}}{n}$$

with equality if and only if G is the star S_n .

Proof. If $\delta = 1$, the result is proved in [17].

Assume that $\delta \geq 2$. From Theorem 2.3, we have

$$\frac{GA}{\sqrt{\delta}} \geq \min \left\{ \frac{n\sqrt{\delta}}{2}, \frac{(\delta+1)(n-1)^{\frac{3}{2}}}{(n-1+\delta)} \right\}.$$

First, we compare between

$$\frac{n\sqrt{\delta}}{2} \quad \text{and} \quad \frac{2(n-1)^{\frac{3}{2}}}{n}.$$

We have

$$\frac{n\sqrt{\delta}}{2} - \frac{2(n-1)^{\frac{3}{2}}}{n} \geq \frac{n}{\sqrt{2}} - \frac{2(n-1)^{\frac{3}{2}}}{n} = \frac{n^2 - 2(n-1)^{\frac{3}{2}}}{\sqrt{2}n}.$$

The study of the function

$$f(t) = \frac{t^2 - 2(t-1)^{\frac{3}{2}}}{\sqrt{2}t}$$

shows that $f(t) > 0$ for all $t \geq 1$. Therefore

$$\frac{n\sqrt{\delta}}{2} > \frac{2(n-1)^{\frac{3}{2}}}{n}$$

for all $n \geq \delta + 1 \geq 3$.

Now, we compare between

$$\frac{(\delta + 1)(n - 1)^{\frac{3}{2}}}{(n - 1 + \delta)} \quad \text{and} \quad \frac{2(n - 1)^{\frac{3}{2}}}{n},$$

or equivalently between

$$\frac{\delta + 1}{(n - 1 + \delta)} \quad \text{and} \quad \frac{2}{n}.$$

We have

$$\begin{aligned} \frac{\delta + 1}{(n - 1 + \delta)} - \frac{2}{n} &= \frac{(\delta + 1)n - 2(n - 1 + \delta)}{n} = \frac{(\delta - 1)n - 2(\delta - 1)}{n} \\ &= \frac{(\delta - 1)(n - 2)}{n} > 0 \quad \text{for all } n \geq \delta + 1 \geq 3. \end{aligned}$$

Thus

$$\frac{\delta + 1}{(n - 1 + \delta)} > \frac{2}{n}$$

for all $n \geq \delta + 1 \geq 3$.

In conclusion

$$\min \left\{ \frac{n\sqrt{\delta}}{2}, \frac{(\delta + 1)(n - 1)^{\frac{3}{2}}}{(n - 1 + \delta)} \right\} \geq \frac{2(n - 1)^{\frac{3}{2}}}{n}$$

for all $n \geq \delta + 1 \geq 3$. This completes the proof. \square

Under certain conditions, the inequality in the above proposition remains valid if $GA/\sqrt{\delta}$ is replaced by GA/δ , as next stated.

Proposition 2.6 *If G is a connected graph on n vertices with minimum degree δ such that $n \geq \max \{13, 2\sqrt{\delta}(\sqrt{\delta} + 1)/(\sqrt{\delta} - 1)\}$, then*

$$\frac{GA}{\delta} \geq \frac{2(n - 1)^{\frac{3}{2}}}{n}$$

with equality if and only if G is the star S_n .

Proof. Again if $\delta = 1$, the result is proved in [17].

Assume that $\delta \geq 2$. In this case, from Theorem 2.3, we have

$$\frac{GA}{\delta} \geq \min \left\{ \frac{n}{2}, \frac{(\delta + 1)(n - 1)^{\frac{3}{2}}}{(n - 1 + \delta)\sqrt{\delta}} \right\}.$$

First, we compare between

$$\frac{n}{2} \quad \text{and} \quad \frac{2(n - 1)^{\frac{3}{2}}}{n}.$$

For $t \geq 2$, consider the function

$$f(t) = \frac{t}{2} - \frac{2(t - 1)^{\frac{3}{2}}}{t}.$$

This function is continuous and its second derivative is

$$f''(t) = \frac{t(t + 4) - 8}{2t^3\sqrt{t - 1}}.$$

We have $f''(t) > 0$ for all $t \geq 2$ and therefore the function $f(t)$ is concave up for $t \geq 2$. Thus the equation $f(t) = 0$ has (at most) two solutions: one of them is $t_1 = 2$ and the other one, say t_2 , satisfies $12 < t_2 < 13$ (since $f(12) < 0$ and $f(13) > 0$). We conclude that $f(t) > 0$, which means

$$\frac{n}{2} > \frac{2(n-1)^{\frac{3}{2}}}{n} \quad \text{for all } n \geq 13.$$

Now, we compare between

$$\frac{(\delta+1)(n-1)^{\frac{3}{2}}}{(n-1+\delta)\sqrt{\delta}} \quad \text{and} \quad \frac{2(n-1)^{\frac{3}{2}}}{n},$$

or equivalently between

$$\frac{\delta+1}{(n-1+\delta)\sqrt{\delta}} \quad \text{and} \quad \frac{2}{n}.$$

We have

$$\begin{aligned} \frac{\delta+1}{(n-1+\delta)\sqrt{\delta}} - \frac{2}{n} &= \frac{(\delta+1)n - 2(n-1+\delta)\sqrt{\delta}}{n(n-1+\delta)\sqrt{\delta}} \\ &= \frac{(\delta - 2\sqrt{\delta} + 1)n - 2(\delta-1)\sqrt{\delta}}{n(n-1+\delta)\sqrt{\delta}} \\ &= \frac{(\sqrt{\delta}-1)^2 n - 2(\sqrt{\delta}-1)(\sqrt{\delta}+1)\sqrt{\delta}}{n(n-1+\delta)\sqrt{\delta}} \\ &= \frac{\sqrt{\delta}-1}{n(n-1+\delta)\sqrt{\delta}} \left((\sqrt{\delta}-1)n - 2(\sqrt{\delta}+1)\sqrt{\delta} \right). \end{aligned}$$

Thus

$$\frac{\delta+1}{(n-1+\delta)\sqrt{\delta}} > \frac{2}{n} \quad \text{if and only if} \quad n > \frac{2\sqrt{\delta}(\sqrt{\delta}+1)}{\sqrt{\delta}-1}.$$

The combination of both cases completes the proof. \square

References

- [1] A. Ali, A. A. Bhatti, and Z. Raza, Further inequalities between vertex-degree-based topological indices. *Int. J. Appl. Comput. Math.* 3 (2017) 1921–1930.
- [2] K. C. Das, I. Gutman, B. Furtula, On the first geometric-arithmetic index of graphs. *Discrete Applied Mathematics*, 159 (2011) 2030–2037.
- [3] K.C. Das, I. Gutman, B. Furtula, Survey on geometric-arithmetic indices of graphs. *MATCH Commun. Math. Comput. Chem.* 65 (2011) 595–644.
- [4] T. Divnić, M. Milivojević and Lj. Pavlović, Extremal graphs for the geometric-arithmetic index with given minimum degree. *Discrete Applied Mathematics* 162 (2014) 386–390.
- [5] B. Furtula, I. Gutman, Geometric-arithmetic Indices. In: I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors—Theory and Applications*, vol. I, Univ. Kragujevac, Kragujevac, 2010, pp. 137–172.
- [6] I. Gutman, B. Furtula (Eds.), *Recent Results in the Theory of Randić Index*. Univ. Kragujevac, Kragujevac, 2008.
- [7] I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors - Theory and Applications I*, Univ. Kragujevac, Kragujevac, 2010.
- [8] I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors - Theory and Applications II*, Univ. Kragujevac, Kragujevac, 2010.
- [9] L. B. Keir, L. H. Hall, *Molecular connectivity in chemistry and drug research*. Academic Press, New York 1976.
- [10] L. B. Keir, L. H. Hall, *Molecular connectivity in structural-activity analysis*. Research study, Letchworth, England, 1986.

-
- [11] X. Li, I. Gutman, *Mathematical Aspects of Randić Type Molecular Structure Descriptors*, Univ. Kragujevac, Kragujevac, 2006.
- [12] X. Li, Y. Shi, A survey on the Randić index, *MATCH Commun. Math. Comput. Chem.* 59 (2008) 127–156.
- [13] M. Randić, On characterization of molecular branching. *J. Am. Chem. Soc.* 97 (1975) 6609–6615.
- [14] J. M. Rodríguez and J. M. Sigarreta, On the Geometric–Arithmetic Index. *MATCH Commun. Math. Comput. Chem.* 74 (2015) 103–120
- [15] M. Sohrabi-Haghighat and M. Rostami, Using linear programming to find the extremal graphs with minimum degree 1 with respect to Geometric-Arithmetic index. *Applied mathematics in Engineering, Management and Technology* 3 (2015) 534–539.
- [16] R. Todeschini, V. Consonni, *Molecular Descriptors for Chemoinformatics*, Wiley-VCH, Weinheim, 2009.
- [17] D. Vukičević and B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end vertex degrees of edges. *Journal of Mathematical Chemistry* 46 (2009) 1369–1376.
- [18] H. Wiener, Structural determination of paraffin boiling points, *J. Am. Chem. Soc.* 69 (1947) 17–20.
- [19] Y. Yuan, B. Zhou, and N. Trinajstić, On geometric-arithmetic index. *Journal of Mathematical Chemistry* 47 (2010) 833–841.